

# ***Interactive comment on “Assessing the role of planetary and gravity waves on the vertical structure of ozone over central Europe” by Peter Križan***

**Anonymous Referee #2**

Received and published: 31 January 2019

Based on the Teitelbaum method, this manuscript studies the characteristics of ozone lamina under the influence of planetary and gravity waves. This article seems to have done a lot of work. Even though I'm not an expert in this area (ozone lamina), there are a few things that make me confused.

Major comments: 1. The formation mechanism of ozone lamina. Tomikawa et al. (2002) reported that the formation of the ozone laminae is closely related to the vertical shear of the subtropical jet. I strongly suggest the authors to discuss in the introduction about the formation mechanisms of the ozone lamina and in which the role of wave activities play.

[Printer-friendly version](#)

[Discussion paper](#)



Tomikawa, Y. , Sato, K. , Kita, K. , Fujiwara, M. , Yamamori, M. , & Sano, T. . (2002). Formation of an ozone lamina due to differential advection revealed by intensive observations. *J. Geophys. Res.*, 107(D10).

2. Lines 101-108, need some references (at least one) or make some explanation: e.g. Why choose 5-15km and 17-22km height area to distinguish. If the identification process is proposed by the author, it appears from the description that the author only uses ozone thin layers at different height region to define whether the ozone laminae is caused by gravity or planetary waves. This makes it very puzzling because gravity waves almost exist anywhere in the earth's atmosphere.

3. The authors have only mentioned the thin layer of ozone caused by gravitational and planetary waves, but I think that some other meso-and small-scale atmospheric processes (such as strong convection, tropopause folding, strong wind shear, stratospheric streamers, etc.) may also responsible for the formation of ozone laminae.

4. Gravity and planetary waves run through the title and the paper, but there is no evidence of their existence in the manuscript (even though the authors indicate that the ozone profile can be used to detect fluctuations)

5. Reading the manuscript, I still didn't understand how gravity and planetary waves affect and lead to ozone laminae. Personally, a detailed case is necessary.

Minor comments: 1. Lines 44-46, as you mentioned, it is the large lamina that has a close correlation with the total ozone content, not the narrow lamina. The actual significance of narrow lamina still not clear throughout the manuscript. 2. Lines 47-48, needs relevant references (at least one), especially about the influence of waves on the laminae. 3. Line 75 from->on ? 4. Line 76 for the approximating-> for approximating 5. Line 125 partitioning of laminae-> partitioning laminae ? 6. Conclusion: as mentioned in the introduction, if the Teitelbaum method is suitable for central Europe? And how well?

[Printer-friendly version](#)

[Discussion paper](#)



---

Interactive  
comment

[Printer-friendly version](#)

[Discussion paper](#)

