



1 **Unknown high-frequency (7-12 kHz) quasi-periodic VLF emissions observed on**  
2 **the ground at L ~ 5.5**

3 Jyrki Manninen<sup>1</sup>, Natalia Kleimenova<sup>2,3</sup>, Tauno Turunen<sup>1</sup>, and Liudmila Gromova<sup>4</sup>

4  
5 <sup>1</sup>Sodankylä Geophysical Observatory, Sodankylä, Finland

6 <sup>2</sup>Schmidt Institute of Physics of the Earth RAN, Moscow, Russia

7 <sup>3</sup>Space Research Institute RAN, Moscow, Russia

8 <sup>4</sup>Pushkov Institute of Terrestrial Magnetism, Ionosphere and Radio Wave Propagation RAN,  
9 Moscow, Troitsk, Russia

10

11 *Correspondence to:* J. Manninen ([jyrki.manninen@sgo.fi](mailto:jyrki.manninen@sgo.fi))

12 **Abstract.** We reveal previously unknown quasi-periodic (QP) VLF emissions at the unusual  
13 high-frequency band of ~7-11 kHz by applying the digital filtering of strong sferics to the ground-  
14 based VLF data recorded at Kannuslehto station (KAN). It is located in Northern Finland at L~5.5.  
15 The frequencies of QP emissions are much higher than the equatorial electron gyrofrequency at L~  
16 5.5. Thus, these emissions must have been generated at much lower L-shells than KAN. Two high-  
17 frequency QP emission events have been studied in detail. The emissions were right-hand  
18 polarized waves indicating an overhead location of the exit area of waves in the ionosphere. In one  
19 event, the spectral-temporal forms of the emissions looked like a series of giant “bullets” with the  
20 very abrupt cessation. Unfortunately, we could not explain such strange shape of the waves. In the  
21 second event, the modulation period was about 3 min under the absence of the simultaneous  
22 geomagnetic pulsations. The studied emissions lasted about 4 hours and were observed under the  
23 very quiet geomagnetic activity. The adequate mechanisms of the generation and propagation of  
24 the revealed high-frequency QP emissions have not yet been established. We speculate that studied  
25 QP emissions can be attributed to the auto-oscillations of the cyclotron instability in the  
26 magnetospheric plasma maser.

27 **Keywords.** Magnetospheric physics (magnetosphere-ionosphere interactions; plasma waves and  
28 instabilities, energetic particles)

29



30

## 31 **1 Introduction**

32 The whistler mode VLF emissions, observed on the ground and onboard satellites, can exhibit a  
33 quasi-periodic (QP) occurrence of the separated signal patches in the frequency range of about  
34 0.5-4 kHz, and due to that, they are termed QP-emissions (e.g. Carson et al., 1965; Kitamura et al.,  
35 1969; Sato et al., 1974; Morrison et al., 1994; Sazhin and Hayakawa, 1994; Engebretson et al.,  
36 2004; Němec et al., 2016). The QP emissions were measured by satellites over a large spatial area  
37 (e.g., Němec et al., 2013, 2014; Titova et al., 2015; Hayosh et al., 2016). A series of expressive  
38 long-lasting QP emissions, observed at ground station at the auroral latitudes, have been shown by  
39 Manninen et al., (2012, 2013, 2014). Note, the ground-based recordings provide important  
40 information about the temporal properties of the studied waves.

41 However, even at the auroral latitudes, the ground-based VLF measurements are mostly  
42 covered by strong atmospheric (sferics) (e.g., Yamashita, 1978; Yedemsky et al., 1992; Volland,  
43 1995) which hide all natural emissions with weaker amplitudes. Weak atmospheric are  
44 electromagnetic pulses with duration of  $\sim 10$ – $100$  milliseconds originated from lightning  
45 discharges and propagating over distances of a few thousands km in the Earth-ionosphere  
46 waveguide (e.g., Ohya et al., 2015). To study the waves of the magnetospheric origin, we applied  
47 the special method of digitally filtering out the strong impulsive atmospheric with duration less  
48 than 30 ms. This processing allowed us to discover completely new types of high-frequency VLF  
49 emissions (Manninen et al., 2016). Beside the separated discrete emissions, discussed by  
50 Manninen et al. (2016), we found also some previously unknown unusual high-frequency quasi-  
51 periodic VLF-emission lasting up to several hours. Such emissions have not been reported earlier.

52 The aim of this paper is to present the spectral feature of the revealed emissions at the  
53 frequency higher than 7 kHz. Our study was based on the VLF measurements (Manninen, 2005) at  
54 the auroral latitudes ( $L \sim 5.5$ ) in Northern Finland at Kannuslehto (KAN, the geographic  
55 coordinates are  $67.74^\circ\text{N}$ ,  $26.27^\circ\text{E}$ ). To reject the influence of the strong narrow-band navigation  
56 transmitter signals, we analyzed the VLF emissions at the frequencies less than 12 kHz.

## 57 **2 Observation results and its discussion**

58 Here we present the analysis of two events of the unusual high-frequency QP emissions  
59 observed on 25 December 2011 and 5 January 2014 both at 16-20 UT (19-23 MLT). Two-hour  
60 non-filtered spectrograms, shown in the upper panels of Figure 1, demonstrate that the strong  
61 sferics hid all other weaker waves at the frequencies higher than 5-6 kHz. The same



62 spectrograms, obtained after the filtering out sferics, are shown in the bottom panels of Fig. 1.  
63 Unexpectedly, after this filtering, the strange QP emissions were detected at the frequency band of  
64 about 7 - 11 kHz. The rather similar event was recorded also on 10 December 2012 (not studied  
65 here). All considered VLF emissions were the right-hand polarized waves indicating the  
66 ionospheric exit point location to be approximately overhead. It is unlike the sferics which  
67 characterized by the left-hand polarization (e.g., Yedemsky et al., 1992; Haykawa et al., 1994).  
68 Some behavior of waves such as the temporal dynamics of the right-hand polarized part of the QP  
69 missions and its arriving direction of both studied events are given in Figure 2 in frequency band  
70 of 6-12 kHz.

71 Description of the VLF receiver is given by Manninen (2005). Sampling frequency was 78125  
72 Hz and the noise level of the receiver is  $10^{-4}$  nT<sup>2</sup>/Hz. The FFT size is 8192 in all spectral analysis.

### 73 ***2.1 Event of 25 Dec 2011***

74 The spectrogram of the first studied event (25 Dec 2011, the left panels in Fig. 1) demonstrates  
75 the strange spectral-temporal forms look like a chain of giant “bullets” with a very sharp end of  
76 each element. The 4-hours total power spectrogram of this event as well as the right-hand (RH)  
77 polarized power of the waves in frequency band of 6-12 kHz and the angles of arrival of the waves  
78 are given in Fig. 2a. It is seen that the RH polarized waves represented two separated frequency  
79 bands: the strongly modulated band of 7.0-8.5 kHz and almost continuous hiss-like band of 9.0-  
80 10.5 kHz. Two strong “bullets” are observed in the beginning of the high-frequency band. It is  
81 interesting to note that the first “bullet” occurred at the higher frequency band, the second “bullet”  
82 appeared at both frequency bands and later the “bullets” were observed only at the lower  
83 frequencies.

84 The direction-finding analysis showed that the high-frequency band arrived mostly along the  
85 meridian during the whole event, but the angles of arrival of the low frequency “bullets” changed  
86 with time deviating from the meridian (Fig. 2a, two lower panels). The angle of arrival has an  
87 uncertainty of 180 degrees (or ambiguity) due to only two orthogonal magnetic loop antennas  
88 oriented in the north-south and east-west directions are used in this study. The angle of arrival is  
89 determined as the direction of the minor axes of the wave polarization ellipses. In order to remove  
90 the ambiguity, an additional vertical electric antenna would be needed.

91 Attributing the emission generation to the electron cyclotron instability (e.g. Trakhtengerts and  
92 Rycroft, 2008), we may suppose the different location of the generation regions responsible for the  
93 high-frequency hiss and for the “bullets”. The higher frequency band was originated at the smaller  
94 L-shells than the lower frequency band, and, the source of the “bullets”-like emissions removed in



95 course of time. Although these two bands may have different source location, the sharp ends of  
96 every burst occur simultaneously in all frequencies.

97 The spectrogram of one separated “bullet” is presented in Fig. 3 both in 20 min (Fig. 3a) and 2  
98 min (Fig. 3b) time scales. The very abrupt cessation of the “bullet” which is clearly seen in Fig. 3a  
99 near 17:29 UT is really steep in the several seconds scale (Fig. 3b). It is important to note that  
100 there were no such remarkable changes of the higher frequency (9.0-10.5 kHz) band.  
101 Unfortunately, we could not find a plausible source of such strong signal rejection.

## 102 *2.2. Event of 5 Jan 2014*

103 The second studied event (5 January 2014, right panels in Fig. 1) represents a series of quasi-  
104 repeated short patches with rather constant low frequencies and gradually increasing upper ones.  
105 The analysis of this QP event (Fig. 2b) demonstrated that the behavior of the event is much more  
106 visible in RH wave power spectrum than in its total power spectrum (upper plot). During first two  
107 hours, the band of emissions gradually shifted to the higher frequencies. At the beginning, the  
108 upper frequency limit increased from ~ 9 kHz to ~ 11 kHz within 1.5 hours. After ~18:30 UT, it  
109 began to decrease down to ~ 10 kHz (at ~19:30 UT). As well as during the 25 Dec 2011 QP event  
110 (Fig. 2a), the angle of arrival of this QP event quickly changed deviating from the north-south  
111 direction. At about 17:30 UT, the angle of arrival got settled approximately along the east-west  
112 direction as if in a daily rotation of the Earth the receiver is removed away from the source of  
113 wave generation. Note, that at this time, the waves remained RH polarized, so, we may conclude  
114 that all this time, the VLF receiver was located in the vicinity of the ionospheric exit area of  
115 waves.

116 The detailed structure of the quasi-repeated patches is shown in Figure 4 as the temporal  
117 variation of total (T) and right-handed (RH) wave power spectra. The bottom panel represents the  
118 variations of the integrated wave intensity in the frequency band of 7-10 kHz demonstrating the  
119 average period of the VLF wave repetition of ~ 3 min. In addition, there were altogether 10 QP  
120 emission bursts with 3 s internal periodicity shown in the bottom panel in Figure 4. According to  
121 ground-based measurements at Scandinavian IMAGE magnetometer chain, there were no  
122 geomagnetic pulsations with similar periods.

123 Besides this 3-min periodicity, Figure 4 demonstrates an appearance of the strong impulsive  
124 packets of very short periods VLF emissions superposing the quasi-repeated VLF patches. An  
125 example of the fine structure of these impulses is shown in Figure 5 at 2-min time scale. It was  
126 found that its structure consists of the long series of the RH polarized QP emissions in the ~ 8.5-  
127 10.5 kHz frequency band with repetition period of about 3 s. The first six emissions were non-  
128 dispersive, but the following ones exhibited smaller intensity decreasing with time and positive



129 time-frequency slope ( $df/dt > 0$ ). The short period QP emissions having similar periodicity have  
130 been recorded previously on the ground and onboard satellites (e.g., Bessalov et al., 2010;  
131 Manninen et al., 2014). However, in these papers the QP emissions were at the frequencies less  
132 than 2.5 kHz, but in our case the QP emissions were observed at frequency higher than 7 kHz.  
133

134 During our events, the conditions in the solar wind and IMF were very quiet shown in Figure  
135 6a. During the studied intervals, there were no geomagnetic disturbances and pulsations at SOD  
136 located 45 km south-east from KAN. The planetary geomagnetic activity was quiet with  $K_p \sim 0$ .  
137 However, some small  $K_p$  enhancements can be seen within preceding 24 hours (Fig. 6b),  
138 providing additional electrons in the radiation belts. This might be due to a small magnetic  
139 substorm in the night side (AL values up to 100 nT) before the 5 Jan 2014 event. There was no  
140 substorm before 25 Dec 2011 event. According to data of RBSP (A and B) satellites  
141 [<http://enarc.space.swri.edu/PTP>], KAN was mapped in the vicinity of the plasmopause (Fig. 6c).

142 There is no complete theory adequately explained the generation of such high-frequency QP  
143 emissions because our finding is so new. Anyway, we attribute the studied QP emissions to the  
144 auto-oscillations of the cyclotron instability of the Earth radiation belts in the magnetospheric  
145 plasma maser (Bessalov and Trakhtengerts, 1986, Trakhtengerts and Rycroft, 2008). The  
146 excitation of these oscillations is possible only during low geomagnetic activity as it was observed  
147 during the studied events.

148 The revealed 7-10 kHz QP emissions could not be generated at L-shell corresponding to KAN  
149 location because the equatorial electron gyrofrequency at  $L \sim 5.5$  is  $\sim 5$  kHz, and this value  
150 controls the upper frequency of the generation of whistler mode waves at given L shell. Moreover,  
151 the ducted propagation of the whistler mode wave in the magnetosphere is possible only at  
152 frequencies less than one half of this value (Carpenter, 1968). So, we suppose that the studied QP  
153 emissions propagated in unducted mode like it was discussed in the papers (Němec et al., 2013;  
154 Titova et al., 2015; Hayosh et al., 2016). We assume that the high-frequency QP emissions are  
155 generated at much lower L shells than KAN, probably even inside of the plasmasphere.

### 156 3 Summary

157 Applying the digital filtering of strong sferics to the ground-based VLF data at  $L \sim 5.5$  we reveal  
158 previously unknown quasi-periodic VLF emissions at unusual high-frequencies of  $\sim 7$ -11 kHz.  
159 These frequencies are much higher than the equatorial gyrofrequency of electrons at  $L \sim 5.5$ .

160 Contrary to typical QP emissions, the discovered high-frequency waves, presented in this paper,  
161 were observed in the late evening. The emissions were right-hand polarized indicating that the  
162 ionospheric exit point of the waves was located approximately overhead. In the first event, the



163 spectral-temporal forms of the emissions looked like a series of giant “bullets” with the very  
164 abrupt cessation. There were no any geomagnetic nor absorption events, which could explain the  
165 sharp stop of generation of the “bullets” on 25 Dec 2011. Maximum variation in geomagnetic field  
166 was only 4 nT and riometer absorption was at the background level. Unfortunately, we could not  
167 explain such strange feature of the emissions. In the second event, the repetition period was about  
168 3 min with the absence of the simultaneous geomagnetic pulsations. Thus, the source of such  
169 modulation remains unknown.

170 The high-frequency QP emissions were observed under very quiet geomagnetic activity.  
171 Apparently, the studied emissions can be attributed to the auto-oscillations of the cyclotron  
172 instability in the magnetospheric plasma maser (Bespalov and Trakhtengerts, 1986; Trakhtengerts  
173 and Rycroft, 2008), and we speculate that the studied waves propagated in the unducted mode.  
174 The source of studied QP emissions could be located at low L-values into the plasmasphere.  
175 However, the adequate mechanism of the generation and propagation of QP emissions at untypical  
176 high frequencies is not yet established.

177

#### 178 *Acknowledgements*

179 The research was supported by the Academy of Finland (grant no. 287988) during stay N.K.  
180 and L.G. in Sodankylä. The work by N.K. was partly supported by the Program of Presidium of  
181 Russian Academy of Science No. 28.

182

#### 183 **References**

- 184 Bespalov, P.A. and Trakhtengerts, V.Yu.: The cyclotron instability in the Earth radiation belts, Rev.  
185 Plasma Physics. Plenum Publ. N.Y, ed. Leontovich, M.A., 10, 155-292, 1986.
- 186 Bespalov, P.A., Parrot, M. and Manninen, J.: Short period VLF emissions as solitary envelope  
187 waves in a magnetospheric plasma maser, J. Atmos. Solar Terr. Phys., 72, 1275–1281, 2010.
- 188 Carpenter, D.L.: Ducted whistler-mode propagation in the magnetosphere; a half-  
189 gyrofrequency upper intensity cut-off and some associated wave growth phenomena, J.  
190 Geophys. Res., 73, 2919–2928. 1968.
- 191 Carson, W.B., Koch, J.A., Pope, J.H., and Gallet, R.M.: Long-period very low frequency emission  
192 pulsations, J. Geophys. Res., 70 (17), 4293–4303, 1965.
- 193 Engebretson, M.J., Posch, J.L., Halford, A.J., Shelburne, G.A., Smith, A.J., Spasojevic, M., Inan,  
194 U.S., and Arnoldy, R.L.: Latitudinal and seasonal variations of quasiperiodic and periodic VLF



- 195 emissions in the outer magnetosphere, *J. Geophys. Res.*, 109, A05216,  
196 doi:10.1029/2003JA010335, 2004.
- 197 Hayakawa, M., Ohta, K., and Baba, K.: Wave characteristics of tweek atmospherics deduced from  
198 the direction-finding measurement and theoretical interpretation, *J. Geophys. Res.*, 99 (05),  
199 10733-10743, 1994.
- 200 Hayosh, M., N̄emec, F., Santolik, O., and Parrot, M.: Propagation properties of quasi-periodic  
201 VLF emissions observed by the DEMETER spacecraft, *J. Geophys. Res. Space Physics*, 43,  
202 1007–1014, doi:10.1002/2015GL067373, 2016.
- 203 Kitamura, T., Jacobs, J.A., Watanabe, T., and Flint, J.R.B.: An investigation of quasi-periodic VLF  
204 emissions, *J. Geophys. Res.*, 74(24), 5652–5664, 1969.
- 205 Manninen, J.: Some aspects of ELF–VLF emissions in geophysical research, Sodankylä Geophys.  
206 Obs. Publ. no. 98, Oulu Univ., Finland, 2005.  
207 <http://www.sgo.fi/Publications/SGO/thesis/ManninenJyrki.pdf>.
- 208 Manninen, J., Kleimenova, N.G., and Kozyreva, O.V.: New type of ensemble of quasi-periodic,  
209 long-lasting VLF emissions at the auroral zone, *Ann. Geophys.*, 30, 1655–1660, 2012.
- 210 Manninen J., Kleimenova, N.G., Kozyreva, O.V., Bepalov, P.A., and Kozlovsky, A.E.: Non-  
211 typical ground-based quasi-periodic VLF emissions observed at  $L \sim 5.3$  under quiet  
212 geomagnetic conditions at night, *J. Atmos. Solar-Terr. Phys.*, 99, 123-128, 2013.
- 213 Manninen, J., Demekhov, A.G., Titova, E.E., Kozlovsky, A.E., and Pasmanik, D.L.: Quasi-  
214 periodic VLF emissions with short-period modulation and their relationship to whistlers: A case  
215 study, *J. Geophys. Res. Space Physics*, 119, 3544–3557, doi:10.1002/2013JA019743, 2014.
- 216 Manninen, J., Turunen, T., Kleimenova, N., Rycroft, M., Gromova, L., and Sirvio, I.: Unusually  
217 high frequency natural VLF radio emissions observed during daytime in Northern Finland,  
218 *Environ. Res. Lett.* 11, 124006, doi:10.1088/1748-9326/11/12/124006, 2016.
- 219 Morrison, K., Engebretson, M. J., Beck, J. R., Johnson, J. E., Arnoldy, R. L., Cahill Jr, L. J.,  
220 Carpenter, D. L., and Gallani, M.: A study of quasi-periodic ELF–VLF emissions at three  
221 Antarctic stations: evidence for off-equatorial generation? *Ann. Geophys.*, 12, 139–146,  
222 doi:10.1007/s00585-994-0139-8, 1994.
- 223 N̄emec, F., Santolik, O., Parrot, M., Pickett, J.S., Hayosh, M., and Cornilleau-Wehrin, N.:  
224 Conjugate observations of quasi-periodic emissions by Cluster and DEMETER spacecraft, *J.*  
225 *Geophys. Res. Space Physics*, 118, 198–208, doi:10.1029/2012JA018380, 2013.
- 226 N̄emec, F., Pickett, J.S., and Santolik, O.: Multispacecraft Cluster observations of quasiperiodic  
227 emissions close to the geomagnetic equator, *J. Geophys. Res. Space Physics*, 119, 9101–9112,  
228 doi:10.1002/2014JA020321., 2014.




- 229 N`emec, F., Bezd`ekova, B., Manninen, J., Parrot, M., Santolik, O., Hayosh, M., and Turunen, T.:  
230 Conjugate observations of a remarkable quasiperiodic event by the low-altitude DEMETER  
231 spacecraft and ground-based instruments, *J. Geophys. Res. Space Physics*, 121, 8790–8803,  
232 doi:10.1002/2016JA022968, 2016.
- 233 Ohya, H., Shiokawa, K., and Miyoshi, Y.: Daytime tweek atmospherics, *J. Geophys. Res. Space*  
234 *Physics*, 120, 654–665, doi:10.1002/2014JA020375, 2015.
- 235 Sato, N., Hayashi, K., Kokubun, S., Oguti, T., and Fukunishi, H.: Relationships between quasi-  
236 periodic VLF emission and geomagnetic pulsation, *J. Atmos. Terr. Phys.*, 36, 1515–1526, 1974.
- 237 Sazhin S. and Hayakawa, M.: Periodic and quasiperiodic VLF emissions, *J. Geophys. Res.*, 56(6),  
238 735–753, 1994.
- 239 Titova, E.E., Kozelov, B.V., Demekhov, A.G., Manninen, J., Santolik, O., Kletzing, C.A., and  
240 Reeves, G.: Identification of the source of quasiperiodic VLF emissions using ground-based  
241 and Van Allen probes satellite observations, *Geophys. Res. Lett.*, 42, 6137–6145,  
242 doi:10.1002/2015GL064911, 2015.
- 243 Trakhtengerts, V.Yu. and Rycroft, M.J.: *Whistler and Alfvén Mode Cyclotron Masers in Space*,  
244 354 pp., Cambridge University Press, 2008.
- 245 Volland, H.: Longwave sferics propagation within the atmospheric waveguide, *Handbook of*  
246 *Atmospheric Electrodynamics*, ed. H Volland (Baton Rouge-London-Tokyo: CRC Press Boca),  
247 2, 65-93, 1995
- 248 Yamashita, M.: Propagation of tweek atmospherics, *J. Atmos. Terr. Phys.*, 40, 151–156, 1978.
- 249 Yedemsky, D.E., Riabov, B.S., Shchekotov, A.I., and Yarotsky, V.S.: Experimental investigation  
250 of the tweek field structure, *Adv. Space Res.*, 12(6), 251-254, 1992.
- 251  
252





253




254

255

256

257 **Fig. 1.** The 2-hour spectrograms of QP events observed on 25 Dec 2011 and 5 Jan 2014. Upper  
258 panels show 0-12 kHz spectrograms without sferics filtering demonstrating that intense sferics  
259 hid all natural VLF emissions at frequencies above 4 kHz. The bottom panels show the same  
260 intervals after filtering out the sferics, and previously unknown events revealed. Colour bars  
261 shown in the right side represent the signal power (in  $\log_{10}(\text{PSD}) [\text{nT}^2/\text{Hz}]$ ).


262



263

264

265 **Fig. 2a.** The 4-hour VLF spectrograms at 6-12 kHz: total power (uppermost panel), right-handed  
266 power (second panel), the wave power/angle distribution based on pure right-hand power at  
267 9.0-10.5 kHz (third panel) and 7.0-8.5 kHz (lowermost panel) on 25 Dec 2011. Colour bars  
268 represent the signal power (in  $\log_{10}(\text{PSD})$  [ $\text{nT}^2/\text{Hz}$ ]).



269

270 **Fig. 2b.** The 4-hour VLF spectrograms at 6-12 kHz: total power (uppermost panel), right-handed  
271 power (middle panel), the wave power/angle distribution based on pure right-hand power at  
272 7.0-11.0 kHz (lowermost panel) on 5 Jan 2014.


273

274

275



276



277


278

279 **Fig. 3a.** The 6-12 kHz VLF spectrograms (total and right-handed powers) of one individual  
280 element ("bullet") of QP emissions at 1710-1730 UT on 25 Dec 2011.

281




282



283

284 **Fig. 3b.** Same event as in Fig 3a., but only last 2 minutes (1728-1730 UT).


285



286

287 **Fig. 4.** The 6-12 kHz VLF spectrograms (total and right-handed powers) and the variation of the  
288 integrated wave intensity in frequency band of 8-11 kHz on 5 Jan 2014.

289




290

291 **Fig. 5.** The 6-12 kHz VLF spectrograms (total and right-handed powers) of two-minutes duration  
292 and the wave power/angle distribution based on pure right-hand power at 8.5-10.5 kHz of the  
293 short-period QP emissions on 5 Jan 2014.

294



295



296

297

298 **Fig. 6.** Geophysical situation during two considered events: (a) – OMNI data (IMF Bz and Psw),

299 the AL and SymH geomagnetic indexes, and the magnetograms from Sodankyla obs;

300 the variation of the planetary Kp –index; (c) – the plasmopause location according to

301 measurements on satellites RBSP (A and B) [<http://enarc.space.swri.edu/PTP>]

302