

Response to the Topical Editor:

Firstly, thank you once again for your valuable comments in an effort to improve the paper. Answers to your concerns follow.

The author corrected the article in accordance with the referees reviews. The discussion section was substantially improved. But there are several technical corrections to be made:

What was changed in line 21 (pg. 1)? Do you removed all the references?

Dear editor, actually on more reference was added to the ones already written in line 21 of page 1. What happened is that in the mark-up version produced with latexdiff the references are marked in red and with a line looking like if they were removed. However, if you look carefully there is a blue bracket at the very end of line 21 near to the right border of the page (in the mark-up version). In the last version of the manuscript (version 4) uploaded on January 7, 2019 the references are in line 20 and 21.

The statement By analyzing the latitudinal profiles, it could be determined that the increment of TEC to produce this LTE was of 241.2% (pg 7, lines 5-6) makes no sense. As far as I understand, LTE is a TEC increment. If you mean to show the LTE intensity, just present TEC value. Currently it is unclear what do you take as a 100%.

Dear editor, indeed, LTE is basically a TEC increment. Therefore, the last sentence of the first paragraph of the conclusions section was slightly changed to indicate better this fact. The increment is calculated with respect to the average of a stable day (being the average 100%). For this case an average of the curves for September 7 and 9, 2017 was taken.

Figures 1 and 5 are to be edited since they take a significant space and do not give too much information. Add a grid and more frequent tick marks to Y-axes for Bz and Dst panels.

Dear editor, a grid was added in every subplot of the two figures and also tick marks in the Y-axes for Bz and Dst were made more frequent

Pictures of differential VTEC (figs. 2 and 3) looks better, but need higher contrast. Try to use colormap seismic from matplotlib package.

Dear editor, colormap changed to seismic in figures.2, 3, and 6 as per requested.

Emergence of a localized total electron content enhancement during the severe geomagnetic storm of September 8, 2017

Carlos Sotomayor-Beltran¹ and Laberiano Andrade-Arenas¹

¹Facultad de Ciencias e Ingeniería, Universidad de Ciencias y Humanidades, Lima 39, Peru

Correspondence: Carlos Sotomayor-Beltran (csotomayor@uch.edu.pe) and Laberiano Andrade-Arenas (landrade@uch.edu.pe)

Abstract.

In this work, the results of the analysis on total electron content (TEC) data before, during and after the geomagnetic storm of September 8, 2017 are reported. One of the responses to geomagnetic storms due to the southern vertical interplanetary magnetic field (B_z) is the enhancement of the electron density in the ionosphere. Vertical TEC (VTEC) from the Center for Orbit determination in Europe (CODE) along with a statistical method were used to identify positive and/or negative ionospheric storms in response to the geomagnetic storm of September 8, 2017. When analysing the response to the storm of September 8, 2017 it was indeed possible to observe an enhancement of the equatorial ionization anomaly (EIA); however what it was unexpected, was the identification of a local TEC enhancement (LTE) to the south of the EIA ($\sim 40^\circ$ S, right over New Zealand and extending towards the south-eastern coast of Australia and also eastward towards the Pacific). This was a very transitory LTE that lasted approximately four hours, starting at $\sim 02:00$ UT on September 8 where its maximum VTEC increase was of 241,2%. Using the same statistical method, comparable LTEs in a similar category geomagnetic storm, the 2015 St. Patrick's day storm, were looked for. However, for the aforementioned storm no LTEs were identified. As also indicated in a past recent study for a LTE detected during the August 15, 2015 geomagnetic storm, an association between the LTE and the excursion of B_z seen during the September 8, 2017 storm was observed as well. Furthermore, it is very likely that a direct impact of the super-fountain effect along with travelling ionospheric disturbances may be playing an important role in the production of this LTE. Finally, it is indicated that the September 8, 2017 LTE is the second one to be detected since the year 2016.

1 Introduction

Anomalies in the ionosphere can be product of different natural phenomena (Afraimovich et al., 2013). For instance earthquakes can produce positive or negative ionospheric anomalies (e.g., Zakharenkova et al., 2008; Yao et al., 2012; Guo et al., 2015; Li et al., 2015; Sotomayor-Beltran, 2019), although such variations are expected to be localized within the earthquake's preparation region (Dobrovolsky et al., 1979). On the other hand, major changes in the ionosphere are caused by geomagnetic storms (e.g., Buonsanto, 1999; Danilov, 2013). The response of the Earth's ionosphere to the geomagnetic storms are known

as ionospheric storms. These ionospheric storms can disrupt technologies relying on transmission of radio frequencies (e.g., Buonsanto, 1999; Borries et al., 2015), and thus they can have an impact in the modern society in general.

In order to understand better ionospheric variability in time and space produced by geomagnetic storms, Global Navigation Satellite System (GNSS) receivers, due to its global coverage, are used as one of the tools for ionospheric studies. According to several studies (e.g., Huang et al., 2005; Mannucci et al., 2005; Astafyeva, 2009), one common response to a geomagnetic storm due to the excursion of the southward interplanetary magnetic field is the significant increment in the equatorial and mid-latitude total electron content (TEC), which manifests as an enhancement of the equatorial ionization anomaly (EIA; Appleton, 1946; McDonald et al., 2011). Such increase of TEC in the EIA is possible to visualize in global ionospheric maps (GIMs). Besides changes in the EIA, it was recently observed by Edemskiy et al. (2018) and Sotomayor-Beltran (2018) that localized TEC enhancements (LTEs) can also emerge as a response to geomagnetic storms.

In this paper vertical TEC (VTEC) maps, also known as global ionospheric maps (GIMs), due to its reliability on ionospheric information (Hernández-Pajares et al., 2009), were used to analyse the response to the geomagnetic storm of September 8, 2017. Section 2 introduces the ionospheric data and the technique for the corresponding analysis. In Sect. 3 the results and the discussion are presented. Section 4 presents the final remarks or conclusions.

2 Data and methods

VTEC maps were downloaded via ftp¹ from the Center for Orbit Determination in Europe (CODE) between August 21, 2017 and September 20, 2017. VTEC maps, which have a resolution of $2.5^\circ \times 5^\circ$ (latitude and longitude, respectively), come in daily IONosphere Map EXchange files (Schaer et al., 1998) and they are produced every hour. Due to the format of the IONEX files, which consists of headers and the actual VTEC data, a code entirely written in Python was implemented for this work. Using the NumPy² library, which handles relatively easily N-dimensional arrays, the VTEC data was stored in a 3D cube for further analysis. The x , y and z axes in the 3D cube are longitude, latitude and number of maps, respectively.

In order to identify ionospheric anomalies a running window of 8 days to every cell in the 3D VTEC cube was applied (e.g., Liu et al., 2004; Zhu et al., 2010; Zou and Zhao, 2010; Li et al., 2015; Sotomayor-Beltran, 2018). Assuming that for each cell or line-of-sight, the VTEC follows a Gaussian distribution, the mean (μ) VTEC and its associated standard deviation (σ) are calculated in order to define the upper and lower bounds:

$$UB = \mu + 2\sigma, \tag{1}$$

$$LB = \mu - 2\sigma. \tag{2}$$

¹<ftp://ftp.aiub.unibe.ch/CODE/>

²<http://www.numpy.org/>

If a VTEC value for a certain day at a particular time falls above the UB , then a positive ionospheric anomaly is detected with a confidence level of 95%. The difference between the VTEC and UB or LB is defined as differential VTEC ($\Delta VTEC$). On the other hand, if the VTEC falls below the LB , then a negative anomaly is detected. In this way, a cube of $\Delta VTEC$ was created, with a total of 744 maps. If $UB > VTEC > LB$, then $\Delta VTEC = 0$

- 5 Some important geomagnetic parameters were also needed to be taken into account for the analysis. The Dst index (Sugiura, 1964) provides information about the strength of the ring current around the Earth. According to Loewe and Pröls (1997) a magnetic storm can be considered as weak when $-50 \text{ nT} < \text{Dst} \leq -30 \text{ nT}$. A moderate and strong storm occurs when $-100 \text{ nT} < \text{Dst} \leq -50 \text{ nT}$ and $-200 \text{ nT} < \text{Dst} \leq -100 \text{ nT}$, respectively. Finally, a severe storm happens when $\text{Dst} \leq -200 \text{ nT}$. For this study Dst data for the month of September 2017 was downloaded from World Data Center for Geomagnetism in Kyoto³.
- 10 Another very important index which measures the fluctuations caused in the Earth's magnetic field by a geomagnetic storm is the Kp index. According to Gosling et al. (1991) when $K_p \geq 8$ - and $K_p \geq 6$ - for at least three 3-h intervals, the storm can be considered a major one. A large storm occurs when $7- \leq K_p \leq 7$ and $K_p \geq 6$ for at least three 3-h intervals. For other cases when $K_p \geq 6$ - for at least three 3-h intervals the storm can be considered of medium strength. Finally, a small storm happens when $5- \leq K_p \leq 5$. Kp data for September 2017 was retrieved from the German Research Centre for Geosciences (GFZ⁴). The
- 15 vertical interplanetary magnetic field (B_z ; Tsurutani et al., 1988) also is a good indicator of a geomagnetic storm. When there is a strong southward B_z for more than 3 hours a geomagnetic storm is in development (Gonzalez et al., 1994; Liu and Li, 2002). Hourly averages for B_z also for the month of September 2017 were downloaded from the OMNI database⁵. In Fig. 1 the Dst and Kp indices and also B_z (in geocentric solar magnetospheric coordinate system) can be observed for a range of days (September 3 – September 16) within the month of September 2017.

20 3 Results and Discussion

Figure 1 shows that $K_p = 8$ during the last 3 hours (UT) of September 7 and the first three hours of September 8. According to the National Oceanic and Atmospheric Administration (NOAA) space weather service⁶, this geomagnetic storm can be classified as a G4 severe storm ($K_p = 8$). Additionally, for September 8, 2017 between 00:00 and 04:00 UT the Dst index had values lower than -100 nT (Fig. 1).

- 25 The origin of this geomagnetic storm lies in the coronal mass ejection (CME) that occurred on September 6, 2017 at $\sim 12:40$ UT. This CME was observed with the Camera 2 of the Large Angle and Spectrometric Coronagraph on board of the Solar and Heliospheric Observatory (SOHO⁷). Figure 1 also shows that on September 8 at $\sim 00:00$ UT the vertical interplanetary magnetic field decreased significantly to a minimum of -24 nT . One hour before (September 7 at 23:00 UT), B_z already decreased considerably to -20.6 nT , time of the storm sudden commencement (Fig. 1). In addition, it can be noticed that almost

³<http://wdc.kugi.kyoto-u.ac.jp/wdc/Sec3.html>

⁴<https://www.gfz-potsdam.de/en/kp-index/>

⁵<https://omniweb.gsfc.nasa.gov/form/dx1.html>

⁶<https://www.swpc.noaa.gov/noaa-scales-explanation>

⁷<https://sohowww.nascom.nasa.gov/>

simultaneously with the drastic change of B_z , the Dst index reached its peak at 01:00 UT on September 8, 2017. As it is already well-known, this relationship between B_z and the Dst index hints to a physical response of the ring current in the magnetosphere to the interplanetary field B_z (Patel and Desai, 1973; Gonzalez and Echer, 2005).

3.1 GIM maps

5 In the left column of Fig. 2, GIMs for September 7, 8 and 9, 2017 at 02:00 UT are presented. It is clearly seen in the GIM of September 8 at 02:00 UT (just three hours after the storm sudden commencement) that the VTEC was enhanced in the EIA region with respect to the day before (September 7) and the day after (September 9) at the same hour. A recent study by Lei et al. (2018), using diverse instruments (e.g., satellites and ionosondes), has also observed this TEC enhancement in the Asian-Australian region for this geomagnetic storm. The increment of VTEC in the EIA was already observed in previous
10 studies about ionospheric responses to geomagnetic storms (e.g., Zhao et al., 2005; Pedatella et al., 2009; Astafyeva et al., 2015; Chakraborty et al., 2015).

3.2 Differential VTEC maps

What it was quite compelling was the detection of a ionospheric localized anomaly ($\sim 40^\circ$ S), or as named by Edemskiy et al. (2018) a localized TEC enhancement (LTE), to the south of the southern conjugate geomagnetic region of the EIA. This LTE
15 can be identified in the GIM map of September 8, 2017 at $\sim 02:00$ UT (Fig. 2). In the right column of Fig. 2, Δ VTEC maps for September 7, 8 and 9, 2017 at 02:00 UT are also presented. It can be seen from these Δ VTEC maps that a day before and after that the LTE appeared, no anomalies were visible. However as already indicated, the day that the ionospheric storm occurred (September 8), the dramatic enhancement of the VTEC to the south of the EIA, manifested as a LTE, was observed.

In Fig. 3 the dynamics of the LTE can be clearly seen. It can be noticed that this LTE was very transitory, in the Δ VTEC
20 maps it appeared at $\sim 02:00$ UT on September 8 and at $\sim 06:00$ UT it was already gone. This unforeseen positive ionospheric storm covers most of New Zealand and extends westward towards the south-eastern part of Australia and eastward towards the Pacific. The maximum peak of this LTE happened as well on September 8 at 02:00 UT with Δ VTEC = 6.47 TECU (where 1 TECU = 10^{16} electrons/m²).

3.3 Shape of the EIA

25 To better visualize this LTE to the south of the EIA, the shape of the VTEC along the meridional line of 170° E is shown in Fig. 4 between September 7 and 9, 2017 at 02:00 UT. From the Δ VTEC maps (Fig. 2), it can be confirmed that the EIA follows its normal variability one day after (September 9 at 02:00 UT) and before (September 7 at 02:00 UT) that the storm occurred (no anomalous VTEC enhancements are visible). However, on September 8 at 02:00 UT the EIA was significantly enhanced and hence this translated in a much sharper definition of the double-crest with a trough shape observed in Fig. 4. This shape is
30 expected because when the LTE is above New Zealand, it is still day time, the local time is 14:00 (02:00 UT). In addition to the two crests from the EIA, a third one in the southern hemisphere is visible (Fig. 4). This third crest is simply the LTE observed

in the Δ VTEC and GIM maps for September 8 at 02:00 UT (Fig. 2 and 3). The peak increment for this day and this time in the southern crest of the EIA is of 172% and in the LTE of 241,2%. Edemskiy et al. (2018) have also reported for the August 15, 2015 G3 geomagnetic storm that the two LTEs they observed were located to the south of the EIA (between Africa and Antarctica), whereas Sotomayor-Beltran (2018) has also identified to the south of the EIA a LTE over the Indian ocean during the G2 moderate storm of April 20, 2018.

3.4 The St. Patrick's day 2015 geomagnetic storm

In order to look for comparable LTEs in a similar geomagnetic storm category, the author turned to the G4 geomagnetic storm that occurred during the St. Patrick's day of 2015 which has been thoroughly studied (Astafyeva et al., 2015; Cherniak et al., 2015; Nava et al., 2016; Yao et al., 2016; Jin et al., 2017; Zhang et al., 2018). In Fig. 5, the variability of the geomagnetic indices, Dst and Kp, and the vertical interplanetary magnetic field for a period of days (March 13 – March 27) in the month of March 2015 can be observed. GIMs and Δ VTEC maps are shown in Fig. 6. In the Δ VTEC maps it was possible to observe a positive ionospheric storm starting on March 17, 2015 at \sim 18:00 UT right over the southern Atlantic, right north off the Antarctic coast. This positive storm started to move westward and it reached its maximum strength on March 18, 2015 at \sim 02:00 UT with a peak of Δ VTEC = 12.88 TECU (Fig. 6). In this case however, the enhancement of VTEC observed in the southern hemisphere is not a LTE, it is only the southern crest of the EIA which underwent an increment of VTEC and shifted several degrees southward. On the other hand in the Δ VTEC maps of March 17, 2015 starting at \sim 22:00 UT, negative ionospheric storms were also observed and they lasted until the end of the day of March 18, 2015. These both results agree well with the ones from previous studies, using different methods, for the St. Patrick's day 2015 storm (Astafyeva et al., 2015; Yao et al., 2016). It can also be finally noticed in Fig. 6, in the Δ VTEC maps, that at 02:00 UT the day before and the day after the maximum peak of the positive ionospheric storm there are no anomalous variations of the observed TEC.

3.5 Creation of September 8, 2017 LTE

For the case of the St. Patrick storm of 2015, for the observed positive one in the southern hemisphere and negative storm in the northern hemisphere, Astafyeva et al. (2015) and Yao et al. (2016) indicated three suitable candidates as the origin mechanisms: the strength of the geomagnetic field, the B_y component of the interplanetary magnetic field and composition changes in the thermosphere. On the other hand, for the moderate G3 storm of August 15, 2015 there was not a clear mechanism put forward by Edemskiy et al. (2018) to account for the observed LTEs. Only a dependence of the emergence of these LTEs to the interplanetary B_z was hinted at, but still as indicated by the authors of that study it was not their definite conclusion.

For the LTE observed during the September 8, 2017 severe storm in this work, an excursion of the interplanetary B_z along with a consequent decrease of the Dst index was also observed (Fig. 1). Thus, it can be suggested that there is as well an association between the interplanetary B_z and the emergence of the LTE. In this vein, Lei et al. (2018) have gone further and indicated that for the September 8, 2017 geomagnetic storm not only B_z could produce prompt penetration electric fields (PPEFs) which enhance the EIA (super-fountain effect) but also could produce traveling atmospheric disturbances (TADs). These TADs, which originate in the polar regions, can transport equatorward winds that drive plasma upwards in the middle

and lower latitudes and as a consequence the ionosphere moves to higher altitudes (Chen et al., 2016). It is very likely then, as suggested by Lei et al. (2018), that the combined effect of TADs and the PPEFs are responsible for the creation of the LTE observed in Fig. 2 and 3. As per to the overall enhancement of the EIA (Fig. 2 and 3) and shifting of the crests in the direction of the poles observed in Fig. 4, as previously mentioned and suggested by many studies (e.g., Tsurutani et al., 2004; Mannucci et al., 2005; Astafyeva, 2009; Astafyeva et al., 2014; Chakraborty et al., 2015) the mechanism at work is the ionospheric super-fountain effect. Finally, it is also worth mentioning that this would be the second time a LTE is detected since 2016, as the first one was the one observed during the April 20, 2018 geomagnetic storm (Sotomayor-Beltran, 2018).

4 Conclusions

Ionospheric response to the G4 severe geomagnetic storm of September 8, 2017 was analysed by using VTEC maps from CODE along with a statistical method to identify ionospheric anomalies. By producing differential VTEC maps it was possible to identify not only an enhancement of the EIA but also a localized TEC enhancement. The maximum intensity of this LTE was on September 8, 2017 at 02:00 UT and it was localized right over New Zealand and extending towards the south-eastern coast of Australia and eastward towards the Pacific. The LTE was quite transitory, it lasted only for about four hours and on September 8 at 06:00 UT it faded away. This LTE is the second one to be observed since 2016. By analyzing the latitudinal profiles, it could be determined that the ~~increment of TEC to produce this LTE was~~ maximum VTEC increment, where the LTE was observed, was of 241.2%.

Due to its category, the G4 storm from March 17, 2015 was also investigated in order to look for comparable LTEs. However, there was no LTE detections and instead a hemispheric asymmetry of ionospheric storms in the northern and southern hemisphere was observed. One geomagnetic storm which presented the same traits (LTEs) as in the one of September 8, 2017 was the G3 August 15, 2015 moderate storm. During this storm also LTEs were identified south of the geomagnetic conjugate region of the EIA. These LTEs, was indicated, seem to be associated with the negative excursion of B_z .

For the September 8, 2017 storm in the present study also such negative excursion of the vertical component of the interplanetary magnetic field was observed; hence, it can be suggested then that this has an effect on the origin of the LTE. Furthermore, it is very likely that TADs along with the super-fountain effect, the two of them due to B_z , are having a significant effect in the generation of the observed LTE. To shed more light into how these LTEs are created, further observations of these events along with physical modeling of the effects of the B_z on the super-fountain effect, TADs and possibly other contributing ionospheric mechanisms would be needed.

Competing interests. The authors declare that they do not have conflict of interest.

Acknowledgements. The ~~author is~~ authors are very grateful to the anonymous referees for their constructive and insightful comments that helped in improving the manuscript. The ~~author~~ authors would like to thank CODE for making publicly available IONEX files and also would like to acknowledge the World Data Center for Geomagnetism in Kyoto, the German Research Center for Geosciences and the OMNI database for providing data for the Dst index, the Kp index and the vertical interplanetary B_z , respectively.

References

- Afraimovich, E. L., Astafyeva, E. I., Demyanov, V. V., Edemskiy, I. K., Gavriluk, N. S., Ishin, A. B., Kosogorov, E. A., Leonovich, L. A., Lesyuta, O. S., Palamartchouk, K. S., Perevalova, N. P., Polyakova, A. S., Smolkov, G. Y., Voeykov, S. V., Yasyukevich, Y. V., and Zhivetiev, I. V.: A review of GPS/GLONASS studies of the ionospheric response to natural and anthropogenic processes and phenomena, *Journal of Space Weather and Space Climate*, 3, A27, 2013.
- 5 Appleton, E. V.: Two anomalies in the Ionosphere, *Nature*, 157, 691, 1946.
- Astafyeva, E.: Effects of strong IMF B_z southward events on the equatorial and mid-latitude ionosphere, *Annales Geophysicae*, 27, 1175–1187, 2009.
- Astafyeva, E., Yasyukevich, Y., Maksikov, A., and Zhivetiev, I.: Geomagnetic storms, super-storms, and their impacts on GPS-based navigation systems, *Space Weather*, 12, 508–525, 2014.
- 10 Astafyeva, E., Zakharenkova, I., and Förster, M.: Ionospheric response to the 2015 St. Patrick’s Day storm: A global multi-instrumental overview, *Journal of Geophysical Research: Space Physics*, 120, 9023–9037, 2015.
- Borries, C., Berdermann, J., Jakowski, N., and Wilken, V.: Ionospheric storms—A challenge for empirical forecast of the total electron content, *Journal of Geophysical Research: Space Physics*, 120, 3175–3186, 2015.
- 15 Buonsanto, M.: Ionospheric Storms - A Review, *Space Science Reviews*, 88, 563–601, 1999.
- Chakraborty, M., Kumar, S., De, B. K., and Guha, A.: Effects of geomagnetic storm on low latitude ionospheric total electron content: A case study from Indian sector, *Journal of Earth System Science*, 124, 1115–1126, 2015.
- Chen, X., Lei, J., Wang, W., Burns, A. G., Luan, X., and Dou, X.: A numerical study of nighttime ionospheric variations in the American sector during 28–29 October 2003, *Journal of Geophysical Research: Space Physics*, 121, 8985–8994, 2016.
- 20 Cherniak, I., Zakharenkova, I., and Redmon, . J.: Dynamics of the high-latitude ionospheric irregularities during the 17 March 2015 St. Patrick’s day storm: Ground-based GPS measurements, *Space Weather*, 13, 585–597, 2015.
- Danilov, A. D.: Ionospheric F-region response to geomagnetic disturbances, *Advances in Space Research*, 52, 343–366, 2013.
- Dobrovolsky, I. P., Zubkov, S. I., and Miachkin, V. I.: Estimation of the size of earthquake preparation zones, *Pure and Applied Geophysics*, 117, 1025–1044, 1979.
- 25 Edemskiy, I., Lastovicka, J., Buresova, D., Habarulema, J. B., and Nepomnyashchikh, I.: Unexpected Southern Hemisphere ionospheric response to geomagnetic storm of 15 August 2015, *Annales Geophysicae*, 36, 71–79, 2018.
- Gonzalez, W. D. and Echer, E.: A study on the peak Dst and peak negative B_z relationship during intense geomagnetic storms, *Geophysical Research Letters*, 32, 2005.
- Gonzalez, W. D., Joselyn, J. A., Kamide, Y., Kroehl, H. W., Rostoker, G., Tsurutani, B. T., and Vasylunas, V. M.: What is a geomagnetic storm?, *Journal of Geophysical Research: Space Physics*, 99, 5771–5792, 1994.
- 30 Gosling, J. T., McComas, D. J., Phillips, J. L., and Bame, S. J.: Geomagnetic activity associated with earth passage of interplanetary shock disturbances and coronal mass ejections, *Journal of Geophysical Research: Space Physics*, 96, 7831–7839, 1991.
- Guo, J., Li, W., Yu, H., Liu, Z., Zhao, C., and Kong, Q.: Impending ionospheric anomaly preceding the Iquique M_w 8.2 earthquake in Chile on 2014 April 1, *Geophysical Journal International*, 203, 1461–1470, 2015.
- 35 Hernández-Pajares, M., Juan, J. M., Sanz, J., Orus, R., Garcia-Rigo, A., Feltens, J., Komjathy, A., Schaer, S. C., and Krankowski, A.: The IGS VTEC maps: a reliable source of ionospheric information since 1998, *Journal of Geodesy*, 83, 263–275, 2009.

- Huang, C.-S., Foster, J. C., and Kelley, M. C.: Long-duration penetration of the interplanetary electric field to the low-latitude ionosphere during the main phase of magnetic storms, *Journal of Geophysical Research: Space Physics*, 110, 2005.
- Jin, S., Jin, R., and Kutoglu, H. J.: Positive and negative ionospheric response to the March 2015 geomagnetic storm from BDS observations, *Journal of Geodesy*, 91, 613–626, 2017.
- 5 Lei, J., Huang, F., Chen, X., Zhong, J., Ren, D., Wang, W., Yue, X., Luan, X., Jia, M., Dou, X., Hu, L., Ning, B., Owolabi, C., Chen, J., Li, G., and Xue, X.: Was Magnetic Storm the Only Driver of the Long-Duration Enhancements of Daytime Total Electron Content in the Asian-Australian Sector Between 7 and 12 September 2017?, *Journal of Geophysical Research: Space Physics*, 123, 3217–3232, 2018.
- Li, J., You, X., Zhang, R., Meng, G., Shi, H., and Han, Y.: Ionospheric total electron content disturbance associated with May 12, 2008, Wenchuan earthquake, *Geodesy and Geodynamics*, 70, 2015.
- 10 Li, J., You, X., Zhang, R., Meng, G., Shi, H., and Han, Y.: Ionospheric total electron content disturbance associated with May 12, 2008, Wenchuan earthquake, *Geodesy and Geodynamics*, 70, 2015.
- Liu, J. Y., Chuo, Y. J., Shan, S. J., Tsai, Y. B., Chen, Y. I., Pulinets, S. A., and Yu, S. B.: Pre-earthquake ionospheric anomalies registered by continuous GPS TEC measurements, *Annales Geophysicae*, 22, 1585–1593, 2004.
- Liu, S.-L. and Li, L.-W.: Study on Relationship between Southward IMF Events and Geomagnetic Storms, *Chinese Journal of Geophysics*, 15 45, 301–310, 2002.
- Loewe, C. A. and Pröls, G. W.: Classification and mean behavior of magnetic storms, *Journal of Geophysical Research: Space Physics*, 102, 14 209–14 213, 1997.
- Mannucci, A. J., Tsurutani, B. T., Iijima, B. A., Komjathy, A., Saito, A., Gonzalez, W. D., Guarnieri, F. L., Kozyra, J. U., and Skoug, R.: Dayside global ionospheric response to the major interplanetary events of October 29–30, 2003 “Halloween Storms”, *Geophysical Research Letters*, 32, 2005.
- 20 McDonald, S. E., Coker, C., Dymond, K. F., Anderson, D. N., and Araujo-Pradere, E. A.: A study of the strong linear relationship between the equatorial ionization anomaly and the prereversal E X B drift velocity at solar minimum, *Radio Science*, 46, 1–9, 2011.
- Nava, B., Rodríguez-Zuluaga, J., Alazo-Cuartas, K., Kashcheyev, A., Migoya-Orué, Y., Radicella, S. M., Amory-Mazaudier, C., and Fleury, R.: Middle- and low-latitude ionosphere response to 2015 St. Patrick’s Day geomagnetic storm, *Journal of Geophysical Research: Space Physics*, 121, 3421–3438, 2016.
- 25 Patel, V. L. and Desai, U. D.: Interplanetary magnetic field and geomagnetic Dst variations, *Astrophysics and Space Science*, 20, 431–437, 1973.
- Pedatella, N. M., Lei, J., Larson, K. M., and Forbes, J. M.: Observations of the ionospheric response to the 15 December 2006 geomagnetic storm: Long-duration positive storm effect, *Journal of Geophysical Research: Space Physics*, 114, 2009.
- 30 Schaer, S., Gurtner, W., and Feltens, J.: IONEX: The ionosphere map exchange format version 1, pp. 233–247, 1998.
- Sotomayor-Beltran, C.: Localized Increment and Decrement in the Total Electron Content of the Ionosphere as a Response to the April 20, 2018, Geomagnetic Storm, *International Journal of Geophysics*, 2018, 2018.
- Sotomayor-Beltran, C.: Ionospheric anomalies preceding the low-latitude earthquake that occurred on April 16, 2016 in Ecuador, *Journal of Atmospheric and Solar-Terrestrial Physics*, 182, 61–66, 2019.
- 35 Sugiura, M.: Hourly value of equatorial Dst for the IGY, *Annals of the International Geophysical Year*, 35, 1964.
- Tsurutani, B., Mannucci, A., Iijima, B., Abdu, M. A., Sobral, J. H. A., Gonzalez, W., Guarnieri, F., Tsuda, T., Saito, A., Yumoto, K., Fejer, B., Fuller-Rowell, T. J., Kozyra, J., Foster, J. C., Coster, A., and Vasyliunas, V. M.: Global dayside ionospheric uplift and enhancement associated with interplanetary electric fields, *Journal of Geophysical Research: Space Physics*, 109, 2004.

- Tsurutani, B. T., Gonzalez, W. D., Tang, F., Akasofu, S. I., and Smith, E. J.: Origin of interplanetary southward magnetic fields responsible for major magnetic storms near solar maximum (1978–1979), *Journal of Geophysical Research: Space Physics*, 93, 8519–8531, 1988.
- Yao, Y., Liu, L., Kong, J., and Zhai, C.: Analysis of the global ionospheric disturbances of the March 2015 great storm, *Journal of Geophysical Research: Space Physics*, 121, 157–170, 2016.
- 5 Yao, Y. B., Chen, P., Wu, H., Zhang, S., and Peng, W. F.: Analysis of ionospheric anomalies before the 2011 M_w 9.0 Japan earthquake, *Chinese Science Bulletin*, 57, 500–510, 2012.
- Zakharenkova, I. E., Shagimuratov, I. I., Tepenitzina, N. Y., and Krankowski, A.: Anomalous modification of the ionospheric total electron content prior to the 26 September 2005 Peru earthquake, *Journal of Atmospheric and Solar-Terrestrial Physics*, 70, 1919–1928, 2008.
- Zhang, W., Zhao, X., Jin, S., and Li, J.: Ionospheric disturbances following the March 2015 geomagnetic storm from GPS observations in
10 China, *Geodesy and Geodynamics*, 2018.
- Zhao, B., Wan, W., and Liu, L.: Response of equatorial anomaly to the October–November 2003 superstorm, *Annales Geophysicae*, 23, 2005.
- Zhu, F., Wu, Y., Lin, J., and Zhou, Y.: Temporal and spatial characteristics of VTEC anomalies before Wenchuan M_s 8.0 earthquake, *Geodesy and Geodynamics*, 1, 23–28, 2010.
- Zou, Y. and Zhao, T.: Ionospheric anomalies detected by GPS TEC measurements during the 15 August 2007 Peru earthquake, in: 2010
15 International Conference on Microwave and Millimeter Wave Technology, pp. 1216–1219, 2010.




Figure 1. The vertical component of interplanetary magnetic field (B_z) and the Dst and Kp indices between September 3 and September 16, 2017. The vertical red dashed line in all the plots points to the storm sudden commencement.




Figure 2. Left column: Global ionospheric maps for September 7, 8 and 9, 2017 at 02:00 UT. Right column: Differential VTEC maps for September 7, 8 and 9, 2017 at 02:00 UT.




Figure 3. Differential VTEC maps for September 8, 2017 between 00:00 and 06:00 UT.




Figure 4. Structure of the VTEC for the 170°E meridian at 02:00 UT between September 7 and 9, 2017. A relevant range of latitudes is shown, 62.5°N–62.5°S. The vertical dashed black line indicates the Equator (latitude = 0°).




Figure 5. The vertical component of the interplanetary magnetic field (B_z) and the Dst and Kp geomagnetic indices between March 13 and March 27, 2015. The vertical red dashed line in all the plots indicates the day that the 2015 St. Patrick's day storm occurred (March 17, 2015).




Figure 6. Left column: Global ionospheric maps for March 17, 18 and 19, 2015 at 02:00 UT. Right column: Differential VTEC maps for March 17, 18 and 19, 2015 at 02:00 UT.