

Reply to the report by Referee_1

1. *The correctness of the transition from equations (1), (2) to equations (5), (6) when considering divergence-free waves ($\text{div}V = 0$) is in doubt. Since in an incompressible medium the speed of sound tends to infinity, the product “infinity to zero” in square brackets of equations (1), (2) becomes uncertain.*

We consider a compressible atmosphere stratified in a field of gravity. In equations (1), (2), the speed of sound refers to a compressible medium and is the final value, whereas the condition $\text{div}V = 0$ determines the properties of the perturbations only.

2. *The realization of the obtained modes was considered in the framework of a simplified model of an infinitely thin discontinuity in the altitude profile of temperature. In the atmospheres of the planets, the situation is most likely realized when the change of the parameters along the vertical occurs on scales of tens of kilometers, or even hundreds, for the Sun. Can the acceptance of the finite thickness of the transition layer significantly affect the conditions of realization of the evanescent wave modes considered in the work?*

The simplest model of a thin temperature gap is considered as an example in order to show the fundamental possibility of implementing the new types of wave modes obtained in the work. To understand how the thickness of the transition layer affects the properties of the modes considered, a separate study is needed. This effect seems to be significant. Especially when the magnitude of the transition layer is commensurate with the atmosphere scale height for the upper or lower isothermal half-spaces. In our opinion, it is more expedient to investigate the implementation of the received modes within the framework of an atmospheric model with a continuous non-isothermal altitude profile of temperature.

3. *In the atmosphere, different types of gravity disturbances may occur: (1) freely propagating waves, having a real and non-zero vertical component of the wave vector; (2) evanescent wave modes, propagating only horizontally. Does it mean that sources of evanescent modes and freely propagating waves are fundamentally different?*

The problem of sources was not analyzed in the work, we considered free waves (on the right side of equations (1), (2) there are zeros). It is unlikely that the sources of atmospheric acoustic-gravity waves of different types must necessarily have a different nature. In our opinion, if the sources are localized in the isothermal interval of the heights of the atmosphere, they generate a freely propagating AGW more effectively. At heights of sharp temperature gradients, evanescent wave modes are preferred, since “surfaces” arise that support the propagation of such waves. This question requires separate study.

4. It is not clear from the manuscript how complete is the list of possible evanescent modes. Are there additional requirements for disturbances that will lead to new solutions?

It is likely that equations (1), (2) admit the possibility of the existence of other types of evanescent wave modes. When imposing other additional conditions on the properties of disturbances, besides those considered in the article, other types of evanescent modes can be obtained and Table 1 can be supplemented.

The following inaccuracies

1) The phrase “the properties of the medium” should be replaced by “the properties of the disturbances”.

2) The bottom line of Table 2 (line 2, p. 20) uses the abbreviations AMp, AM, which, apparently, should be replaced by AE and AEp.

3) Mistakes are made in the names of subsections 2 and 2.1 (line 1, page 3 and line 19, page 3). Probably should be

2 Evanescent modes in the isothermal atmosphere

2.1 Non-divergent and pseudo-non-divergent modes

We have checked and corrected these inaccuracies in the text.

Reply to to the report by Referee_2

The title: Perhaps, the authors should modify the title of the paper, as it addresses also the case of a vertically non-isothermal atmosphere.

In the first part of the work, we considered separate types of modes in an unlimited isothermal atmosphere. In the second part, we studied the possibility of realizing the modes at a temperature discontinuity, but for each half-space within the isothermal model. Therefore, we are of the opinion that the term “isothermic atmosphere” in the title is appropriate.

p. 1, l. 23 "consisting of acoustic and gravity regions" - are those regions on the dispersion plane or in different parts of the atmosphere?

We changed the sentence to «consisting of acoustic and gravity regions on the dispersion plane».

p. 2, l. 30: the authors claim that the possibility of the existence of a new type of evanescent acoustic-gravity modes is proved in the paper. Could the authors explain why this mode has been missing from the vast amount of previous studies of this problem? In other words, which novel element (assumption or method) allowed the authors to identify this previously unknown mode.

We assumed that wave disturbances may exist in a stratified compressible atmosphere that satisfy the new additional conditions. Under the assumption of perturbation incompressibility ($\text{div}V = 0$), the known ND mode was obtained, and under the assumption of perturbation inelasticity ($\text{div}(\rho_0 V) = 0$) a new AE mode was obtained. In the text of the article indicated, under which assumption each of these modes is obtained.

p. 3, l. 8: Please mention that the sound speed is determined by the temperature.

We added in the text the definition of the atmospheric scale height in the form $H = kT / mg$, clearly indicating the dependence on temperature.

Eq. (11): the RHS of the equation may be confusing: it is not clear that it actually consists

of two different lines corresponding to different signs on the LHS. Please modify the equation, by, e.g., adding a comma after k_x in the top row, and a full stop after k_x in the bottom row.

Equation (11) was written in one line.

It would be instructive to link the term "anelastic" with the terms "compressive" or "incompressive", which are commonly used in the solar atmospheric research.

According to the physical meaning, ND mode is “incompressible” ($\text{div}V = 0$), the other considered modes, that is, NDp, AE, AEp modes, are “compressible” ($\text{div}V \neq 0$). We use the term "anelastic" for disturbances with $\text{div}(\rho_0 V) = 0$.

The term "an unlimited atmosphere" would perhaps sound better as "an unbounded atmosphere".

Replaced.

p. 10, l. 8: Please give the physical meaning of this boundary condition. In other words, the continuity of which physical quantity or quantities should be kept across the interface?

Obviously, if the atmosphere is barometric, then the equilibrium pressure p_0 should be continuous across the interface, and hence the value of ρc^2 also. In addition, for the perturbed values, we require continuity of the vertical velocity component V_z (kinematic condition) and perturbed pressure (dynamic condition). Under these assumptions, we obtained equations (29), (30). For more details see, for example, Tolstoy (1963), Rosental and Gough (1994), Cheremnykh et al. (2018a).

Throughout the paper: please use "equation" instead of "equality".

Corrected.

p. 11, l. 1: It is not clear how the 8th order polynomial in Eq. (31) is obtained from Eq.(30) which has a 4th order polynomial in the numerator.

To get rid of the radicals in expressions (28), (29), which determine the values of a_1 and a_2 , these expressions were squared several times when a polynomial was obtained. The procedure for obtaining a polynomial from boundary conditions (30) is described in Miles and Roberts (1992). Note that expression (31) is only part of the full polynomial expression obtained in Miles and Roberts (1992). Moreover, in (31) two non-physical roots are omitted.

p. 15, l. 4: "the f-mode observed on the Sun should not be compared with the nondivergent ND mode, but with non-divergent pseudo-mode NDp." First of all, I think that the word "associated" would be better than "compared" in this context. Anyway, please explain the physical implications of this association (or comparison).

We agree that "associated" better reflects the meaning of the statement. ND mode and pseudo – mode NDp have the same variance. Therefore, when observing $\omega(kx)$, these modes are indistinguishable. The modes differ in the sign of polarization (in one mode, the V_x oscillations are ahead of V_z by 90° , and in the other mode they are 90° behind) and the pattern of amplitude variation with height. Physically, ND mode is "incompressible" ($\text{div}V = 0$), and NDp mode is "compressive" ($\text{div}V \neq 0$). In the framework of the considered model, only the NDp mode can satisfy the condition of energy reduction in both sides of the interface if the temperature in the upper half-space is higher than in the lower half-space.

Table 2 and 3: Please remind the abbreviations used in the tables (i.e., "L", "Lp", "BV", "BVp", etc.) in the captions. It would allow using those tables in review papers and presentations.

We gave the full names of the modes in the headings of Tables 2 and 3.

EVANESCENT ACOUSTIC-GRAVITY MODES IN THE ISOTHERMAL ATMOSPHERE: SYSTEMATIZATION, APPLICATIONS TO THE EARTH'S AND SOLAR ATMOSPHERES

Oleg K. Cheremnykh, Alla K. Fedorenko, Evgen I. Kryuchkov, Yuriy A. Selivanov

Space Research Institute NASU-SSAU, Kyiv, 03187, Ukraine

Correspondence to: Yuriy A. Selivanov (yuraslv@gmail.com)

Abstract. The objects of research in this work are evanescent wave modes in a gravitationally stratified atmosphere and their associated pseudo-modes. Whereas the former, according to the dispersion relation, rapidly decrease with distance from a certain surface, the latter, having the same dispersion law, differ from the first by the form of polarization and the nature of decreasing from the surface. Within a linear hydrodynamic model, the propagation features of evanescent wave modes in an isothermal atmosphere are studied. Research carried out for different assumptions about the properties of the **disturbances**. On this way, a new wave mode - anelastic evanescent wave mode - was discovered that satisfies the dispersion relation $\omega^2 = k_x g (\gamma - 1)$. Also, the possibility of the existence of a pseudo-mode related to it is indicated. The case of two isothermal media differing in temperature at the interface is studied in detail. It is shown that a non-divergent pseudo-mode with a horizontal scale $k_x \sim 1/2H_1$ can be realized on the interface with dispersion $\omega^2 = k_x g$. Dispersion relation $\omega^2 = k_x g (\gamma - 1)$ at the interface of two media is satisfied by the wave mode, which has different types of amplitude versus height dependencies at different horizontal scales k_x . The applicability of the obtained results to clarify the properties of f -mode observed on the Sun is analyzed.

Keywords: acoustic-gravity waves, evanescent wave modes, isothermal atmosphere, solar atmosphere, earth's atmosphere

1 Introduction

Acoustic - gravity waves (AGWs) in the Earth's atmosphere are studied theoretically and experimentally for more than 60 years. The linear theory of AGW (Hines, 1960; Yeh and Liu, 1974; Francis, 1975) admits the existence in the atmosphere of a continuous spectrum of freely propagating waves, consisting of acoustic and gravity regions on the dispersion plane, as well as of evanescent modes, which can only propagate horizontally.

The freely propagating AGWs effectively transfer the energy and momentum between various atmospheric layers and thus play an important role in the dynamics and energy balance of the atmosphere. These waves are generated by various sources (both natural and technogenic ones), which are accompanied by a significant energy output into the atmosphere. Further, when the AGWs propagate upward the energy conservation compensates for the decrease of the atmospheric density with the height by exponentially increasing amplitude. Therefore at a certain height the waves become nonlinear. Significant progress in the development of the nonlinear theory of AGW was achieved by a number of authors, in particular, Belashov (1990), Nekrasov et al. (1995), Kaladze et al. (2008), Stenflo and Shukla (2009), Huang et al. (2014). Numerical modeling of the freely propagating AGWs in the realistic viscous and heat-conducting atmosphere is an important area of modern studies of these waves (i.e. Vadas, 2012; Cheremnykh et al., 2010).

Satellite observations of AGWs in the Earth's polar thermosphere indicate a prevailing presence of waves with oscillation periods concentrated around the Brunt-Väisälä period and of horizontal scale of about 500 - 700 km (Johnson et al., 1995; Innis and Conde, 2002; Fedorenko et al., 2015). Azimuths of the propagation of these AGW demonstrate the close connection with the directions of background winds in the thermosphere. Moreover, the amplitudes of the waves depend on the speed of headwind but do not depend on height (Fedorenko and Kryuchkov, 2013; Fedorenko et al., 2018). These experimental results cannot be sufficiently explained by the theory of freely propagating AGWs. They may indicate waveguide or evanescent (along a horizontal surface) propagation of at least part of the observed waves.

As well as freely propagating AGWs, evanescent wave modes also play an important role in atmospheric dynamics of the Sun and planets. Evanescent waves propagate horizontally in an atmosphere, vertically stratified by gravity, subject to the presence of vertical gradients of parameters. The energy of these waves should decrease both up and down from the level at which they are generated. Therefore, evanescent waves are most effectively generated in areas of presence of significant vertical gradients of temperature and density or strong local currents. For example, in the solar atmosphere suitable conditions for realization evanescent modes occur at the boundary

between the chromosphere and corona (Jones, 1969). In the Earth's atmosphere, such waves can be efficiently generated at sharp vertical temperature gradients, for example, at the base of the thermosphere or at the heights of the tropo- and mesopause. Also, evanescent wave modes can emerge in the presence of strong inhomogeneous winds, for example, in the region of the polar circulation of the thermosphere.

The study of evanescent waves traditionally gets less attention than the study of freely propagating AGWs. The most known of them are the horizontal Lamb wave and vertical oscillations with Brunt–Väisälä (BV) frequency (Beer, 1974; Waltercheid and Hecht, 2003). In hydrodynamics, physics of terrestrial and solar atmosphere, the surface gravity mode with dispersion $\omega^2 = k_x g$ is also well studied (Tolstoy, 1963; Jones, 1969). In particular, it was shown that it is the fundamental mode (f - mode) of oscillations in the solar atmosphere (Jones, 1969). Experimental f - mode observations are used to study flows, refinement of the solar radius and other parameters of the Sun (Ghosh et al., 1995; Antia, 1998). In the earth's atmosphere, evanescent waves are often observed at altitudes near the mesopause using ground-based instrumentation (Shimkhada et al., 2009).

In this paper, different types of evanescent acoustic-gravity modes characteristic of an isothermal atmosphere are investigated using a set of linearized hydrodynamic equations. In particular, the possibility of the existence of a new type of evanescent acoustic-gravity modes with the dispersion $\omega^2 = k_x g(\gamma - 1)$ is proved **in the assumption of anelasticity of the disturbance**. Also the possibility of realization the evanescent modes in the model of a thin temperature gap studied.

2 Evanescent modes in the isothermal atmosphere

Consider an unbounded ideal isothermal atmosphere, stratified in a field of gravity. Linear perturbations in such a medium satisfy a set of four first order hydrodynamic equations (Hines, 1960). These equations are convenient to bring to a set of two second order equations for the perturbations of the horizontal V_x and vertical V_z particle velocities (Tolstoy, 1963):

$$\rho_0 \frac{\partial^2 V_x}{\partial t^2} = -\rho_0 g \frac{\partial V_z}{\partial x} + \frac{\partial}{\partial x} \left[\rho_0 c^2 \left(\frac{\partial V_x}{\partial x} + \frac{\partial V_z}{\partial z} \right) \right], \quad (1)$$

$$\rho_0 \frac{\partial^2 V_z}{\partial t^2} = \rho_0 g \frac{\partial V_x}{\partial z} + \frac{\partial}{\partial z} \left[\rho_0 c^2 \left(\frac{\partial V_x}{\partial x} + \frac{\partial V_z}{\partial z} \right) \right], \quad (2)$$

where ρ_0 , γ , g denote background atmosphere density, ratio of specific heats, acceleration of gravity, respectively; $c = \sqrt{\gamma g H}$ is the sound speed, $H = -\rho_0 / (d\rho_0/dz) = kT / mg$ is the density scale height, T is the temperature, k is the Boltzmann constant, m is the molecular mass of the atmospheric gas.

Solutions to the system (1), (2) are usually searched for in the form:

$$V_x, V_z \sim \exp(az) \exp[i(\omega t - k_x x)] , \quad (3)$$

where ω , k_x are cyclic frequency and horizontal component of the wave vector, respectively; parameter a sets the vertical scale of the change in the amplitude of velocities, V_x and V_z , with the height, z . For brevity, we will refer to a as the stratification of the corresponding mode.

The system (1), (2) allows, on the plot “frequency-wave number”, for the existence of gravity and acoustic regions of freely propagating waves, for which $a = \frac{1}{2H} \pm ik_z$ (Hines, 1960), where k_z is the vertical component of the wave vector. Also, from (1), (2) we get the solutions in the form of evanescent wave modes having real a and propagating horizontally (Waltercheid and Hecht, 2003). Solutions in the form of evanescent modes are usually obtained by imposing additional conditions on the perturbation properties.

2.1 Non - divergent and pseudo - non - divergent modes

Let us note the well known in hydrodynamics approximation of perturbations incompressibility (see, e.g., Ladikov-Roev et al., 2010), for which

$$\text{div} \vec{V} = \frac{\partial V_x}{\partial x} + \frac{\partial V_z}{\partial z} = 0 . \quad (4)$$

In frames of this approximation, we obtain the following equations from (1), (2):

$$\frac{\partial^2 V_x}{\partial t^2} = -g \frac{\partial V_z}{\partial x} , \quad (5)$$

$$\frac{\partial^2 V_z}{\partial t^2} = g \frac{\partial V_x}{\partial x} . \quad (6)$$

After substituting (3) into equations (5), (6) we find:

$$-\omega^2 V_x = ik_x g V_z ,$$

$$-\omega^2 V_z = -ik_x g V_x .$$

This yields a dispersion equation for incompressible wave modes in the form

$$\omega^2 = k_x g . \quad (7)$$

Given the dispersion found, we obtain an expression for the polarization of the incompressible modes:

$$V_z = iV_x . \quad (8)$$

Further, from the condition (4) and polarization (8) we get $a = k_x$. Insofar as a is real value then non - divergent (ND) wave mode has no periodic vertical solution and is horizontally propagating.

Let us show that the dispersion relation (7) is also satisfied by another wave mode. After using this relation in (1), (2) we get:

$$V_x (\gamma H k_x - 1) - iV_z (1 - \gamma a H) = 0 , \quad (9)$$

$$iV_x (1 - \gamma H a - \gamma) - V_z \left(1 + \frac{\gamma H a^2}{k_x} - \frac{\gamma a}{k_x} \right) = 0 . \quad (10)$$

From the system (9), (10) follows:

$$a^2 - \frac{a}{H} + \frac{k_x}{H} (1 - k_x H) = 0 ,$$

which implies that there are two solutions to this equation:

$$a = k_x , \quad a = \frac{1}{H} - k_x . \quad (11)$$

The **first** solution in (11) corresponds to the non-divergent (ND) wave mode, and the **second** one we call pseudo-non-divergent mode (NDp). The expression for polarization NDp is obtained from (9) and has the form:

$$V_x \left(\frac{1}{\gamma H} - k_x \right) = -i \left(k_x - \frac{\gamma - 1}{\gamma H} \right) V_z .$$

Also for this mode holds the equation

$$\text{div} \vec{V} = \frac{V_z}{H} \frac{(1 - 2k_x H)}{(1 - \gamma k_x H)} ,$$

which shows that for NDp mode $\text{div} \vec{V} = 0$ only when $k_x = 1/2H$.

2.2 Anelastic and pseudo-anelastic modes

Let us show that equations (1), (2) indicate that another wave mode, not previously studied, may exist. To do this, we introduce, according to Bannon (1996), the anelastic linear perturbations, which satisfy the condition

$$\text{div}(\rho_0 \vec{V}) = 0 . \quad (12)$$

In the isothermal atmosphere with barometric density distribution we have

$$\frac{\partial \rho_0}{\partial z} = -\frac{\rho_0}{H} ,$$

therefore, for such *anelastic* perturbations, the following equation holds:

$$\text{div} \vec{V} = \frac{V_z}{H} . \quad (13)$$

Substituting (13) into equations (1), (2) we get:

$$\frac{\partial^2 V_x}{\partial t^2} = g(\gamma - 1) \frac{\partial V_z}{\partial x} ,$$

$$\frac{\partial^2 V_z}{\partial t^2} = -g(\gamma - 1) \frac{\partial V_x}{\partial x} .$$

Thus, given (3), should:

$$\omega^2 V_x = ik_x g(\gamma - 1) V_z , \quad (14)$$

$$\omega^2 V_z = -ik_x (\gamma - 1) g V_x . \quad (15)$$

Then the dispersion equation for *anelastic* (AE) *modes* takes the form:

$$\omega^2 = k_x g(\gamma - 1) . \quad (16)$$

With the resulting dispersion, polarization follows from equations (14), (15):

$$V_x = i V_z . \quad (17)$$

Further, taking into account (13), we obtain $a = \frac{1}{H} - k_x$. Consequently, the AE mode also does not have a solution periodic vertically and can only propagate horizontally.

After substituting the dispersion (16) into equations (1), (2) we get:

$$V_x (1 - \gamma + \gamma H k_x) - i V_z (1 - \gamma a H) = 0 , \quad (18)$$

$$iV_x(1-\gamma+\gamma Ha)+V_z\left(1-\gamma-\frac{\gamma Ha^2}{k_x}+\frac{\gamma a}{k_x}\right)=0, \quad (19)$$

whence we get a pair of values a , identical to (11). Consequently, there is another wave solution that satisfies the equation (16), we call it *pseudo-anelastic (AEp) mode*. **The first value in (11) corresponds to the AEp wave mode, and the second – to the AE one.**

Polarization of the AEp mode has the form:

$$V_x\left(k_x-\frac{\gamma-1}{\gamma H}\right)=-i\left(\frac{1}{\gamma H}-k_x\right)V_z$$

that follows from (18) or (19).

3 General properties of evanescent modes

Let us prove that the different types of evanescent modes characteristic of an isothermal atmosphere are related. We substitute (3) into system (1), (2) without additional conditions that were imposed in Section 2 when deriving ND and AE mode. As a result, we get:

$$V_z\left(a-\frac{g}{c^2}\right)-ik_xV_x\left(1-\frac{\omega^2}{k_x^2c^2}\right)=0, \quad (20)$$

$$V_x\left(a-\frac{N^2}{g}\right)-ik_xV_z\left(\frac{N^2}{\omega^2}-1\right)=0, \quad (21)$$

where $N^2 = \frac{g}{H} \frac{(\gamma-1)}{\gamma}$ is the square of the Brunt-Väisälä frequency.

From system (20), (21) we obtain the dispersion equation:

$$\left(\frac{1}{\gamma H}-a\right)\left(a-\frac{\gamma-1}{\gamma H}\right)=k_x^2\left(\frac{N^2}{\omega^2}-1\right)\left(1-\frac{\omega^2}{k_x^2c^2}\right). \quad (22)$$

Expressions $\omega^2 = N^2$ and $\omega^2 = k_x^2c^2$ are well-known dispersions of Brunt-Väisälä oscillations with $a = 1/\gamma H$ and Lamb waves (L) with $a = (\gamma-1)/\gamma H$. In addition to these known modes, dispersion (22) also admits the existence of additional solutions in the form of BV pseudo-modes (BVp) with $\omega^2 = N^2$, $a = (\gamma-1)/\gamma H$ and Lamb pseudo-modes (Lp) with $\omega^2 = k_x^2c^2$, $a = 1/\gamma H$ (Beer, 1974; Waltercheid and Hecht, 2003).

Then represent (22) in the form of a quadratic equation with respect to a :

$$a^2 - \frac{a}{H} + \frac{\omega^2}{c^2} - k_x^2 + \frac{k_x^2 g^2}{\omega^2 c^2} (\gamma - 1) = 0 .$$

The solution to this equation is:

$$a = \frac{1}{2H} \pm \sqrt{\frac{(k_x g - \omega^2)(\omega^2 - k_x g(\gamma - 1))}{\omega^2 c^2} + \left(k_x - \frac{1}{2H}\right)^2} . \quad (23)$$

From which it follows that for modes with dispersions $\omega^2 = k_x g(\gamma - 1)$ and $\omega^2 = k_x g$ there are two possible values: $a = k_x$ and $a = \frac{1}{H} - k_x$. The first value corresponds to modes ND and AEp, and the second - NDp and AE.

Thus, each evanescent mode can be associated with a pseudo-mode, which satisfies the same dispersion relation, but differs in polarization and dependence of the amplitude from the height, i.e., in its stratification. Table 1 presents the properties of different evanescent modes, characteristic of the isothermal atmosphere: BV oscillations, Lamb waves, non-divergent and anelastic modes, along with associated pseudo-modes: BVp, Lp, NDp, AEp. Table 1 shows that for all pseudo-modes, the polarization changes depending on the value of k_x . Wave modes AE and ND at $k_x = 1/2H$ completely coincide with AEp and NDp, respectively.

The location of the dispersion curves for anelastic and non-divergent modes relative to gravity and acoustic regions in the (ω, k_x) plane is shown in Fig. 1. The $\omega^2 = k_x g(\gamma - 1)$ mode touches the gravity region of freely propagating AGWs at the same value $k_x = 1/2H$ at which the $\omega^2 = k_x g$ curve touches the acoustic region (see Figure 1). In this case, the dispersion curves of AE and ND modes are symmetric relative to the ‘‘characteristic’’ curve (see Beer, 1974), which separates the AGW acoustic region from the AGW gravity region. In fact, the characteristic curve is the geometric mean of the dispersion curves of AE and ND modes with $\omega^2 = \sqrt{k_x^2 g^2 (\gamma - 1)} = N k_x c$.

From Figure 1 we see that the dispersion curves of different evanescent modes have intersection at separate points. Lamb dispersion curve with $\omega^2 = k_x^2 c^2$ intersects with BV curve with $\omega^2 = N^2$ at the point $k_x = N/c$. However, these modes cannot interact with each other by reason of different polarizations and values of a . At the same time, the pairs Lp - BV and L - BVp completely coincide in these properties and are indistinguishable at the intersection points.

Dispersion curves $\omega^2 = k_x g$ and $\omega^2 = k_x g(\gamma - 1)$ intersect with the Lamb curve and the BV curve at points $k_x = 1/\gamma H$, $k_x = (\gamma - 1)/\gamma H$. In addition, the ND mode curve intersects with the Lamb curve at the same value k_x , at which the AE mode curve intersects with the BV curve (see Fig. 1). ND and AE modes cannot interact with the Lamb mode and BV oscillations due to different polarizations (Table 1). Pseudo-modes NDp and AEp, at the points of intersection with the Lamb wave and the BV oscillations, have the same polarization and values of a . Similarly, ND and AE are indistinguishable at the points of intersection with Lp and BVp. Table 2 shows all evanescent modes that coincide with each other at the points of intersection of the dispersion curves, and between which interaction is possible. The cases of ND and AE mode curves intersection with curves ($a = 1/2H$), which separate the area of freely propagating AGWs from the evanescent area, are not presented in Table 2.

4 The energy of evanescent modes in an isothermal atmosphere

In Sections 2 and 3, we considered a model of an **unbounded** isothermal stratified atmosphere to determine which types of evanescent modes can satisfy the initial system of equations (1), (2). However, in an infinitely extended medium, the necessary condition for the existence of evanescent modes is the absence of unlimited growth of oscillation energy above and below the height level at which they are generated. It is easy to verify that in an isothermal infinite atmosphere, none of the modes listed in Table 1 satisfy this condition.

Suppose further that an evanescent wave is generated at a certain altitude level $z = 0$. The kinetic energy density $E \sim \rho(z)(V_x^2 + V_z^2)$ of waves should decrease both up and down from the level $z = 0$. When $z \rightarrow +\infty$ the energy density $E \sim \exp\left(2a - \frac{1}{H}\right)z \rightarrow 0$, if $a < 1/2H$, and $E \rightarrow \infty$, if $a > 1/2H$. When $z \rightarrow -\infty$ the energy density $E \rightarrow 0$, if $a > 1/2H$ and $E \rightarrow \infty$, if $a < 1/2H$. Based on these considerations, it is not difficult to understand how the energy density varies with height for different types of evanescent modes in an infinite isothermal atmosphere (see Table 3). Therefore, for the realization of such modes, it is necessary to have boundaries in the medium at which the condition for reducing energy in both directions from this boundary can be satisfied.

The presence of boundaries is not the only condition that can limit the energy of the evanescent mode. If the equality $a = 1/2H$ holds for these modes, then their energy is not varies with height in an isothermal atmosphere. For an infinite atmosphere, this solution does not seem to be physical, but it can make sense for a real atmosphere of finite height. As follows from (11), for the ND and AE modes, as well as their pseudo-modes, the condition $a = 1/2H$ performed at the point $k_x = 1/2H$. Also, at this point, the ND mode is identical to the NDp mode, and the AE mode completely coincides with AEp. In addition, when $k_x = 1/2H$ these evanescent modes adjoin the border of regions of freely propagating AGW (see Fig.1).

Consider some features of the energy balance for the evanescent modes. It follows from equation (20) that:

$$|V_z|^2 \left(a - \frac{g}{c^2} \right)^2 = k_x^2 |V_x|^2 \left(1 - \frac{\omega^2}{k_x^2 c^2} \right)^2. \quad (24)$$

Combining equations (22) and (24) gives the relation:

$$\rho_0 |V_x|^2 \left(1 - \frac{\omega^2}{k_x^2 c^2} \right) \left(a - \frac{N^2}{g} \right) = \rho_0 |V_z|^2 \left(\frac{N^2}{\omega^2} - 1 \right) \left(\frac{g}{c^2} - a \right). \quad (25)$$

The average density of the kinetic energy of the perturbations is $E_k = \frac{1}{4} \rho_0 (V_x^2 + V_z^2)$ and of the potential energy is $E_p = \frac{1}{4} \rho_0 \left(V_x^2 \frac{\omega^2}{k_x^2 c^2} + V_z^2 \frac{N^2}{\omega^2} \right)$ (Yeh, Liu, 1974; Fedorenko, 2010). Therefore, from equation (25) it follows that for the evanescent modes $E_k \neq E_p$. At the same time, for freely propagating AGWs is always fulfilled the equality $E_k = E_p$ (Yeh, Liu, 1974). At the point $a = 1/2H$ where evanescent modes on the plane (ω, k_x) in Fig. 1 are adjacent to areas of freely propagating AGWs, the equality $a - \frac{N^2}{g} = \frac{g}{c^2} - a$ holds. Taking this circumstance into account, from (25) we obtain:

$$\frac{\rho_0}{4} (|V_x|^2 + |V_z|^2) = \frac{\rho_0}{4} \left(|V_x|^2 \frac{\omega^2}{k_x^2 c^2} + |V_z|^2 \frac{N^2}{\omega^2} \right), \quad (26)$$

that is, at this point $E_k = E_p$.

5 Evanescent modes at the interface of isothermal media

Let us consider the possibility of realization of evanescent modes in the atmosphere at a thin interface between two isothermal half-spaces of infinite extent, which differ in temperature T . Let the boundary be localized at some altitude level $z=0$. In the lower half-space ($z < 0$) we have $T = T_1$, while in the upper half-space ($z > 0$) we have $T = T_2$ and it is assumed that $T_2 > T_1$. Note that a similar model was considered by Rosental and Gough (1994). We will search for solutions to the system (1), (2) in the form of $V_x, V_z \sim \exp(a_1 z) \exp[i(\omega t - k_x x)]$ for the lower half-plane and in the form $V_x, V_z \sim \exp(a_2 z) \exp[i(\omega t - k_x x)]$ for the upper half-plane. Substituting these dependencies into (1), (2) yields:

$$a_1 = \frac{1}{2H_1} \pm \left(\frac{1}{4H_1^2} - \frac{\omega^2}{c_1^2} + k_x^2 - k_x^2 \frac{N_1^2}{\omega^2} \right)^{1/2}, \quad (27)$$

$$a_2 = \frac{1}{2H_2} \pm \left(\frac{1}{4H_2^2} - \frac{\omega^2}{c_2^2} + k_x^2 - k_x^2 \frac{N_2^2}{\omega^2} \right)^{1/2}. \quad (28)$$

Here indices 1 and 2 denote the values in the lower and upper half-spaces, respectively.

The density of the kinetic energy of evanescent waves should decrease from the level $z=0$ both up and down. This condition limits the possible values of a_1 and a_2 . In the upper half-space ($z > 0$), when $z \rightarrow +\infty$ the energy density $E_2 \sim \exp\left(2a_2 - \frac{1}{H_2}\right)z \rightarrow 0$, if $a_2 < 1/2H_2$. In the lower half-space ($z < 0$), when $z \rightarrow -\infty$ the energy density $E_1 \sim \exp\left(2a_1 - \frac{1}{H_1}\right)z \rightarrow 0$, if $a_1 > 1/2H_1$. Therefore, it is necessary to take in the expression (27) for a_1 the solution with a “+” sign and in expression (28) for a_2 , with a “-” sign, so that the energy decreases on both sides of the interface.

It is also necessary to consider that the possible values of a_1 and a_2 must satisfy the boundary condition (Tolstoy, 1963; Rosental and Gough, 1994), arising from (1), (2):

$$\rho_1 c_1^2 \frac{gk_x^2 - \omega^2 a_1}{\omega^2 - c_1^2 k_x^2} \Big|_{z=0} = \rho_2 c_2^2 \frac{gk_x^2 - \omega^2 a_2}{\omega^2 - c_2^2 k_x^2} \Big|_{z=0}, \quad (29)$$

where ρ_1 and ρ_2 are the densities on both sides of the boundary. The procedure for deriving equality (29) is exactly the same as in the papers by Cheremnykh et al. (2018a) and Cheremnykh et al. (2018b). **When obtaining (29) we require continuity of the vertical velocity component (kinematic condition) and perturbed pressure (dynamic condition).** In the barometric atmosphere we have $\rho c^2 = \gamma p_0$, where p_0 is the equilibrium pressure, which must be continuous across the interface. Therefore, when $\gamma_1 = \gamma_2$ equation (29) can be written as:

$$\frac{gk_x^2 - \omega^2 a_1}{\omega^2 - c_1^2 k_x^2} = \frac{gk_x^2 - \omega^2 a_2}{\omega^2 - c_2^2 k_x^2} . \quad (30)$$

Dispersion dependencies of $\omega = f(k_x)$ calculated numerically by means of the expression (30) are shown in Fig. 2a for different values of the parameter $d = H_2 / H_1$. On each of these curves, the condition for decreasing energy up and down from the interface is satisfied. The long-wavelength part of the spectrum, where the most interesting features appear, is shown in more detail in Fig. 2b. Also shown in these figures are the dispersion curves $\omega = \sqrt{k_x g}$ and $\omega = \sqrt{k_x g(\gamma - 1)}$ for the ND and AE wave modes. The discontinuities of the $\omega = f(k_x)$ curves, as well as their cut-off for smaller k_x values, are due to requirements $a_1 > 1/2H_1$ and $a_2 < 1/2H_2$. Some features of the behavior of $\omega = f(k_x)$ will be discussed below.

As shown by Miles and Roberts (1992), the dispersion equation (30) can be rewritten to a polynomial form suitable for analysis:

$$\omega^8 - 2c_1^2(d+1)k_x^2\omega^6 + [c_1^4(d+1)^2k_x^4 + (2\gamma-1)k_x^2g^2]\omega^4 - 2(\gamma-1)c_1^2(d+1)k_x^4g^2\omega^2 - c_1^4(d-1)^2k_x^6g^2 = 0 . \quad (31)$$

Non-physical solutions (Miles and Roberts, 1992) arising from quadratic expressions under the radicals were excluded from consideration while obtaining equation (31) (see (27), (28)). Expressions (31) can be analyzed by examining their asymptotic behavior.

If $k_x^2 c_1^2 \gg \omega^2$, then from (31) we get:

$$\omega^4 - \frac{2N_1^2}{d+1}\omega^2 - \frac{(d-1)^2}{(d+1)^2}k_x^2g^2 \approx 0 .$$

It follows from this expression:

$$\omega^2 = \frac{1}{d+1} \left[N_1^2 + \sqrt{N_1^4 + (d-1)^2 k_x^2 g^2} \right]. \quad (32)$$

The expression (32) contains an interesting dependence of the frequency on the parameter d . In the limit $d \rightarrow \infty$, the dispersion $\omega^2 \approx k_x g$ of the ND (NDp) mode, independent of the properties of both environments, follows from (32). With $d \rightarrow 1$ and using (32), we obtain the dispersion of the BV (BVp) mode with the parameters of the lower medium, that is, $\omega^2 \approx N_1^2$. The indicated asymptotic features are visible on the curves shown in Fig. 2 below.

In the long-wave limit, i.e., at $k_x \rightarrow 0$, from (31) it follows:

$$(2\gamma - 1)\omega^4 - 2(\gamma - 1)c_1^2(d+1)k_x^2\omega^2 - c_1^4(d-1)^2k_x^4 \approx 0.$$

Hence we find:

$$\omega^2 = \frac{c_1^2 k_x^2}{2\gamma - 1} \left[(\gamma - 1)(d+1) + \sqrt{\gamma^2(d+1)^2 - 4d(2\gamma - 1)} \right]. \quad (33)$$

For the considered small k_x , for different values of d , from (33) we obtain the family of Lamb-type acoustic modes (see Fig. 2b). For large values of d , using (33), we obtain the expression $\omega^2 \approx c_1^2 k_x^2 d = c_2^2 k_x^2$, i.e. the oscillation frequency is determined by the characteristics of the medium in the upper half-space.

The evanescent modes frequencies lie on the (ω, k_x) plane between the acoustic and gravity regions of freely propagating AGW determined for upper and lower media separately (see Fig. 1). It is necessary to take into account when considering evanescent modes at the boundary of two isothermal media with different temperatures, that the evanescent regions are different in the upper and lower half-planes. On the (ω, k_x) plane, these regions are shifted relative to each other the more, the more is the value of d . At the same time, the wave modes at the interface of the media should remain evanescent in both media, and their dispersions should be enclosed within the overlap region of two evanescent regions. The cut-off curves for evanescent regions in the media under consideration are obtained in case of the null expressions under the radicals in (27) and (28). Gaps on the $\omega = f(k_x)$ dispersion curves are due to the evanescent areas of the two media do not match (see. Fig. 3).

Note that the dispersion curves $\omega = f(k_x)$ for values $d \leq 4$ are mostly inside both evanescent regions (see Fig. 3a, 3b), except for the longest waves. When $d \geq 4$, the dispersion curve $\omega = f(k_x)$ breaks into two separate branches (see Fig. 3 c , 3 d). The long-wave branch is acoustic, and another branch with $k_x \geq 0.4H_1$ is surface gravity by its physical nature.

6 Characteristic scales of ND and AE evanescent modes on the discontinuity

In an unlimited isothermal medium, evanescent modes are separate “pure” solutions of hydrodynamic equations. At the interface between two isothermal media with different temperatures, dispersion of the evanescent modes have a combined character, composing different types of “pure” modes, depending on the value of the parameter d and spectral properties $\omega(k_x)$.

For some values of d , the curves of the dispersion equation (30) approach fairly close to the curves $\omega^2 = k_x g$ and $\omega^2 = k_x g(\gamma - 1)$, and also intersect them at different points. These intersection points correspond to the specific value of k_x , at which the dispersions of the ND and AE modes are realized, in the model under consideration, in a “pure” form. Let us now examine these cases in more detail. For this purpose, we substitute the dispersion relations $\omega^2 = k_x g$ and $\omega^2 = k_x g(\gamma - 1)$ directly into (27), (28), and then into the boundary condition (30).

As was shown in Section 2, for dispersion relations $\omega^2 = k_x g$ and $\omega^2 = k_x g(\gamma - 1)$, values a_1 and a_2 coincide and are determined by expressions (11). Consider the valid values of a_1 and a_2 for these dispersions with regard to the requirement of energy decay in both directions from the interface $a_1 > 1/2H_1$ and $a_2 < 1/2H_2$.

6.1 Dispersion of the form $\omega^2 = k_x g$

For a dispersion of the form $\omega^2 = k_x g$, we first analyze the stratification of the ND mode with $a_1 = k_x$, $a_2 = k_x$. In order for the energy of this mode to decay in both directions from the discontinuity, the following inequalities $1/2H_1 < k_x < 1/2H_2$ must be satisfied, i.e., $H_1 > H_2$. Therefore, ND mode can be realized at the discontinuity, if the ambient temperature in the upper

region is less and the density is greater than they are in the lower region. This situation corresponds to the unstable state of the atmosphere (see Roberts, B., 1991).

Take the stratification of the NDp modes in the form of $a_1 = \frac{1}{H_1} - k_x$, $a_2 = \frac{1}{H_2} - k_x$. The energy in this case decreases both ways from the discontinuity, if $1/2H_2 < k_x < 1/2H_1$, i.e. when $H_2 > H_1$. This condition corresponds to the stable state and the case under consideration. For the NDp mode from the dispersion equation (30) we get:

$$H_2 \left(\frac{1}{\gamma H_2} - k_x \right) \left(2k_x - \frac{1}{H_1} \right) = H_1 \left(\frac{1}{\gamma H_1} - k_x \right) \left(2k_x - \frac{1}{H_2} \right), \quad k_x \neq 1/\gamma H_1, \quad k_x \neq 1/\gamma H_2. \quad (34)$$

From (34) it follows:

$$k_x = \frac{d+1}{4dH_1} \left(1 \pm \sqrt{1 - \frac{8d}{\gamma(d^2-1)}} \right). \quad (35)$$

Figure 4a shows values of k_x for which the dispersion curve $\omega^2 = k_x g$ intersects with the calculated dispersion curve $\omega = f(k_x)$ depending on the parameter d . The upper solid curve in this figure corresponds to the solution (35) with the sign “+” before the radical and shows the points of intersection with the shorter wavelength branch. The lower dashed curve corresponds to the solution with a sign “-” and represents the points of intersection with the long-wavelength branch. For the upper curve $k_x \rightarrow 1/2H_1$ when $d \rightarrow \infty$. For $d < 2.5$, there are no intersections of the curve $\omega = f(k_x)$ calculated numerically from (30) with the curve for the dispersion $\omega^2 = k_x g$.

When combining the stratifications for ND modes as $a_1 = k_x$ and for NDp modes as $a_2 = \frac{1}{H_2} - k_x$, equation (30) yields the only possible value of $k_x = 1/2H_2$. For a combination of stratifications $a_1 = \frac{1}{H_1} - k_x$ (NDp), $a_2 = k_x$ (ND) we get $k_x = 1/2H_1$. Both of these cases do not satisfy the condition of energy decrease with height.

Thus, consideration of the possible values of a_1 and a_2 leads to the conclusion that on the interface of two isothermal media with $H_2 > H_1$ can only be implemented NDp mode with a dispersion $\omega^2 = k_x g$ and a specific scale $k_x \sim 1/2H_1$.

6.2 Dispersion of the form $\omega^2 = k_x g(\gamma - 1)$

For the AE stratification of the form $a_1 = \frac{1}{H_1} - k_x$, $a_2 = \frac{1}{H_2} - k_x$ and for the AEp stratification of the form $a_1 = k_x$, $a_2 = k_x$, from the dispersion equation (30) follows the identity $H_1 = H_2$. Therefore, such modes do not realize at a temperature discontinuity. Apparently, to study the conditions of realization of AE and AEp modes, it is necessary to consider atmospheric models in which height profile $H(z)$ is continuous.

It should be noted that for the dispersion of the form $\omega^2 = k_x g(\gamma - 1)$, cases of combined modes stratifications are possible, satisfying the condition of decreasing energy on both sides of the boundary. So, for the combination of stratifications $a_1 = k_x$ (AEp), $a_2 = \frac{1}{H_2} - k_x$ (AE) from (30) we obtain the relation:

$$H_2 k_x (2 - \gamma) = H_1 \gamma \left(k_x - \frac{\gamma - 1}{\gamma H_1} \right).$$

Whence $k_x = \frac{\gamma - 1}{H_1 [\gamma - (2 - \gamma)d]}$. In this case, the inequality $d < \gamma / (2 - \gamma)$ must be satisfied. When $\gamma = 5/3$ we get the following restriction: $d < 5$. Given this limitation and condition $k_x > 1/2dH_1$, we obtain that a mode with a dispersion of $\omega^2 = k_x g(\gamma - 1)$ and stratification of AE type for the upper half-space and of AEp type for the lower half-space can propagate at the boundary in the range $1 < d < 5$ and for $k_x > 1/2H_1$. For the stratifications $a_1 = \frac{1}{H_1} - k_x$ (AE), $a_2 = k_x$ (AEp) from equation (30) we obtain the relation:

$$H_2 \gamma \left(k_x - \frac{\gamma - 1}{\gamma H_2} \right) = H_1 k_x (2 - \gamma).$$

It implies the ratio $k_x = \frac{\gamma - 1}{H_1 [\gamma d - (2 - \gamma)]}$, in which the parameter d can take any values with $d > 1$, and the horizontal wave number is limited by the inequality $k_x < 1/2H_1$. Features of the behavior of the $\omega^2 = k_x g(\gamma - 1)$ mode at the discontinuity, depending on the scale k_x are shown in Fig. 4b.

7 Discussion

Let us dwell on some of the results in terms of their use for the analysis of experimental data.

With the f -mode observed on the Sun, one should identify the mode that we classify as ND mode, for which $\omega^2 = k_x g$, $V_z \sim \exp(k_x z)$ and $\text{div}\vec{V} = 0$ (Roberts, B., 1991). In the framework of the considered temperature discontinuity model, it was shown that with $T_1 < T_2$ (corresponds to the chromosphere-corona interface) the condition for decreasing amplitude with height to both sides of the interface is satisfied only by the NDp mode with $\omega^2 = k_x g$, $V_z \sim \exp\left(\frac{1}{H} - k_x\right)z$ and $\text{div}\vec{V} \neq 0$.

When the ratio $d \rightarrow \infty$ (i.e., $H_2/H_1 \rightarrow \infty$), the NDp mode with $k_x \rightarrow 1/2H_1$ asymptotically approaches ND mode. On the interface between the chromosphere and the solar corona d is large, but of finite magnitude: $d \sim 50$ (Athay, 1976; Jones, 1969). Therefore, the condition of the presence of a free surface, which is required for the realization of the ND mode, is fulfilled only approximately. Therefore, in the framework of the temperature discontinuity model, the f -mode observed on the Sun should not be **associated** with the non-divergent ND mode, but with non-divergent *pseudo*-mode NDp.

For the Earth's atmosphere, the maximum possible value of d is observed at the interface between the thermosphere with $T_2 \sim 800 - 1500K$ (depending on solar activity) and the underlying atmosphere with $T_1 \sim 300K$. When $d = 5$, the dispersion (30) asymptotically tends to $\omega^2 = k_x g(\gamma - 1)$ with $k_x \rightarrow \infty$. Therefore, it can be expected that evanescent modes in this case will be close to $\omega^2 = k_x g(\gamma - 1)$.

In other layers of the earth's atmosphere we have $d \leq 1.3$ (Jursa, 1985). As follows from (33), for small values of $d \leq 1.3$ and for the wavelengths in the interval $k_x \sim (0.5 - 1.5)H_1$, the **relation** $\omega^2 \rightarrow N^2$ is satisfied (see Fig. 2). Therefore, it can be expected that at small positive temperature gradients in the atmosphere, waves with a frequency close to the frequency of Brent-Väisälä should prevail. These conclusions experimentally confirm (Shimkhada et al., 2009) the results of observations of short-period evanescent waves with small wavelengths at altitudes near the mesopause.

8 Main results

In the paper, different types of evanescent acoustic-gravity modes characteristic of an isothermal atmosphere are investigated. A new mode was derived in the form of anelastic acoustic-gravity wave mode with the dispersion equation $\omega^2 = k_x g (\gamma - 1)$. The main properties of the AE mode are presented in Table 1 in comparison with other known evanescent modes. It is shown that for both anelastic and non-divergent modes there are pseudo-modes that satisfy the same dispersions, but having different polarization and the dependence of the amplitude of the disturbances on the height.

For AE and ND evanescent modes, the value of $k_x \rightarrow 1/2H$ sets a special scale (wavelength) at which these modes are identical to their pseudo-modes AEp and NDp. In addition, at the same point they are adjacent to the boundaries of the continuous spectrum (AE mode to the gravity region, and ND mode to the acoustic region, respectively).

The features of the evanescent modes realization at the interface of two isothermal media are considered. It is shown that in this case, dispersions of evanescent modes are combined, merging the features of different types of modes characteristic of an **unbounded** isothermal atmosphere. This effect is most pronounced in the following asymptotic cases: 1) when $d \rightarrow \infty$, we obtain the dispersion for the ND (NDp) mode in the form $\omega^2 \approx k_x g$; 2) when $d \rightarrow 1$, for scales $k_x \sim H_1$, a mode with $\omega^2 \approx N_1^2$ is realized; 3) for $k_x \rightarrow 0$, a Lamb wave with a dispersion relation of the form $\omega^2 \approx c_2^2 k_x^2$ is obtained, which depends only on the parameters of the medium in the upper half-space.

It was demonstrated that on the interface of two isothermal media with $T_2 > T_1$, the NDp mode with the dispersion $\omega^2 = k_x g$ and the selected scale $k_x \sim 1/2H_1$ is realized. At the same time, the ND mode does not satisfy the condition of decreasing energy on each side of the interface. Dispersion $\omega^2 = k_x g (\gamma - 1)$ on the interface of two media is satisfied by the wave mode, which has different types of amplitude versus height dependencies at different horizontal scales k_x . When $k_x > 1/2H_1$, the height dependence of AE amplitude for $z > 0$ and AEp amplitude for $z < 0$ satisfy the condition of decreasing energy from the interface. On the contrary, when $k_x < 1/2H_1$, this condition is satisfied by AEp amplitude for $z > 0$ and AE amplitude for $z < 0$.

It is important to note that according to our analysis in the framework of the temperature discontinuity model: (1) the f - mode observed on the Sun should not be **associated** with the non-

divergent ($\omega^2 = k_x g, \text{div}V = 0$) mode, but with its non-divergent *pseudo*-mode ($\omega^2 = k_x g, \text{div}V \neq 0$). (2) At the interface between the earth's thermosphere and the underlying atmosphere it can be expected that evanescent modes with short wavelengths will be close to the new mode ($\omega^2 = k_x g(\gamma - 1)$). (3) Oscillations with a frequency close to the frequency of Brent-Väisälä should prevail at altitudes near the earth's mesopause.

***Acknowledgements.* This publication is based on work supported in part by Integrated Scientific Programmes of the National Academy of Science of Ukraine on Space Research and Plasma Physics.**

References

- Antia, H.M.: Estimate of solar radius from f-mode, *Astron. Astrophys.*, 330, 336-340, 1998.
- Athay, R.G.: *The Solar Chromosphere and Corona: Quiet Sun*. Dordrecht, Holland: Reidel, 504, 1976.
- Beer, T.: *Atmospheric Waves*, John Wiley, New York, 300 pp, 1974.
- Belashov, V.Yu.: Dynamics of nonlinear internal gravity waves at ionosphere F-region heights, *Geomagn. Aeron. (Engl. Transl.)*, 1990. 30, 637-641, 1990.
- Bannon, P.R.: On the anelastic approximation for a compressible atmosphere, *J. Atmos. Sci.*, 53, 23, 3618 – 3628, 1996.
- Cheremnykh, O., Cheremnykh, S., Kozak, L., Kronberg, E.: Magnetohydrodynamics waves and the Kelvin_Helmholtz instability at the boundary of plasma mediums, *Phys. Plasmas*, 25, 10, 102119, 2018a.
- Cheremnykh, O.K., Fedun, V., Ladikov-Roey, Y.P., Verth, G.: On the stability of incompressible MHD modes in magnetic cylinder with twisted magnetic field and flow, *Astrophys. Journ.*, 866, 2, 86, 2018b.
- Cheremnykh, O.K., Selivanov, Yu.A., Zakharov, I.V.: The influence of compressibility and non-isothermality of the atmosphere on the propagation of acoustic-gravity waves, *Kosm. nauka tehnol.*, 16, 1, 09–19, doi.org/10.15407/knit2010.01.009, 2010.

- Fedorenko, A.K.: Energy Balance of Acoustic Gravity Waves above the Polar Caps According to the Data of Satellite Measurements, *Geomagn. Aeron.* 50, 107–118, 2010.
- Fedorenko, A.K., Kryuchkov, Y.I.: Wind Control of the Propagation of Acoustic Gravity Waves in the Polar Atmosphere, *Geomagn. Aeron. (Engl. Transl.)* 53, 377-388, 2013.
- Fedorenko, A.K., Bespalova, A.V., Cheremnykh, O.K., Kryuchkov, E.I.: A dominant acoustic-gravity mode in the polar thermosphere, *Ann. Geophys.*, 33, 101-108, 2015. [doi:10.5194/angeo-33-101-2015].
- Fedorenko, A.K., Kryuchkov, E.I., Cheremnykh, O.K., Klymenko, Yu.O., Yampolski Yu.M.: Peculiarities of acoustic-gravity waves in inhomogeneous flows of the polar thermosphere, *J. Atmos. Solar-Ter. Phys.*, 178, 17-23, 2018. doi.org/10.1016/j.jastp.2018.05.009
- Francis, S.H.: Global propagation of atmospheric gravity waves: A review, *J. Atmos. Terr. Phys.*, 37, 1011-1054, 1975.
- Ghosh, P., Antia, H.M., Chitre, S.M.: Seismology of the solar f-mode. I. Basic signatures of shearing velocity fields, *Astrophys. J.*, 451, 851-858, 1995.
- Hines, C.O.: Internal gravity waves at ionospheric heights, *Can. J. Phys.*, 38, 1441–1481, 1960.
- Huang, K.M., Zhang, S.D., Yi, F., Huang, C.M., Gan, Q., Gong, Y., Zhang, Y. H.: Nonlinear interaction of gravity waves in a nonisothermal and dissipative atmosphere, *Ann. Geophys.*, 32, 263-275, 2014. [doi:10.5194/angeo-32-263-2014].
- Innis, J.L., Conde, M.: Characterization of acoustic-gravity waves in the upper thermosphere using Dynamics Explorer 2 Wind and Temperature Spectrometer (WATS) and Neutral Atmosphere Composition Spectrometer (NACS) data, *J. Geophys. Res.*, 107, NoA12, 2002. [doi: 10.1029/2002 JA009370].
- Johnson, F.S., Hanson, W.B., Hodges, R.R., Coley, W.R., Carignan, G.R., Spencer, N.W.: Gravity waves near 300 km over the polar caps, *J. Geophys. Res.*, 100, 23993-24002, 1995.
- Jones, Walter L.: Non-divergent oscillations in the Solar Atmosphere, *Solar Phys.*, 7, 204-209, 1969.
- Jursa, Adolph S.: Handbook of geophysics and the space environment. Vol. 1. Hanscom Air Force Base, MA: Air Force Geophysics Laboratory, Air Force Systems Command, United States Air Force, 1040, 1985.

- Kaladze, T.D., Pokhotelov, O.A., Shah, H.A., Khan, M.I., Stenflo, L.: Acoustic-gravity waves in the Earth's ionosphere, *J. Atmos. Solar-Terr. Phys.*, 70, 1607–1616, 2008.
- Ladikov-Roev, Y.P., Cheremnykh, O.K.: *Mathematical Models of Continuous Media*, Naukova Dumka, Kyiv, 552, 2010 (in Russian).
- Miles, Alan, J., Roberts, B.: Magnetoacoustic-gravity surface waves. I. Constant Alfvén Speed, *Solar Phys.*, 141, 205-234, 1992.
- Nekrasov, A.K., Shalimov, S.L., Shukla, P.K., Stenflo, L.: Nonlinear disturbances in the ionosphere due to acoustic gravity waves, *J. Atmos. Terr. Phys.*, 57, 732–742, 1995.
- Roberts, B.: MHD Waves in the Sun, in: *Advances in solar system magnetohydrodynamics*, CUP, 239, 1991.
- Rosental, C.S., Gough, D.O.: The Solar f-mode as interfacial mode at the chromosphere-corona transition. *The Astrophysical Journal*, 423, 488-495, 1994.
- Simkhada, D.B., Snively, J.B., Taylor, M.J., Franke, S.J.: Analysis and modeling of ducted and evanescent gravity waves observed in the Hawaiian airglow. *Ann. Geophys.* 27, 3213-3224, 2009.
- Stenflo, L., Shukla, P. K.: Nonlinear acoustic-gravity waves, *J. Plasma Phys.*, 75, 841–847, 2009. [doi.org/10.1017/S0022377809007892].
- Tolstoy, I.: The theory of waves in stratified fluids including the effects of gravity and rotation, *Rev. of Modern Phys.*, Vol. 35, N1, 1963.
- Vadas, S. L., Nicolls, M. J.: The phases and amplitudes of gravity waves propagating and dissipating in the thermosphere: Theory, *J. Geophys. Res.*, 117, A05322, doi:10.1029/2011JA017426, 2012.
- Waltercheid, R.L., Hecht, J.H.: A reexamination of evanescent acoustic-gravity waves: Special properties and aeronautical significance, *J. Geophys. Res.*, 108, D11, 4340, doi:10.1029/2002JD002421, 2003.
- Yeh, K.S., Liu, C.H.: Acoustic-gravity waves in the upper atmosphere, *Rev. Geophys. Space. Phys.*, 12, 193-216, 1974.

Table 1. Properties of different evanescent acoustic-gravity modes

Mode type	Dispersion	a	Polarization
Lamb Wave (L)	$\omega^2 = k_x^2 c^2$	$\frac{\gamma-1}{\gamma H}$	$V_z = 0; V_x \neq 0$
Lamb's Pseudo-mode (Lp)		$\frac{1}{\gamma H}$	$V_x(2-\gamma)k_x g = i(N^2 - k_x^2 c^2)V_z$
BV Oscillations (BV)	$\omega^2 = N^2$	$\frac{1}{\gamma H}$	$V_x = 0; V_z \neq 0$
BV Pseudo-mode (BVp)		$\frac{\gamma-1}{\gamma H}$	$V_x(k_x^2 c^2 - N^2) = i(2-\gamma)k_x g V_z$
Non-divergent (ND) mode, $div \vec{V} = 0$	$\omega^2 = k_x g$	k_x	$V_x = -iV_z$
Pseudo-non-divergent mode (NDp), $div \vec{V} \neq 0$		$\frac{1}{H} - k_x$	$V_x \left(\frac{1}{\gamma H} - k_x \right) = -i \left(k_x - \frac{\gamma-1}{\gamma H} \right) V_z$
Anelastic mode (AE), $div(\rho_0 \vec{V}) = 0$	$\omega^2 = k_x g(\gamma-1)$	$\frac{1}{H} - k_x$	$V_x = iV_z$
Pseudo-anelastic mode (AEp), $div(\rho_0 \vec{V}) \neq 0$		k_x	$V_x \left(k_x - \frac{\gamma-1}{\gamma H} \right) = i \left(\frac{1}{\gamma H} - k_x \right) V_z$

Table 2. The coincidence of the evanescent mode properties at the intersection points of the dispersion curves *

Lamb Wave (L)	Lamb's Pseudo-mode (Lp)	BV Oscillations (Bv)	BV Pseudo-mode (Bvp)	Non-divergent mode (ND)	Pseudo-non-divergent mode (NDp)	Anelastic mode (AE)	Pseudo-anelastic mode (AEp)
BVp NDp AEp	BV ND AE	Lp NDp AEp	L ND AE	Lp BVp	L BV	Lp BVp	L BV

* *Note.* The bottom rows show the modes that are indistinguishable from the corresponding mode of the top row at the point of intersection of the dispersion curves.

Table 3. The change in energy density of evanescent modes with height in an infinite isothermal atmosphere

Domain	Lamb Wave (L)	Lamb's Pseudo-mode	BV Oscillations (BV)	BV Pseudo-mode	Non-divergent mode (ND)	Pseudo-non-divergent mode (NDp)	Anelastic mode (AE)	Pseudo-anelastic mode (AEp)
$z \rightarrow +\infty$	$E \rightarrow 0$	$E \rightarrow \infty$	$E \rightarrow \infty$	$E \rightarrow 0$	$E \rightarrow 0,$ $k_x < 1/2H$ $E \rightarrow \infty,$ $k_x > 1/2H$	$E \rightarrow 0,$ $k_x > 1/2H$ $E \rightarrow \infty,$ $k_x < 1/2H$	$E \rightarrow 0,$ $k_x > 1/2H$ $E \rightarrow \infty,$ $k_x < 1/2H$	$E \rightarrow 0,$ $k_x < 1/2H$ $E \rightarrow \infty,$ $k_x > 1/2H$
$z \rightarrow -\infty$	$E \rightarrow \infty$	$E \rightarrow 0$	$E \rightarrow 0$	$E \rightarrow \infty$	$E \rightarrow \infty,$ $k_x < 1/2H$ $E \rightarrow 0,$ $k_x > 1/2H$	$E \rightarrow \infty,$ $k_x > 1/2H$ $E \rightarrow 0,$ $k_x < 1/2H$	$E \rightarrow \infty,$ $k_x > 1/2H$ $E \rightarrow 0,$ $k_x < 1/2H$	$E \rightarrow \infty,$ $k_x < 1/2H$ $E \rightarrow 0,$ $k_x > 1/2H$




Fig. 1. Dispersion dependencies $\omega = f(k_x)$: 1) boundaries between acoustic and gravity regions for freely propagating waves (dashed lines); 2) evanescent mode: $\omega = \sqrt{k_x g}$ (upper solid curve) and $\omega = \sqrt{k_x g(\gamma - 1)}$ (lower solid curve), $\omega = N$ (thin horizontal line), $\omega = k_x c$ (thin sloping straight line)




Fig. 2. Dispersion dependencies $\omega = f(k_x)$ at the boundary of the discontinuity for different values of the parameter d . General dependence (a), long-wave part in more detail (b). Thin curves denote $\omega = \sqrt{k_x g}$ (upper curve) and $\omega = \sqrt{k_x g(\gamma - 1)}$ (lower curve), $\omega = N$ (horizontal line)




Fig. 3. Dispersion dependencies of the $\omega = f(k_x)$ type at the temperature discontinuity boundary for $d = 2$ (a), $d = 3$ (b), $d = 5$ (c), $d = 20$ (d). **The dashed curves represent the boundaries of the areas with free propagation of AGW in the upper and lower half-space. Thin curves denote $\omega = \sqrt{k_x g}$ (upper curve) and $\omega = \sqrt{k_x g(\gamma - 1)}$ (lower curve)**




Fig. 4. Horizontal scales $k_x H_1$, on which the modes with the dispersion $\omega^2 = k_x g$ (a) and $\omega^2 = k_x g(\gamma - 1)$ (b) are realized, depending on $d = H_2 / H_1$