REPLY TO EDITOR

Reply: The authors reviewed the text according to the suggestions and comments proposed by the Referees. In addition, we highlight the modifications in the manuscript to be followed in the review process. Lastly, we would like to take this opportunity to thank the editor for considering this article for publication in Annales Geophysicae.

REPLY TO REFEREE#1 (Thana Yeeram)

Reviewer comments for authors:

I have read the manuscript that reports the responses of TEC to HILDCAA intervals over the two Brazilian GNSS stations. The manuscript is fairly good in presentation, particularly in reporting an equinoctial anomaly of the TEC during HILDCAAs. As known, this topic has not been extensively studied and is in progress in the field. Since some of issues in the present forms are not adequately explained for the underlying Physics, I decided a minor revision for this manuscript.

Reply: We would like to thank the useful suggestions and comments given by the Referee to improve this paper.

In what follows, we describe how we revised the manuscript and provided the answers to the specific and technical comments of the Referee. We try to make the manuscript clearer and more consistent with what was actually done in this work. We would like to take this opportunity to thank the Referee for his contribution to improving this work.

Specific comment:

1. The abstract should be written in a concise form for the lines 32 - 39.

Reply: We would like to thank the Reviewer for this suggestion. We modified the abstract in order to let it more concise.

2. I think that one of the HILDCAA's criteria is there are HSS and high frequency fluctuations of IMF Bz about zero value.

Reply: Actually, the criteria of the fluctuation of IMF Bz around zero was added in a later study carried out by Koga et al. 2011 (doi: 10.1016/j.jastp.2010.09.002). According to Tsurutani et al., 2004 (doi: 10.1016/j.jastp.2003.08.015) the same physical process may occur weather one criteria is not strictly followed.

3. In lines 65 - 67, the authors may refer to n° 2 for HILDCAAs' properties as well.

Reply: We would like to thank the Referee for this suggestion. However, as the information written in the lines 65-67 is well-known, we have preferred to let it in this way.

4. In lines 74 - 75, the authors should mention the references.

Reply: Thank you for the suggestion. We added the reference.

5. In section Data and Methodology, the authors should remove the links of data since they are already in the acknowledgement.

Reply: Thanks for your suggestion. We removed the link of data from the Data and Methodology section and left it in the Acknowledgement section.

6. In line 218 please give more details about the related mechanism for the equinoctial anomaly.

Reply: Information about the equinoctial anomaly is already written in the manuscript in lines 223 - 225. For more details, please see Balan et al., 1998 (doi:10.1029/97JA03137), Mansilla et al., 2005 (doi:10.1016/j.jastp.2005.02.024) and Chen et al., 2012 (doi:10.5194/angeo-30-613-2012).

7. In section 3.3 why the solar wind speed is thought to be a main factor that affects the TEC?

Reply: We would like to thank the referee for this question. We realize that several papers address the relation between HILDCAAs and HSS/CIRs. However, how this relation is done is still an open question. Contrary to our expectations, the TEC answers during HILDCAAs are no direct relation to fast speeds.

- 8. I would like to suggest some related work that may fulfill the discussion: For recent study of TEC and HILDCAAA:
- de Siqueira Negreti, P. M., de Paula, E. R., and Candido, C. M. N.: Total electron content responses to HILDCAAs and geomagnetic storms over South America, Ann. Geophys., 35, 1309–1326, https://doi.org/10.5194/angeo-35-1309-2017, 2017.
 For PPEF and DDEF during HILDCAA:
- Yeeram, T. (2019). The solar wind magnetospheric coupling and daytime disturbance electric fields in equatorial ionosphere during consecutive recurrent geomagnetic storms, Journal of Atmospheric and Solar-Terrestrial Physics, 187, 40-52.
- Yeeram, T., and Paratrasri, A. (2018). Recurrent geomagnetic storms and equinoctial ionospheric F-region in the low magnetic latitude: a case study, IOP Conf. Series: Journal of Physics: Conf. Series 1144: 012024(1-4).

Reply: We would like to thank the Referee for the suggestions. We read them and added the citation in lines 75, 95 and 237.

9. Line 251, the geoeffectiveness of HILDCAA can be different and separated from CIR. Please see Hajra et al. 2015. Relativistic electron acceleration during HILDCAA events: are precursor CIR magnetic storms important?

Reply: Yes, the geoeffectiveness of HILDCAAs can be separated from CIRs, since to a lesser extent; there are HILDCAA events related to CME.

10. The conclusions should be written in a short and concise form.

Reply: We would like to thank the suggestion. The main results were written in topics in the Conclusions section.

11. I do not understand the x-axis of Figures 2 - 5. Why the scale is for one day and for what? I see HILDCAA intervals are longer than that for each event H? Please describe in the text.

Reply: The scale is not for one single day. The x-axis in Figures 2, 3 and 4 represent the mean dTEC hourly values. For Figures 2 and 3, the mean values were represented in this way in order to allow the comparison between one interval and other (please, see the lines 147 - 148). For Figure 4 the x-axis represents the central tendency for all dTEC values, minute-to-minute, as can be seen in lines 215 - 218.

Lastly, in Figure 5, the x-axis represents the time duration of all intervals, represented each 12 hours.

The technical comments:

 Some of the English must be corrected. For example, line 251, gerund "Summarizing" should be →In summary/In conclusion.

Reply: We would like to thank the Referee for this correction. The word was changed in the manuscript.

2. The sentences must be simple and concise.

Reply: We are very thankful to the Referee to help us improve the manuscript.

REPLY TO REFEREE#2

Reviewer comments for authors:

This work deals with a topical issue, which is space weather and its effect on the Ionosphere. However I cannot see which is the real message. If I am right, I understand that the authors want to show that HILDCAAs affect TEC at low and equatorial latitudes. But in my opinion, they need to show that what they find is not something at random, and to exclude any other source of disturbance. I think that the paper may become acceptable for publication after some revision.

Reply: We would like to thank useful suggestions and comments given by the Referee to improve this paper.

In what follows, we describe how we revised the manuscript and provided the answers to the main and minor comments of the Referee. We try to make the manuscript clearer and more consistent with what was actually done in this work. Again, we would like to take this opportunity to thank the Referee for his/her contribution to improving this work.

Main comments:

1. I would add some statistical analysis, or qualitative analyses, to show that the TEC disturbances they observe are due to HILDCAAs; by showing that variation they see is not at random and is not due to any other mechanism. Is it possible that it is due to geomagnetic storms, for example? Maybe I am not fully understanding HILDCAAs.

Reply: Unfortunately, do a statistical analysis with ten HILDCAA events as a sample is not significant. However, we believe that carried out a qualitative analysis when we seek to comprise the HILDCAAs disturbances regards seasonal behavior or local time, for example. All TEC disturbances showed in the study is about HILDCAAs influences, since all analysis was done taking into account the dTEC (dTEC = TEC mean – TEC quiet days). Please, see lines 125 - 128.

HILDCAAs are manifestations of space weather events in the form of geomagnetic activities. Their main feature is a continuous flux of energetic particles into the magnetosphere. The average Dst remain suppressed for days or weeks, and appear as an

unnaturally long storm recovery phase. For more, please see Tsurutani and Gonzalez, 1987 (doi: 10.1016/0032-0633(87)90097-3), Tsurutani et al., 2004 (doi: 10.1016/j.jastp.2003.08.015), Hajra et al., 2013 (doi: 10.1002/jgra.50530) and Silva et al., 2017 (doi: 10.5194/angeo-35-1165-2017).

2. Line 165: "The HILDCAA intervals present the positive dTEC predominance" can you quantify this?

Reply: We would like to thank the Referee for this suggestion. We realized that the sentence was short and no concise. To improve the understanding, we have adding the following sentences (lines 168 - 170):

"60% (70%) of all intervals present a positive dTEC response during the whole event for São Luís (Cachoeira Paulista)."

The percentage is regarding the predominance of the positive dTEC during the whole hourly distribution. Only four events for São Luís (three for Cachoeira Paulista) had negative predominance of the TEC behavior.

3. In Figures 2 and 3: what is the "y" axis?

Reply: Figures 2 and 3 are the subplots of all HILDCAA intervals regarding mean dTEC hourly distribution, i. e., each panel refer to each interval. The y-axis is the HILDCAA interval according to its identification (H01, H02, etc).

However, the y-axis legend was added to the manuscript in order to avoid misinterpret.

4. In Figure 5, for H03 and H04 a clear daily variation can be noticed. This 24-hour oscillation is absent in the others. What does this mean?

Reply: This result surprised us! Figure 5 shows the solar velocity for each one of the HILDCAA intervals, where intervals H03 and H04 are representative for the autumn season in the South hemisphere (Table 1). Figure 4 also clearly shows the 24-hour oscillation (top left panel).

The behavior is very similar to the daily oscillation, where the maximum density occurs during the daytime, while the minimum density occurs at night. However, as has been said, this behavior is already subtracted from the quiet day pattern. Our hypothesis is that because geomagnetic activity during HILDCAAs events remains high, continuous injection of energetic particles can cause prolonged changes in the global wind circulation, causing the ionospheric F layer to remain elevated during the day causing increases in ionization.

However, we would like to emphasize that only two events occurring in each season were chosen for this study, and therefore further studies on this specific behavior of dTEC during HILDCAA intervals will need to be performed.

Minor comments:

 Line 53: "The", should be lower-case, that is "the", and also in the following "The" which appear after a semi colon.

Reply: We would like to thank the Reviewer for this note. The words were corrected in the manuscript.

2. Line 67: I thinking that may be "differential" should be "different".

Reply: We would like to thank the Reviewer for this suggestion. The word was changed to the manuscript. Please, check the line 67.

3. Line 97: "taking account" should be "taking into account".

Reply: We would like to thank the Referee for this observation. The sentence was corrected. Please, see the line 97.

3. Line 118: use dot for decimal of latitude and longitude instead comma. That is "2,59" should be 2.59.

Reply: We would like to thank the Referee for this correction.

4. Line 153: Why "to São Luís". Shouldn't it be "for São Luís" or at São Luís? The same in the case of Cachoeira Paulista.

Reply: The sentence was corrected changing "to" to "for". Please, verify the lines 154 - 155.

5. Line 154: What do you means by "It was considered the same minimum and maximum values occurred to all intervals, for each station." That you used the same scale range? If this is the case, then you should explain it properly.

Reply: Yes, we used the same scale range for all HILDCAA intervals observed in each GNSS station. We seek the maximum and minimum values to use as a pattern to analyze the hourly distribution. Doing this way, it allows us to compare events with each other.

Following the Referee's suggestion, the below sentences were added to the paper (lines 155 - 158):

"These values were considered to perform the TEC hourly distribution, i. e., for each specific GNSS station, the maximum and minimum TEC values were used to analyze all HILDCAAs in the same range."

6. Line 162: "...phases that over equatorial latitudes", may be you mean "...phases over equatorial latitudes"?

Reply: We would like to thank the Referee for this observation.

7. Line 228: "There are considerable works whose show" should be "There are considerable works that show..."

Reply: We would like to thank the Referee for this observation. The sentence was corrected in the manuscript.

1	Ionospheric Total Electron Content responses to HILDCAAs intervals
2	R. P. Silva ^{*1,2} , C. M. Denardini ¹ , M. S. Marques ³ , L. C. A. Resende ^{1, 4} , J. Moro ^{4, 5} , G.
3	A. S. Picanço ¹ , G. L. Borba ³ , M. A. F. Santos ² .
4	
5	
6	¹ National Institute for Space Research – INPE, São José dos Campos-SP, Brazil.
7	² Northeast Regional Center – CRN/INPE, Natal-RN, Brazil.
8	³ Geophysics Department (DGEF), Federal University of Rio Grande do Norte
9	(UFRN), Natal-RN, Brazil.
10	⁴ National Space Science Center, China Academy of Science, CAS, Beijing, China;
11	⁵ Southern Regional Space Research Center – CRS/COCRE/INPE, Santa Maria-RS,
12	Brazil.
13	
14	
15	
16	*Corresponding author: INPE/DIDAE, Av. dos Astronautas, 1758, Jd. da Granja,
17	12227-010, São José dos Campos, SP, Brazil; Tel.: +55 12 3208-7184; e-mail:
18	regia.pereira@inpe.br; regiapereira@gmail.com
19	
20	

22 Abstract

23 The High-Intensity Long-Duration and Continuous AE Activities (HILDCAA) 24 intervals are capable of causing a global disturbance in the terrestrial ionosphere. 25 However, the ionospheric storms' behavior due to these intervals geomagnetic 26 activity forms is still not widely understood. In the current this study, we seek to 27 comprise the HILDCAAs disturbance time effects in the Total Electron Content 28 (TEC) values with respect to the quiet days' pattern analyzing local time and seasonal 29 dependences, and the influences of the solar wind velocity to a sample of ten intervals 30 occurred in 2015 and 2016 years. The main results showed that the hourly distribution 31 of the disturbance TEC may vary substantially between one HILDCAA interval and 32 another. Doing a comparative to geomagnetic storms, while the positive ionospheric 33 storms are more pronounced in the winter, this season presents less geoeffectiveness 34 or almost none to HILDCAA intervals. It was find found an equinoctial anomaly, 35 since the equinoxes represent more ionospheric TEC responses during HILDCAA 36 intervals than the solstices. Regarding to the solar wind velocities, although HILDCAA intervals are associated to with High Speed Streams, this association does 37 38 not present a direct relation regards to TEC disturbances magnitudes in low and 39 equatorial latitudes.

40

41

- 42
- 43
- 44

45 Keywords: HILDCAA, TEC, Equatorial Ionosphere

47 **1. Introduction**

As similar to geomagnetic storms, High-Intensity Long-Duration and Continuous AE Activities (HILDCAA) intervals can influence the ionosphere, leading to disturbances in the ionospheric F2-region. It is well known that these intervals can change the F2region peak height being, generally, less intense than those observed during typical geomagnetic storm events (Sobral et al., 2006; Koga et al., 2011, Silva et al., 2017).

53 In fact, HILDCAAs are characterized by present some criteria: i) the AE index must 54 reach an intensity peak greater than or equal to 1000 nT; ii) the The AE index needs 55 to be almost continuous and never drops below 200 nT for more than two hours at a 56 time; iii) the The event must have a duration of at least two days, and iv) the The 57 event occurred after the main phase of magnetic storms. However, the same physical 58 process may occur whether one of the four criteria are not strictly followed (Tsurutani 59 and Gonzalez, 1987; Tsurutani et al., 2004; Sobral et al., 2006, Tsurutani et al., 2006; 60 Hajra et al., 2013, Silva et al., 2017). As the main feature is the high AE index levels, 61 in this study we have considered drops below 200 nT for more than two hours as long 62 as the AE index value returns in high activity for prolonged hours.

The electron density perturbation in the ionosphere during HILDCAA events is different from that one occurred during geomagnetic storms in the equatorial and low latitudes stations. Since the HILDCAA presents a weak/moderate geoeffectiveness when it compares to the other forms of space disturbances, it is expected that the ionosphere response presents a different differential behavior.

The Total Electron Content (TEC) is an important ionospheric parameter to several studies and technologic applications. As HILDCAAs can cause F2-region peak alterations, it can be observed the enhancements/depletions in TEC profile. In fact, the TEC response to the geomagnetic storms is a well-known issue in the space physics field (Lu et al., 2001; Kutiev et al., 2005; Mendillo, 2006; Maruyama and
Nakamura, 2007; Biqiang et al., 2007; de Siqueira et al., 2011). However, only few
studies about TEC pattern during HILDCAAs intervals have been found in the
literature (de Siqueira et al., 2011).

76 Ionospheric storms are manifestations of space weather events, which are caused by 77 energy inputs in the upper atmosphere in the form of enhanced electric fields, 78 currents, and energetic particle precipitation (Buonsanto, 1999; Mendillo, 2006). 79 Usually, ionospheric storms are associated with ionosphere responses to geomagnetic 80 storm events. However, in a broader way, these responses happen due to 81 magnetospheric energy inputs to the Earth's upper atmosphere, and this can occur to all kind of geomagnetic activity form. Park (1974) pointed that ionospheric storms 82 83 can be understood in terms of the superposed effects of many substorm. In view of 84 the foregoing and considering that the development of ionospheric storms during 85 HILDCAAs intervals has not been dealt with in depth, in the current study we have 86 focused the TEC pattern during this kind of event.

87 Recently, Verkhoglyadova et al. (2013) suggested that HILDCAAs associated with 88 High Speed Streams (HSS) can be one of the external driving TEC variabilities. 89 Indeed, the continuous energy injection and energetic particles precipitation into the 90 polar upper atmosphere during HILDCAA intervals could modify the dynamic and 91 chemical coupling process of the thermosphere-ionosphere system resulting in 92 changes in the electron density. These modifications, beyond to change the auroral 93 electron density, can be mapped to low latitudes involving electric fields 94 disturbances, as prompt penetration electric fields (PPEF) and disturbance dynamo 95 (DD) (Koga et al., 2011; Silva et al., 2017; Yeeram and Paratrasri, 2019).

96 Therefore, in the current study we have focused the TEC pattern during HILDCAAs 97 intervals, taking into account taking account local time dependence, seasonal 98 dependence and high/slow speed streams influences in the equatorial and low latitude 99 ionosphere. This paper is structured as followed: in the next section we present the 100 HILDCAA intervals chosen to support this study as well as the GNSS receivers 101 locations over the Brazilian region. In section 3 we show the results and discussion of 102 the analysis and the conclusions are presented in the last section.

103

104 **2. Data and Methodology**

In this study was possible to construct an overall perception of the ionospheric storms occurred during HILDCAA disturbance time intervals that affect the TEC values with respect to the expected behavior for quiet days. The features studied are local time and seasonal dependences, and solar wind velocity influences.

109 We have selected ten HILDCAA intervals occurred during the 2015 – 2016 period.

110 These intervals are listed in Table 1, where the two columns present the identification

111 and the data range of each interval. The geomagnetic indices and interplanetary data

112 used to classify the HILDCAA events were obtained from OMNIWeb Plus data and

113 service (https://omniweb.gsfc.nasa.gov/ow.html). The Kp index data were obtained

114 from the World Data Center for Geomagnetism, Kyoto, Japan (<u>http://wde.kugi.kyoto-</u>

115 <u>u.ac.jp/kp/index.html</u>). In this work it was used the daily Kp sum value.

116 The TEC mean was initially processed by a program developed at the Institute for

117 Space Research, Boston College, USA (Krishna, 2017). The mean values of vertical

118 TEC (VTEC) were obtained from two Brazilian GNSS stations, São Luís (SL) (2.,59

119 S; 44., 21 W) and Cachoeira Paulista (CP) (22., 68 S; 44., 98 W), representing the

120 station closest to the equator and the low latitude station, respectively. The Rinex files

121 used in this study were obtained from Brazilian Network for Continuous Monitoring

122 of the GNSS-RBMC Systems (RBMC)

123 (https://www.ibge.gov.br/en/geosciences/geodetic-positioning/geodetic-

124 <u>networks/20079-brazilian-network-for-continuous-monitoring-of-the-gnss-systems-</u>

125 $\frac{2?=\&t=o-que-e}{}$. Besides that, the TEC data during HILDCAA events were analyzed 126 and then compared with a set of three days average belonging to a quiet period, in 127 which it refers to the three days less disturbed ($\Sigma Kp < 24$) of the month of the 128 occurrence of each HILDCAA interval.

Figure 1 shows a map with the location of each GNSS station, which is represented by a red triangle. The dashed line represents the magnetic equator. The TEC data obtained during the HILDCAA intervals were analyzed and then compared to the TEC data during the selected quiet days, resulting in dTEC (dTEC = TEC mean – TEC quiet days). All the analyses done in this work took into account the dTEC values.

135

136 **3. Results and Discussions**

In this section, we will present the ionospheric TEC responses observed during ten
HILDCAA intervals focusing on local time dependence and seasonal features and the
solar wind velocity influences.

140

141 3.1 Local time dependence

A common feature of ionospheric storms is to be associated with dependence on local
time, mainly when they are caused by geomagnetic storms (Titheridge and
Buonsanto, 1988; Pedatella et al., 2010). However, to the best of the authors'

145 knowledge, no study has been found analyzing this aspect when regarding HILDCAA146 intervals.

Figures 2 and 3 show the mean dTEC hourly values related to all HILDCAA intervals for São Luís and Cachoeira Paulista, respectively. Each panel represents a single interval from the bottom (H01) to the top (H10). The x axis is given in the Universal Time (LT = UT - 3) and the color scale represents the dTEC values in TEC units (TECu).

152 Notice that the dTEC values have a greater magnitude for the low latitude GNSS 153 station to the detriment of the closer equatorial GNSS station. The minimum and 154 maximum values are, respectively, -16.00 TECu and 27.40 TECu for to São Luís, and 155 -37.60 TECu and 48.80 TECu for to Cachoeira Paulista. These values were 156 considered to perform the TEC hourly distribution, i. e., for each specific GNSS 157 station, the maximum and minimum TEC values were used to analyze all HILDCAAs 158 in the same range. It was considered the same minimum and maximum values 159 occurred to all intervals, for each station. This fact explains why some intervals 160 appear too close to the quiet time pattern. We believed that since the HILDCAA 161 events has low/moderate geoeffectiveness it was not expected high values of the dTEC. 162

The distribution of the dTEC effects hour-to-hour during HILDCAA intervals shows a substantial variability from one event to another. Habarulema et al. (2013) found that the negative storms effects are observed during geomagnetic storms recovery phases that over equatorial latitudes. However, since HILDCAAs intervals are characterized by a long continuous phase of Dst index recovery, this does not apply. The HILDCAA intervals present the positive dTEC predominance. 60% (70%) of all intervals present a positive dTEC response during the whole event for São Luís

170 (Cachoeira Paulista). In a more simplified definition, HILDCAA means an interval 171 where there is always energy injection (Søraas et al., 2004; Sandanger et al., 2005). 172 Silva et al. (2017) observed that during HILDCAA intervals it was seen the uplift of 173 the equatorial F2 region peak height, probably due to prompt penetration electric 174 fields. One of the main mechanisms of TEC enhancements is the rise of the 175 ionosphere to higher altitudes where the recombination rates are small. Besides that, 176 our results are in agreement with the results found by de Siqueira et al. (2017). They 177 did a study comparing the TEC responses between two magnetic storms and two 178 HILDCAAs intervals following by them, and found a great TEC variability pattern 179 from one to another event. Hereupon, it was not possible to find a response pattern to 180 the HILDCAA effects in the equatorial and low latitude TEC considering only the 181 local time. There is great variability, and it is important to consider the day-to-day 182 ionospheric variabilities as well as the separate effect of each electric fields 183 disturbance (PPEF/DD).

Comparing both stations, Cachoeira Paulista GNSS station presented higher values both to positive as negative ionospheric storms. During the daytime hours, the latitude is responsible for the different ionospheric responses due to the presence of photoionization. This probably explains the dTEC higher sensibility to low latitude station in detriment of the closer equatorial latitude station.

Analyzing the hourly behavior of each interval from Figures 2 and 3, we observed more intensity in TEC disturbances, both for positive and negative storms, during some specific intervals. This aspect led us to make a seasonal analysis, which will be presented in the next section.

193

194 3.2 Seasonal Dependence

195 It is well known for geomagnetic storms that the influence of the season entails on 196 positive/negative ionospheric storms is more pronounced in winter/summer than in 197 equinox months (Matsushita, 1959; Prölss and Najita, 1975; Mendillo, 2006, among 198 others). However, has not yet been established whether the occurrence of HILDCAA 199 interval in different seasons can do different TEC disturbances.

200 In a recent study involving more than one hundred HILDCAA events, Hajra et al. 201 (2013) reported no seasonal dependence, in what regards to predominant occurrence 202 rate in any specific epoch of the year due to the solar cycle influences. They 203 announced the HILDCAAs may occur during any month and any year, with increases 204 in the numbers of events occurring during the solar cycle descending phase. In the 205 current study, it was considered as seasonal dependence feature the TEC disturbances 206 responses at HILDCAA intervals already classified in a seasonal way. The years 2015 and 2016 years comprise the descending phase of the 24th solar cycle, which 207 208 made it possible to catalog an expressive number of HILDCAAs events in a short 209 time. Among the ten intervals chosen for this study, we have separated eight ones to 210 represent the seasonal variability, being two events for each season station, taking 211 into account the month of occurrence of each interval, and considering the seasons as 212 they occur in South Hemisphere. The intervals are distributed according to the Table 213 2.

Figure 4 shows the disturbed TEC according to the seasonal classification which the blue and coral colors refer to São Luís and Cachoeira Paulista, respectively. The solid lines show an estimate of the central tendency for all values, minute-to-minute, for all days of the events belongs to the season, while the shaded area represents the confidence interval for that estimate. While the positive storms are more pronounced in the winter for geomagnetic storms, to HILDCAA intervals this season presents less geoeffectiveness, or almost none. Our results show that the equinoxes represent more ionospheric TEC responses during HILDCAA intervals than the solstices. Both equatorial and low latitude stations present positive storms during the autumn, while the spring presents a negative behavior, mainly. This equinoctial anomaly may be originated from the equinoctial differences in neutral winds, thermospheric composition, and electric fields. Additional studies are necessary to quantify how each factor can play an important role in HILDCAA seasonal TEC disturbances.

227

228 3.3 Solar wind velocities analysis

During the solar cycle descending phase, polar coronal holes migrate to lower latitudes emanating intense magnetic fields. When HSS from these low latitudinal coronal holes interact with slow speed streams (SSS) a region called Corotating Interaction Regions (CIR) is formed and it is well characterized by compressions of the magnetic field and plasma.

There are considerable works that whose show how HILDCAA is well associate with HSS and CIRs (Tsurutani et al., 2006; Verkhoglyadova et al., 2013). However, to be associated not necessarily means that the degree of geoeffectiveness is directly related to high speeds. Including, Yeeram (2019) suggest that Alfvén waves present during HILDCAA interval are more dominant than CIR-storms, revealing that both are controlled by different interplanetary drivers.

Figure 5 shows the solar wind velocities (V_{SW}) during each HILDCAA interval. As the Figure 4, the blue and coral colors refer to São Luís and Cachoeira Paulista, respectively. The diameter of the bubble is related to the velocity. The results showed great variability from one interval to another, even considering the intervals that occurred in the same year. In our first analysis (not shown here) we did not find a direct association or cross-correlation between the $V_{SW}SW$ magnitude and the dTEC in the equatorial and low latitude GNSS stations. Kim (2007) indicated that HILDCAA intervals can be accompanied by HSS as well as SSS. It is possible to see in our results that the dTEC responses to some intervals present similar behavior to both HSS and SSS (e.g. H03, H07 and H08). This means that HILDCAA intervals can affect the ionospheric TEC, but not in a direct correlation.

251

4. Conclusions

For this work, the ionospheric TEC response to a sample of ten HILDCAA intervals has been studied. We have used two GNSS stations from RBMC network representing equatorial and low latitude locations. As HILDCAA can affect the equatorial ionospheric F2 region, some disturbed TEC from its quiet time pattern is found. Addressing how the ionospheric storms behave during the HILDCAA intervals is our main goal.

259 In summary Summarizing, HILDCAAs geoeffectiveness in Earth is mainly associated 260 with CIRs, for this reason, the HILDCAA occurrence is more recurrent in the solar 261 cycle descending phase since CIRs play a major role during this phase. Their effects 262 occur during magnetic reconnection due to association with southward z component 263 of the interplanetary magnetic field and Alfvén waves present in it (Tsurutani et al., 264 2004). These long-lasting intervals are due to continuous injection of energy and 265 precipitation of particles, which disturb the high latitude ionosphere. The mainly 266 disturbs are changes in thermospheric neutral composition, temperature, winds and 267 electric fields. Similar to geomagnetic storms, theses disturbs can be mapped to low 268 and equatorial latitude and alter the quiet time ionosphere. However, generally, they 269 are less intense because in one astronomical unit the CIRs are not fully developed. In this study we seek to understand the behavior of the ionospheric storm duringHILDCAA intervals. The main results are highlighted below:

The hourly distribution of the dTEC during HILDCAAs intervals may vary
substantially between low and equatorial latitude. Probably, the photoionization
associated with latitude is responsible for these variations;

• Despite the geomagnetic storms recovery phase presents negative ionospheric storms, this pattern do not occur during HILDCAA intervals. There is great variability from one interval to another, but, predominantly, occurs positive phase;

Regarding seasonal features, while the positive storms are more pronounced in the
 winter for geomagnetic storms, this season present less geoeffectiveness, or almost
 none to HILDCAA intervals. The equinoxes represent more ionospheric responses
 to HILDCAA intervals presenting positive/negative phase predominance during
 the autumn/spring;

A well-known HILDCAA feature is its association with HSS present in the solar
 wind. However, this association does not present a direct relation regards to TEC
 disturbances in low and equatorial latitudes.

To conclude, the upshot of this study is the possibility to understand how ionospheric storms behave during some HILDCAA intervals and to contribute to improving the discussions about this issue.

289

290 Data availability

- 291 The data used in this work are made publicly available on the following sites:
- 292 <u>https://omniweb.gsfc.nasa.gov/ow.html</u>, <u>http://wdc.kugi.kyoto-u.ac.jp/kp/index.html</u>,
- and https://www.ibge.gov.br/en/geosciences/geodetic-positioning/geodetic-
- 294 <u>networks/20079-brazilian-network-for-continuous-monitoring-of-the-gnss-systems-</u>
- 295 2?=&t=o-que-e. The GPS-TEC program used in this work is available in
- 296 <u>http://seemala.blogspot.com/</u>
- 297

298 Author contributions

- 299 R. P. Silva conceived the study, designed the data analysis, discussed the results and
- 300 leaded writing this manuscript.
- 301 C. M. Denardini assisted to conceive the study, to design the GNSS data analysis and302 discuss the final results.
- 303 M. S. Marques assisted with the GNSS data analysis and with designing the figures.
- 304 L. C. A. Resende assisted to design the study and discuss the results of the study.
- 305 J. Moro assisted to design the study and discuss the results of the study.
- 306 G. A. S. Picanço assisted to discuss the results of the study and review the 307 manuscript.
- 308 G. L. Borba assisted to discuss the results of the study and review the manuscript.
- 309 M. A. F. Santos assisted to discuss the results of the study and review the manuscript.
- 310 All the authors helped to write and to revise the manuscript.
- 311

312 **Competing interests**

313 The authors declare that they have no conflict of interest.

315 Special issue statement

This article is part of the special issue "7th Brazilian meeting on space geophysics and aeronomy". It is a result of the Brazilian meeting on Space Geophysics and Aeronomy, Santa Maria/RS, Brazil, 5–9 November 2018.

319

320 Acknowledgements

321 R. P. Silva acknowledges the supports from Conselho Nacional de Desenvolvimento 322 Científico e Tecnológico (CNPq) through the grant 300329/2019-9. C. M. Denardini 323 thanks to CNPq/MCTIC (Grant 303643/2017-0). L. C. A. Resende would like to thank the National Space Science Center (NSSC), Chinese Academy of Sciences 324 (CAS) for supporting her postdoctoral. J. Moro would like to acknowledge the China-325 326 Brazil Joint Laboratory for Space Weather (CBJLSW), National Space Science 327 Center (NSSC), Chinese Academy of Sciences (CAS) for supporting his Postdoctoral 328 fellowship, and the CNPq for the grant 429517/2018-01. G. A. S. Picanço thanks 329 CAPES for supporting his Ph.D. (Grant 88887.351778/2019-00). We also would like 330 **OMNIWeb** thank the Plus data and service **OMNI** -data to 331 (https://omniweb.gsfc.nasa.gov/ow.html), and the World Data Center for Geomagnetism, Kyoto (http://wdc.kugi.kyoto-u.ac.jp/kp/index.html). The Rinex files 332 333 were obtained from Brazilian Network for Continuous Monitoring of the GNSS-334 RBMC Systems (RBMC) interface at 335 https://www.ibge.gov.br/en/geosciences/geodetic-positioning/geodeticnetworks/20079-brazilian-network-for-continuous-monitoring-of-the-gnss-systems-336 337 2?=&t=o-que-e. The authors acknowledge Gopi Seemala for making available the

- 338 GPS-TEC program (<u>http://seemala.blogspot.com/</u>).
- 339
- 340

341 **References**

- 342 Biqiang, Z., Weixing, W., Libo, L., and Tian, M.: Morphology in the total electron
- 343 content under geomagnetic disturbed conditions: results from global ionosphere
- 344 maps, Ann. Geophys., 25, 1555-1568, <u>https://doi.org/10.5194/angeo-25-1555-2007</u>,
- 345 2007.
- Buonsanto, M. J.: Ionospheric storms A review. Space Science Reviews, 88(3-4),
- 347 563-601, <u>https://doi.org/10.1023/A:100510753</u>, 1999.
- de Siqueira, P. M., de Paula, E. R., Muella, M. T. A. H., Rezende, L. F. C., Abdu, M.
- 349 A., and Gonzalez, W. D.: Storm-time total electron content and its response to
- 350 penetration electric fields over South America, Ann. Geophys., 29, 1765-1778,
- 351 <u>https://doi.org/10.5194/angeo-29-1765-2011</u>, 2011.
- de Siqueira Negreti, P. M., de Paula, E. R., and Candido, C. M. N.: Total electron
- 353 content responses to HILDCAAs and geomagnetic storms over South America, Ann.
- 354 Geophys., 35, 1309-1326, <u>https://doi.org/10.5194/angeo-35-1309-2017</u>, 2017.
- 355 Habarulema, J. B., McKinnell, L. A., Burešová, D., Zhang, Y., Seemala, G., Ngwira,
- 356 C., Chum, J., Opperman, B.: A comparative study of TEC response for the African
- ad mid-latitudes during storm conditions J. Atmos. Sol. Terr. Phys. 102,
- 358 105-114, <u>https://doi.org/10.1016/j.jastp.2013.05.008</u>, 2013.
- 359 Hajra, R., Echer, E., Tsurutani, B. T., Gonzalez, W. D.: Solar cycle dependence of
- 360 High-Intensity Long-Duration Continuous AE Activity (HILDCAA) events,
- 361 relativistic electron predictors?: J. Geophys. Res. 118, 5626-5638,
- 362 <u>https://doi.org/10.1002/jgra.50530</u>, 2013.
- 363 Kim, H.: Study on the particle injections during HILDCAA intervals. Journal of
- Astronomy and Space Sciences, v. 24, <u>https://doi.org/10.5140/JASS.2007.24.2.119</u>,
- 365 2007.
- 366 Koga, D., Sobral, J. H. A., Gonzalez, W. D., Arruda, D. C. S., Abdu, M. A., Castilho,
- 367 V. M., Mascarenhas, M., Gonzalez, A. C., Tsurutani, B. T., Denardini, C. M.,
- 368 Zamlutti, C. J.: Electrodynamic coupling process between the magnetosphere and the

- 369 equatorial ionosphere during a 5-day HILDCAA event. J. Atmos. Sol. Terr. Phys. 73,
- 370 148-155, <u>https://doi.org/10.1016/j.jastp.2010.09.002</u>, 2011.
- 371 Krishna, S. G., GPS-TEC Analysis Software Version 2.9.5, Sep. 2017, [online]
- 372 Available: <u>http://seemala.blogspot.in/</u>.
- 373 Kutiev, I., Watanabe, S., Otsuka, Y., Saito, A.: Total electron content behavior over
- Japan during geomagnetic storms. J. Geophys. Res, 110(A1),
- 375 <u>https://doi.org/10.1029/2004JA010586</u>, 2005.
- Lu, G., Richmond, A. D., Roble, R. G., Emery, B. A.: Coexistence of ionospheric
- 377 positive and negative storm phases under northern winter conditions: A case study. J.
- 378 Geophys. Res, 106(A11), 24493-24504, <u>https://doi.org/10.1029/2001JA000003</u>, 2001
- 379 Maruyama, T., & Nakamura, M.: Conditions for intense ionospheric storms
- 380 expanding to lower midlatitudes. J. Geophys. Res, 112(A5),
- 381 <u>https://doi.org/10.1029/2006JA012226</u>, 2007.
- 382 Matsushita, S.: A study of the morphology of ionospheric storms. J. Geophys. Res. v.
- 383 64, n. 3, p. 305-321, <u>https://doi.org/10.1029/JZ064i003p00305</u>, 1959.
- 384 Mendillo, M.: Storms in the ionosphere: Patterns and processes for total electron
- content. Reviews of Geophysics, v. 44, n. 4, <u>https://doi.org/10.1029/2005RG000193</u>,
 2006.
- 387 Park, C. G.: A morphological study of substorm-associated disturbances in the
- ionosphere. Journal of Geophysical Research, 79(19), 2821-2827,
- 389 <u>https://doi.org/10.1029/JA079i019p02821</u>, 1974.
- 390 Pedatella, N. M., Lei, J., Thayer, J. P., & Forbes, J. M.: Ionosphere response to
- 391 recurrent geomagnetic activity: Local time dependency. Journal of Geophysical
- 392 Research: Space Physics, 115(A2), <u>https://doi.org/10.1029/2009JA014712</u>, 2010.
- 393 Prölss, G. W., Najita, K.: Magnetic storm associated changes in the electron content
- at low latitudes. Journal of Atmospheric and Terrestrial Physics, v. 37, n. 4, p. 635-
- 395 643, <u>https://doi.org/10.1016/0021-9169(75)90058-6</u>, 1975.

- 396 Silva, R. P., Sobral, J. H. A., Koga, D., and Souza, J. R.: Evidence of prompt
- 397 penetration electric fields during HILDCAA events, Ann. Geophys., 35, 1165-1176,
- 398 <u>https://doi.org/10.5194/angeo-35-1165-2017</u>, 2017.
- 399 Sobral, J. H. A., Abdu, M. A., Gonzalez, W. D., Clua De Gonzalez, A. L., Tsurutani,
- 400 B. T., Da Silva, R. R. L, Barbosa, I. G., Arruda, D. C. S., Denardini, C. M., Zamlutti,
- 401 C. J., Guarnieri, F.: Equatorial ionospheric responses to high-intensity long-duration
- 402 auroral electrojet activity (HILDCAA), J. Geophys. Res. 111, A07S02,
- 403 <u>https://doi.org/10.1029/2005JA011393</u>, 2006.
- 404 Sandanger, M. I., Søraas, F., Aarsnes, K., Oksavik, K., Evans, D. S., & Greer, M. S.:
- 405 Proton Injections Into the Ring Current Associated With B~ z Variations During
- 406 HILDCAA Events. GEOPHYSICAL MONOGRAPH-AMERICAN GEOPHYSICAL
- 407 UNION, 155, 249, <u>https://doi.org/10.1029/155GM26</u>, 2005.
- 408 Søraas, F., Aarsnes, K., Oksavik, K., Sandanger, M. I., Evans, D. S., Greer, M. S.
- 409 (2004). Evidence for particle injection as the cause of Dst reduction during
- 410 HILDCAA events. J. Atmos. Sol. Terr. Phys. 66(2), 177-186,
- 411 <u>https://doi.org/10.1016/j.jastp.2003.05.001</u>, 2004.
- 412 Titheridge, J. E., Buonsanto, M. J.: A comparison of northern and southern
- 413 hemisphere TEC storm behaviour. J. Atmos. Sol. Terr. Phys. 50(9), 763-780,
- 414 <u>https://doi.org/10.1016/0021-9169(88)90100-6</u>, 1988.
- 415 Tsurutani, B. T., Gonzalez, W.D.: The cause of high intensity long-duration
- 416 continuous AE activity (HILDCAA): interplanetary Alfvén wave trains. Planet Space
- 417 Sci. 35, 405-412, <u>https://doi.org/10.1016/0032-0633(87)90097-3</u>, 1987.
- 418 Tsurutani, B. T., Gonzalez, W.D., Guarnieri, F., Kamide, Y., Zhou, X., Arballo, J. K.:
- 419 Are high-intensity long-duration continuous AE activity (HILDCAA) events
- 420 substorm expansion events?, J. Atmos. Sol. Terr. Phys. 66,167-176,
- 421 <u>https://doi.org/10.1016/j.jastp.2003.08.015</u>, 2004.
- 422 Tsurutani, B. T., Gonzalez, W.D., Gonzalez, A. L. C., Guarnieri, F. L., Gopalswamy,
- 423 N., Grande, M., Kamide, Y., Kasahara, Y., Lu, G., Mann, I., McPherron, R., Soraas,
- 424 F., Vasyliunas, V.: Corotating solar wind streams and recurrent geomagnetic activity:

- 425 A review. J. Geophys. Res. 111, A07S01, <u>https://doi.org/10.1029/2005JA011273</u>,
 426 2006.
- 427 Verkhoglyadova, O. P., Tsurutani, B. T., Mannucci, A. J., Mlynczak, M. G., Hunt, L.
- 428 A., and Runge, T.: Variability of ionospheric TEC during solar and geomagnetic
- 429 minima (2008 and 2009): external high speed stream drivers, Ann. Geophys., 31,
- 430 263-276, <u>https://doi.org/10.5194/angeo-31-263-2013</u>, 2013.
- 431 Yeeram, T.; Paratrasri, A.: Recurrent geomagnetic storms and equinoctial ionospheric
- 432 F-region in the low magnetic latitude: a case study. In: Journal of Physics:
- 433 Conference Series. IOP Publishing, 012024, https://doi.org/10.1088/1742-
- 434 <u>6596/1144/1/012024</u>, 2018.
- 435 Yeeram, T.: The solar wind-magnetosphere coupling and daytime disturbance electric
- 436 fields in equatorial ionosphere during consecutive recurrent geomagnetic storms. J.
- 437 Atmos. Sol. Terr. Phys, 187, 40-52, <u>https://doi.org/10.1016/j.jastp.2019.03.004</u>, 2019.

438

440 **Figure captions**

- 441 **FIGURE 1** Map showing the locations of the GNSS stations used in the present
- study. Both stations are localized in the Brazilian region and are marked by a red triangle,
- 443 where SL and CP are, respectively, São Luís and Cachoeira Paulista.
- 444 **FIGURE 2** dTEC hourly values to all HILDCAA intervals to São Luís (equatorial
- 445 station).
- 446 **FIGURE 3** dTEC hourly values to all HILDCAA intervals to Cachoeira Paulista
- 447 (low latitude station).
- 448 **FIGURE 4** Seasonal dTEC response to HILDCAA intervals. The blue and coral
- 449 lines refer to São Luís and Cachoeira Paulista, respectively.
- 450 **FIGURE 5** Solar wind velocities analysis during HILDCAA intervals. The blue
- 451 and coral colors refer to São Luís and Cachoeira Paulista stations, respectively, while
- 452 the bubble diameter is related to velocity (km/s).
- 453

Table captions

- **TABLE 1** The date range for HILDCAA intervals identified during 2015 2016
- 457 years
- **TABLE 2** Seasonal classification of HILDCAA intervals (according to the seasons
- 459 in the Southern hemisphere).

FIGURE 1 –





FIGURE 2 –



FIGURE 2 –





FIGURE 3 –



FIGURE 3 –



FIGURE 4 –



FIGURE 5 –



TABLE 1 –

ID	Date range
H01	2015/03/01 - 03
H02	2015/03/17 - 21
H03	2015/04/16 - 20
H04	2015/06/08 - 11
H05	2015/07/11 - 14
H06	2015/08/15 - 18
H07	2015/10/07 - 14
H08	2016/07/09 - 12
H09	2016/08/03 - 07
H10	2016/12/08 - 11

TABLE 2 –

Season	HILDCAA Intervals
Autumn	H03 and H04
Winter	H05 and H06
Spring	H07 and H10
Summer	H01 and H02