

---

# An inter-hemispheric seasonal comparison of Polar Amplification using radiative forcing of quadrupling CO<sub>2</sub> experiment

Fernanda Casagrande<sup>1</sup>, Ronald Buss de Souza<sup>1</sup>, Paulo Nobre<sup>1</sup>, Andre Lanfer Marquez<sup>1</sup>

5 <sup>1</sup>Center for Weather Forecasting and Climate Studies, National Institute for Space Research, Cachoeira Paulista, 13620-000 Brazil.

*Correspondence to:* Fernanda Casagrande (Fernanda.casagrande@inpe.br)

**Abstract.** The numerical climate simulations from the Brazilian Earth System Model (BESM) are used here to investigate the response of Polar Regions to a forced increase of CO<sub>2</sub> (Abrupt-4xCO<sub>2</sub>) and compared with Coupled Model Intercomparison Project phases 5 (CMIP5) and 6 (CMIP6) simulations. The main objective here is to investigate the seasonality of the surface and vertical warming as well as the coupled processes underlying the polar amplification, as changes in sea ice cover. Polar Regions are described as the most climatically sensitive areas of the globe, with an enhanced warming occurring during the cold seasons. The asymmetry between the two poles is related to the thermal inertia and the coupled ocean atmosphere processes involved. While in the northern high latitudes the amplified warming signal is associated to a positive snow and sea ice albedo feedback, for southern high latitudes the warming is related to a combination of ozone depletion and changes in the winds pattern. The numerical experiments conducted here demonstrated a very clear evidence of seasonality in the polar amplification response, as well as, linkage with sea ice changes. In winter, for the northern high latitudes (southern high latitudes) the range of simulated polar warming varied from 10 K to 39 K (-0.5 K to 13 K). In summer, for northern high latitudes (southern high latitudes) the simulated warming varies from 0 K to 23 K (0.5 K to 14 K). The vertical profiles of air temperature indicated stronger warming at the surface, particularly for the Arctic region, suggesting that the albedo-sea ice feedback overlaps with the warming caused by meridional transport of heat in the atmosphere. The latitude of the maximum warming was inversely correlated with changes in the sea ice within the model's control run. Three climate models were identified as having high polar amplification for the Arctic cold season (DJF): IPSL-CM6A-LR (CMIP6), HadGEM2-ES (CMIP5) and CanESM5 (CMIP6). For the Antarctic, in the cold season (JJA), the climate model identified as having high polar amplification were: IPSL-CM6A-LR (CMIP6), CanESM5 (CMIP6) and FGOALS-s2 (CMIP5). The large decrease in sea ice

---

30 concentration is more evident in models with great Polar Amplification, and for the same range of  
latitude (75°N – 90°N). Also, we found, for models with enhanced warming, expressive changes in the  
sea ice annual amplitude with outstanding ice-free conditions from May to December (EC-Earth3-Veg)  
and June to December (HadGEM2-ES). We suggest that the large bias found among models can be  
35 related to the differences in each model to represent the feedback process and also as a consequence of  
each distinct sea ice initial conditions. The polar amplification phenomenon has been observed  
previously and is expected to become stronger in the coming decades. The consequences for the  
atmospheric and ocean circulation are still subject to intense debate in the scientific community.

## 1 Introduction

Polar regions have been shown to be more sensitive to climate change than the rest of the world  
40 (Smith et al., 2019; Serreze and Barry, 2011). The Arctic is warming at least twice as fast as the  
northern hemisphere and as the globe as a whole. This phenomenon is known as the Arctic  
Amplification (AA) and is combined with a fast shrinking of the sea ice cover (Serreze and Barry, 2011;  
Kumar et al., 2010; Screen and Simmonds, 2010). Previous research has indicated that the enhanced  
Arctic warming is a response to anthropogenic Greenhouse Gas (GHG) forcing, which, in turn,  
45 intensifies many complex non-linear coupled ocean-atmosphere feedbacks (e.g. the sea ice albedo  
feedback) (Stuecker et al., 2018; Pithan and Mauritsen, 2014; Alexeev et al., 2005). The sea ice-albedo  
feedback is one of the key mechanisms to amplify the Arctic warming, playing an important role in  
global climate change (Stuecker et al., 2018; Pithan and Mauritsen, 2014). In contrast to the Arctic sea  
ice, the total sea ice cover surrounding the Antarctic continent has increased in association with cooling  
50 over eastern Antarctica and warming over the Antarctic Peninsula. The physical ocean atmosphere  
coupled processes responsible for Antarctic sea ice rising are still unclear. Turner et al., (2017) show the  
unprecedented springtime retreat of Antarctic sea ice in 2016. However, results derived from numerical  
simulations and observations point to a combination of changes in the wind pattern, the ocean  
circulation, accelerated basal melting Antarctica's ice shelf and the ozone depletion (Marshall et al.,  
55 2014; Bintanja et al., 2013; Thompson et al., 2011; Thompson and Solomon, 2002). According to  
Marshall et al., (2014), these two-poles inter-hemispheric asymmetries strongly influence the Sea

---

Surface Temperature (SST) response to an increase in the global CO<sub>2</sub> forcing, accelerating the warming in the Arctic while delaying it in Antarctica.

Numerous scientific publications based on both, observations and state-of-the-art Global Climate Model simulations for the high latitudes of the northern hemisphere have shown that AA is an intrinsic feature of the Earth's climate system (Smith et al., 2019; Vaughan et al., 2013; Serreze and Barry, 2011; Screen and Simmonds, 2010). These works suggested that the Surface Air Temperature (SAT) will continue to increase with effects extending beyond the Arctic region (Dethloff et al., 2019; Smith et al., 2019; Bintanja et al., 2013; Serreze and Barry, 2011; Winton 2006; Holland and Bitz, 2003). Although the annual average SAT at northern mid- and high latitudes is increasing, the wintertime SAT has decreased since the 1990 (Zhang et al., 2016; Mori et al., 2014; Cohen et al., 2012; Honda et al., 2009).

Bekryaev et al., (2010), for instance, found a warming rate of 1.36° C century<sup>-1</sup> for the period from 1875 to 2008 using an extensive set of observational data from meteorological stations located at high latitudes of the northern hemisphere (> 60° N). That trend is almost double that of the northern hemisphere trend as a whole (0.79° C century<sup>-1</sup>), with an accelerated warming rate in the most recent decade. Rigor et al., (2002) also using an observational dataset showed that the Arctic warming varies largely among regions and that changes in SAT are also related to the Arctic Oscillation (Ambaum et al., 2001).

The Arctic Ocean temperature and ocean heat fluxes also have increased over the past several decades (Walsh, 2014; Polyakov et al., 2010; Polyakov et al., 2008). According to Polyakov et al., (2017), the recent sea ice shrinking, weakening of the halocline and shoaling of the intermediate-deep Atlantic water masses layer in eastern Eurasia Basin have increased the winter ventilation in the ocean interior, making the region structurally similar to the western Eurasian Basin. The authors described these processes as an “Atlantification” phenomenon and represent an essential step toward a new Arctic climate state.

Holland and Bitz, (2003) using a set of 15 state-of-the-art CMIP models found that the range of simulated Arctic warming as response to a doubling of CO<sub>2</sub> concentration varies largely between the models ranging from 1.5 to 4.5 times the global mean warming. The large differences among the

---

models are related to differences in simulating the ocean's meridional heat transport, the polar cloud cover and the sea ice (e.g. a simulation with thinner sea ice cover presents a higher polar amplification).

According to Shu et al., (2015), Global Climate Models in general offer much better simulations for the Arctic than for the Antarctica. Turner et al., (2015) suggested that the main problem of climate models in the high latitudes of the southern hemisphere is their inability to reproduce the observed (although slight) increase in Sea Ice Extent (SIE). Bintanja et al., (2015) and Swart and Fyfe, (2013) have demonstrated the importance to include the effect of the increasing freshwater input from Antarctic continental ice into the Southern Ocean. The authors described that the ice sheet dynamics, essential for having accurate sea ice simulations, is currently disregarded in all CMIP5 models. Swart and Fyfe (2013) also suggested that this deficiency may significantly influence the simulated sea ice trend because the subsurface ocean warming causes basal ice-shelf melt, freshening the surface waters, which eventually leads to an increase in sea ice formation. Moreover, the instrumental network for data collection in Antarctica and the Southern Ocean is considered scarce (even more than in the Arctic), inhomogeneous and insufficiently dense to validate climate models. Therefore, for the high latitudes regions of the southern hemisphere, the effects of the ongoing climate change and its associated processes are still considered hot topics that lack conclusive answers.

How the polar climate will change as response to an external forcing deeply depends on feedback processes, which operates to amplify or diminish the effects of climate change forcing. These feedbacks depend on the integrated coupled processes between ocean-atmosphere-cryosphere over a large spectrum of spatial and temporal scales, which makes the quantification of them even more complicated.

Here the seasonal sensitivity of high latitudes as a response to quadrupling atmospheric CO<sub>2</sub> is investigated using the recently developed Brazilian Earth System Model, coupled ocean-atmosphere version 2.5 (BESM-OA V2.5) and comparing its results with those from 32 other Coupled General Circulation Models participating in CMIP5 and CMIP6. Our goal is to investigate the coupled processes underlying the polar warming by seasons. The paper is organized as follows: Section 2 provides a description of the climate models and experimental design[s] used in this work, focusing on the BESM-OA V2.5 model (Veiga et al., 2019; Giarolla et al., 2015; Nobre et al., 2013). In Section 3, the

---

seasonality in the surface warming in high latitudes is examined of both northern and southern hemispheres and results from different models are compared. Section 4 provides an analysis of the vertical structure of air temperature warming, spatial pattern of sea ice changes and a discussion about the coupled ocean atmosphere processes and feedback mechanisms involved. A summary of results and conclusions are presented in Section 5.

## 2 Data Sources

### 2.1 Numerical Design

This study used two numerical experiments from CMIP5 and CMIP6: (i) piControl: it runs for 700 years, forced by invariant pre-industrial atmospheric CO<sub>2</sub> concentration level (280ppmv) and (ii) Abrupt 4xCO<sub>2</sub>: it runs for 460 years, comprising an abrupt instantaneous quadrupling of atmospheric CO<sub>2</sub> level concentration from the piControl simulation. The design of both experiments follows the CMIP5 protocol (Taylor et al., 2012) and Eyring et al. (2016) for CMIP6 numerical experiments.

Although an instantaneous quadrupling CO<sub>2</sub> scenario is not realistic for the 21st century compared with RCP scenarios and observations, this scenario can give us a measure of climate sensitivity and how large can be the response of the polar region in comparison to the globe as a whole. The results are compared for polar amplification (changes in air temperature) and sea ice cover, for the same numerical experiment.

For CMIP5 numerical experiment, the follow models are used: BESM-OA V2.5 (Nobre et al., 2013; Veiga et al., 2019), ACCESS-3 (Bi et al., 2013; Collier and Uhe, 2012), GFDL-ESM2M (Griffies, 2012), IPSL-CM5-LR (Dufresne et al., 2013), MIROC-ESM (Watanabe et al., 2011), MPI-ESM-LR (Stevens et al., 2013), NCAR-CCSM4 (Gent et al., 2011), CanESM2 (Chylek et al., 2011), FGOALS-s2 (Bao et al., 2013), GFDL-ESM2G (Delworth et al., 2006), GISS-E2\_H (Schmidt et al., 2006), HadGEM2-ES (Collins et al., 2008), MIROC5(Watanabe et al., 2010), MPI-ESM-P(Giorgetta et al., 2013), MRI-CGCM3(Yukimoto et al., 2012).

For CMIP6 numerical experiments, the follow models are used: ACCESS-CM2 (Bi et al., 2013), CAMS-CSM1-0 (Rong, 2019), CanESM5 (Swart et al., 2019), CMCC-CM2-SR5 (Fogli et al., 2019),

---

CNRM-ESM2-1 (Séferian et al., 2019), ACCESS-ESM1-5 (Ziehn et al., 2019), E3SM-1-0 (Bader et al., 2019), EC-Earth3-Veg, FGOALS-G3 (Li et al., 2020), GISS-E2-1-H (Schmidt et al., 2006), INM-CM4-  
140 8 (Volodin et al., 2019), MIROC6 (Tatebe et al., 2018), MIROC-ES2L (Hajima et al., 2019), MPI-  
ESM1-2-LR (Fiedler et al., 2019), MRI-ESM2-0 (Yukimoto et al., 2019).

## 1.2 Brazilian Earth System Model

The Brazilian Earth System Model, Version 2.5 (BESM-OA2.5) used here is a global climate coupled ocean-atmosphere-sea ice model, and is part of CMIP5 project. The atmospheric component of  
145 BESM-OA2.5 is BAM (Brazilian Atmospheric Model) and was described in detail by Figueroa et al.,  
(2016). The latest version of BAM, used here and described by Figueroa et al., (2016) and Veiga et al.,  
(2019), has spectral horizontal representation truncated at triangular wave number 62, grid resolution of  
approximately  $1.875^\circ \times 1.875^\circ$ , and 28 sigma levels in the vertical, with unequal increments between the  
vertical levels (i.e., a T62L28). Two important changes were implemented on the BESM last version: (i)  
150 a new microphysics scheme, described by Ferrier et al., (2002) and Capistrano et al., (2020) and (ii) a  
new surface layer scheme, described by Capistrano et al., (2020) and Jimenez and Dudhia, (2012).  
These key changes represent an improvement in surface layer, resulting in better representation of near-  
surface air temperature, wind and humidity at 10 m. The main improvements occur over the ocean,  
where temperature, wind and humidity are important to calculate the heat fluxes at ocean-atmosphere-  
155 sea ice interface.

The oceanic component of BESM-OA2.5 is the Modular Ocean Model, Version 4p1, from  
National Oceanic and Atmospheric Administration-Geophysical Fluid Dynamics Laboratory  
(MOM4p1/NOAA-GFDL), described in detail by Griffies, (2009). The MOM4p1 includes a Sea Ice  
 Simulator (SIS) built-in ice model (Winton 2000). The SIS has five ice thickness categories and three  
160 vertical layers (one snow and two ice). To calculate ice internal stresses are used the elastic-viscous-  
plastic technique described by Hunke and Dukowicz, (1997). The thermodynamics is given by a  
modified Semtner's three-layer scheme (Semtner, 1976). SIS is able to calculate sea ice concentrations,  
snow cover, thickness, brine content and temperature. The horizontal grid resolution of MOM4p1 in the  
longitudinal direction is a set to  $1^\circ$ . The latitudinal direction varies uniformly, in both hemispheres,


165 from  $1/4^\circ$  between  $10^\circ$  S and  $10^\circ$  N to  $1^\circ$  of resolution at  $45^\circ$  and to  $2^\circ$  of resolution at  $90^\circ$ . The vertical  
axis has 50 levels (upper 220m, has 10 m resolution, increasing to about 360 at deeper levels. The  
MOM4p1 and BAM models were coupled using FMS coupler. FMS coupled was developed by  
NOAA-GFDL. The BAM model receives SST and ocean albedo from MOM4p1 and SIS (hour by  
hour). The MOM4p1 receives momentum fluxes, specific humidity, pressure, heat fluxes, vertical  
170 diffusion of velocity components and freshwater. The Monin-Obukhov scheme is used to calculate the  
wind stress fields.

### 3 Results and Discussion

First we discuss the seasonality of polar warming near the surface in the Arctic, vertical profile,  
sea ice changes, differences among models and coupled process involved. Follow, we do the same  
175 analysis for the southern high latitudes and assess the reasons for asymmetries between poles.

#### 3.1 Polar Amplification

In order to evaluate the seasonality of near surface polar warming, the seasons are defined as  
follows: December to February (DJF) as boreal winter, March to May (MAM) as boreal spring, June to  
August (JJA) as boreal summer, and September to November (SON) as boreal fall.



180

**Figure 1. Polar Amplification using Long-term observations of Surface Air Temperatures ( $^\circ$ C) at 2008-2018 (seasonal average) relative to 1979-1989 (seasonal average) in (a) Winter (DJF) and (b) Summer (JJA). Source: Era Interim Reanalysis.**

---

Figure 1 shows the enhanced surface warming at high latitudes compared to the rest of globe, with a slightly greater rate of warming in the 20<sup>th</sup> century. This Polar Amplification is not symmetric, most evidence is from Arctic region (during the boreal winter). According to Stocker et al., (2013), the enhanced warming at northern high latitudes was linked with decrease in snow cover and sea ice concentration, sea level rise and increase in land precipitation. Furthermore, changes in atmospheric and ocean circulations (Pedersen et al., 2019; Pithan and Mauritsen, 2014; Stocker et al., 2013; Yang et al., 2010; Graversen et al., 2008). Polar Amplification is also reported by Climate Models, driven by solar or natural carbon cycle perturbations (Sundqvist et al., 2010; O'ishi and Abe-Ouchi, 2009; Mann et al., 2009; Masson-Delmotte et al., 2006).

Figure 2 and Figure 3 show the seasonality of the Polar Amplification (change in zonally SAT average) simulated by BESM-AO V2.5 and 32 state-of-art CMIP5 and CMIP6 models. To assess the climate sensitivity of polar amplification, seasonally and coupled processes involved, we used the difference between Abrupt 4xCO<sub>2</sub> and piControl numerical experiments, considering only the last 30 years of the 150 years of model integration after quadrupling CO<sub>2</sub> concentration (when the model reaches a new equilibrium state). This procedure has been largely used by researchers, since it allows us to evaluate and compare potential warming and sensitivities among low and high latitudes as well as to compare differences between models (Van der Linden et al., 2019; Cvijanovic et al., 2015; Manabe et al., 2004; Holand and Bitz, 2003).

Under the largest future GHG forcing (4xCO<sub>2</sub>), the polar regions are found to be the most sensitive areas of the globe, with a very pronounced seasonality (Figure 2 and Figure 3). The high southern latitudes warming predicted by the models analyzed is modest in relation to the Arctic's, but still not negligible. This asymmetry is partly due to the smaller area covered by ocean in Northern Hemisphere that induces a smaller thermal inertia. Contrasting to the high latitudes, the tropical warming is similar for both hemispheres, without the robust warming pattern as showed in high latitudes. Salzmann (2017) suggested that the overall weaker warming in Antarctica is due to a more efficient ocean heat uptake in the southern ocean, weaker surface albedo feedback in combination with ozone depletion. BESM-OA V2.5 model has no ozone chemistry as a climate component, so we suggest that even neglecting the ozone depletion, the weaker warming in Antarctica will be shown. Also is



---

215 expected a weak albedo sea ice feedback compared with Arctic region (because the fast retreat of sea ice on the Northern Hemisphere). The role of the Antarctica surface height for both feedbacks processes and meridional transports is similarly important to consider. According to Salzmann (2017), the polar amplification asymmetry is explained by the difference in surface height. If Antarctica is considered to be flat in a climate simulation with CO<sub>2</sub>-doubling experiment, the north-south asymmetry is reduced.

220 From September to February (boreal autumn and winter), the surface warming is maximum at northern high latitudes, decreasing with latitude to reaching a minimum at 60°S and then increasing towards the South Pole. Consistent with previous analyses based on climate simulations and observations, this enhanced Arctic Amplification appears as an inherent characteristic for the Arctic region (Pithan and Mauritsen, 2014). From March to August, the reverse signal shows the maximum warming close to 70°S, decreasing towards to tropical region, and lacking the enhanced warming at the northern high latitudes.

225 The main reason for winter (DJF) Arctic Amplification pointed by Serreze et al., (2009) is largely driven by changes in sea ice, allowing for intense heat transfers from the ocean to the atmosphere. During boreal summer, when Arctic warming is not prominent and solar radiation is maximal, the energy is used to melt sea ice and increase the sensible heat content of the upper ocean. The atmosphere heats the ocean during summer whereas the flux of heat is reverse in winter. The sea ice loss in summer allows a large warming of the upper ocean but the atmospheric warming at the surface or lower troposphere is modest (promoting more open water). The excess heat stored in the upper ocean is subsequently released to the atmosphere during winter (Serreze et al., 2009). According to Lu and Cai, (2009), in summertime the positive surface albedo feedback is mainly canceled out by the negative cloud radiative forcing feedback. The positive surface albedo feedback is relatively much weaker in winter when compared to its counterpart in summer, therefore it does not contribute to the pronounced polar amplification in winter.

235 For southern high latitudes, a pronounced warming appears from March to August (boreal summer and spring), predominantly close from 70°S. This enhanced warming tends to decrease in the direction of the South Pole. This pattern is similar to the one obtained by Goosse and Renssen, (2001). The authors used a coupled climate model to investigate the response of the Southern Ocean to an increase

240 in GHG concentration. They found that the response could occur separately in two distinct phases. At the first moment, the ocean damps the surface warming (because of its large heat capacity). Then, after 100 years of run simulation, the warming is enhanced due to a positive feedback that is linked to a stronger oceanic meridional heat transport toward the southern ocean.

When comparing the seasonal response to CO<sub>2</sub> forcing among CMIP5 and CMIP6 models, for 245 boreal winter (DJF), the enhanced Arctic warming at 75-90° N is shown to be a robust feature of all CMIP5 and CMIP6 climate models simulations presented here. For high Northern Hemisphere (high southern Hemisphere) the warming (difference between piControl and 4xCO<sub>2</sub>) ranged from 10 K to 39 K (-0.5 K – 13 K). CAMS-CSM1-0 (CMIP6) and INMC-CM4 (CMIP5) presented the lowest warming, close from 12\_K for Northern high latitudes. On the other hand, IPSL-CM6A-LR (CMIP6), HadGEM2- 250 ES (CMIP5) and CanESM5 (CMIP6) outputs presented warming almost twice as large, with a high amplification close from 30 K. BESM model, for winter (DJF) season, presented Polar Amplification for Northern high latitudes, close from 27 K.

One interesting feature shown in Figure 2 is related to the maximum Arctic warming obtained in different simulations. Many models have shown that the maximum warming does not always occur at 255 highest northern latitudes instead, it occurs between 80° N - 85° N decreasing toward 90° N. According to Holland and Bitz, (2003) the localization of the maximum warming varies widely among CMIP outputs, but models with high polar amplification generally presented a maximum warming over the Arctic Basin. Therefore, we suggest that the spatial distribution of maximum Arctic Amplification can be closely related to sea ice conditions through a sea ice albedo feedback, and this region (Arctic Basin) 260 presents the major taxes of decrease in sea ice concentration. Similar result was found for the sea ice simulation from BESM model, as discussed below (Figure 5 and Table 1). Additionally, Casagrande et al., (2016), using BESM-OA V2.3 model, showed that the sea ice spatial pattern could vary largely between CMIP5 models, especially in frontiers areas.

For the southern high latitudes, in wintertime (DJF- Figure 2), the warming decreases to close to 265 60° S for most CMIP5 and CMIP6 models, increasing toward South Pole, with the maximum warming close to 11 K. The minimum warming is registered by GFDL-ESM2M and GFDL-ESM2G, both from CMIP5 simulations (close to 0K in 60° S) and the maximum south polar amplification between models

---

is presented by E3SM-1, close to 90° S. In summer (JJA), the compared response to CO<sub>2</sub> forcing in CMIP5 models is amplified (damped) at southern (northern) hemisphere. A pronounced amplification was found close to 70°S with a range of 1 K to 17K, decreasing towards the South Pole. In this region the maximum was obtained by BESM-OA V2.5 model, close to 13K.

The pronounced seasonality of near surface warming in Polar Regions has been found in observations (Bekryaev et al., 2010) and climate simulations (Holland and Bitz, 2003), but less emphasis has been placed in the vertical structure of the atmosphere. To understand if this enhanced warming occurs only in surface or also well above, Figure 4 presents results obtained with three different CMIP5 models with moderate (BESM-OA V2.5/MPI-ESM-LR) and low (NCAR-CCSM4) Polar Amplification (based on Figure 2 and Figure 3).

Figure 4 shows evidence of temperature amplification well above the surface with enhanced warming during the cold season for both, northern and southern high latitudes. Snow and ice feedback cannot explain the warming above the lowermost part of the atmosphere because this feedback is expected to primarily affect the air temperature near surface. Part of the vertical warming may be explained by physical mechanisms that induce to warming as changes in the atmospheric heat transport into the Arctic. According to Graversen et al., (2008), a substantial proportion of the vertical warming can be caused by changes in this variable, especially in summertime (JJA). Graversen and Wang (2009) used an idealized numerical experiment (doubling CO<sub>2</sub>) with a climate model that has no ice albedo feedback. Their results also revealed a polar warming as a response to anthropogenic forcing (doubling CO<sub>2</sub>). It was found that the enhanced Arctic warming is due to an increase of the atmospheric northward transport of heat and moisture. These results are supported by observational and numerical analyses (Graversen et al., 2014; Graversen et al., 2008). In addition to ice-albedo feedback, the strength of the atmospheric stratification is an important factor to explain the vertical warming. The troposphere is more stably stratified in high latitudes. An increase in GHG forcing generates an increase in downwelling long-wave radiation at the surface, consequently causing warming, which in Polar Regions is confined to the lower troposphere (Graversen et al., 2014; Graversen et al., 2009).

When examining Arctic warming at different levels computed by the three different models shown in Figure 4, we find that MPI-ESM-LR presented the strongest warming in both, near surface temperature and at high levels. Similar behavior is found at tropical regions, with robust warming at high levels (400-200 hPa). Holland and Bitz, (2003) suggested that sea ice conditions are more important than continental ice and snow cover to enhanced polar warming. According to these authors, models with relatively thin sea ice in the control run tend to have higher warming. The same feature was

300 found in BESM-OA V2.5. According to Casagrande et al. (2016) and Casagrande (2016), the last  
version of BESM model (Version 2.5) is considered to be a climate model with high polar amplification  
exhibiting thin sea ice conditions on the control run. This occurs, in part, because of the new surface  
scheme based on Jimenez and Dudhia, (2012) and the microphysics of Ferrier et al. (2002). The  
305 advantage of these changes in the BESM's last version is an improvement in the representation of  
precipitation, wind and humidity at tropical regions. Comparatively, NCAR-CCSM4 is considered a  
model with moderate polar amplification for both, Northern and Southern Ocean. The warming at high  
levels in boreal summer is not as amplified as in boreal winter. These results are in agreement with  
Holland and Bitz, (2003).

Figure 5 shows, under the largest future GHG (4xCO<sub>2</sub>), the spatial pattern of sea ice changes for  
310 both, Arctic and Antarctic (difference between sea ice concentration for the last 30 years of  
abrupt4xCO<sub>2</sub> numerical experiment and the last 30 years of the piControl run). The maximum of the  
Arctic warming obtained from observations (Figure 1) and different CMIP5 simulations (Figure 2 and  
Figure 3) occurs in boreal winter (DJF).

According to Figure 2, the following models, in descending order, appear as having greater  
315 amplification: IPSL-CM6A-LR (CMIP6), HadGEM2-ES (CMIP5) and CanESM5 (CMIP6). Similar  
response, for the same period, is observed in Figure 5 and Figure 6, related to sea ice changes. Figure 6  
shows the climatology of maximum and minimum sea ice area for the last 30 years of the abrupt 4xCO<sub>2</sub>  
numerical experiment minus the last 30 years of the piControl run. For the Arctic, in March, EC-Earth3-  
Veg (NCAR-CCSM4) shows the highest (lowest) value, close to 15 x 10<sup>6</sup> km<sup>-2</sup> (3 x 10<sup>6</sup> km<sup>-2</sup>). For  
320 September month, in agreement with Figure 2, the Polar Amplification is not evident as in the cold  
period. For the Antarctica (Figure 6), in the cold period (September), the difference between abrupt  
4xCO<sub>2</sub> numerical experiment and the piControl run is higher for models with enhanced Polar  
Amplification, as FGOALS-S2 (13 x 10<sup>6</sup> km<sup>-2</sup>). Both, Figure 5, Figure 6 and Table 1 are in agreement  
with Figure 2, showing that the large decrease in sea ice concentration is more evident in models with  
325 great Polar Amplification, and for the same range of latitude (75° N – 90° N). The end of melting period  
(when sea ice reaches its minimum annual value) for all models shows sea ice-free conditions (Table 1).  
Models that have strong Polar Amplification, also exhibit expressive changes in the annual amplitude of  
sea ice with outstanding ice-free condition from May to December (EC-Earth3-Veg) and June to

---


December (HadGEM2-ES). Then, the end of melting period is expected early, likely, associated a large  
330 decrease in sea ice thickness, which contributes to a delay in sea ice formation. For BESM-OA V2.5,  
Arctic ice-free conditions are found from August to November. We suggest that, the Arctic will become  
covered only by first year sea ice (more vulnerable to melting), making the region more sensitive  
thermodynamically and dynamically to temperature changes. These evidences, corroborates with the  
theory, that the Arctic Polar Amplification is closely linked to sea ice albedo feedback. For Antarctica,  
335 however, the same physical processes cannot be used to explain the Polar Amplification (as discussed  
previously). Although, according to Figure 2 and Figure 4, there is a small indication of the contribution  
of sea ice albedo feedback in Antarctic Polar Amplification, however, this still remain as a open  
discussion and we suggest that is important to consider the contribution of the ice sheet in Polar  
Amplification.

340 Previous researchers, using observational and modeling dataset, have found that shrinking of sea  
ice (Figure 5) and enhanced Arctic warming may affect the middle latitudes (Coumou et al., 2018;  
Screen, 2017; Walsh, 2014). According to Walsh (2014), the AA affects by the weakening the west-to-  
east wind speed in the upper atmosphere, by increasing the frequency of wintertime blocking events that  
in turn lead to persistence or slower propagation of anomalous temperature in middle latitudes, and by  
345 increasing in continental snow cover that can in turn influence the atmospheric circulation. Finally, in  
view of the results it is important to consider the limitations and differences among each climate model  
in order to improve the understanding of the physical process in climate simulations that represent large  
bias among the models belonging to CMIP5 project.

350

(a)


(b)




355 **Figure 2. Seasonal zonal mean surface temperature differences (K) for the last 30 years of Abrupt4xCO<sub>2</sub> numerical experiment minus the last 30 years of the piControl run for CMIP5 and CMIP6 models for (a) Winter (DJF), (b) Spring (MAM), (c) Summer (JJA) and (d) autumn (SON).**

360


(a)




(b)




(c)



(d)




**Figure 3. Seasonal zonal mean surface temperature differences (K) for the last 30 years of Abrupt4xCO<sub>2</sub> numerical experiment minus the last 30 years of the piControl run for CMIP5 and CMIP6 models, box plot in 75°N – 90°N (left) and 60°S -80°S (right) for (a) Winter (DJF), (b) Spring (MAM), (c) Summer (JJA) and (d) autumn (SON).**




**Figure 4. Zonal-average atmosphere temperature changes, in °C (Abrupt 4xCO<sub>2</sub> minus piControl) at each pressure level, in mb (solid line) for the last 30 years run for (a) BESM OA V2.5, (b) NCAR-CCSM4 and (c) MPI-ESM-LR model, in DJF (left) and JJA (right) columns.**





375 **Figure 5. Sea ice concentration for the last 30 years of abrupt4xCO<sub>2</sub> numerical experiment minus the last 30 years of the piControl run for the following models: (a) BESM-OA V2.5, (b) NCAR-CCSM4, (c) FGOALS-S2, (d) CanESM5, (e) HadGEM2-ES (f) EC-Earth3-Veg in March (left column).**



**Figure 6. Climatological maximum and minimum Sea ice area (million square kilometers) for the last 30 years of the abrupt 4xCO<sub>2</sub> numerical experiment minus the last 30 years of the piControl run for CMIP5 and CMIP6 models (a) Arctic, black colors represents the March month and Gray colors represent September month (b) Antarctic: Black colors represents the September month and Gray colors represent February month.**

385

CMIP6 Models	Arctic		Antarctic		CMIP5 Models		Arctic		Antarctic	
	March	Sept	Feb	Sept	March	Sept	Feb	Sept	March	Sept
ACCESS-CM2	14.2	5.3	0.5	13.4	BESM-OA2L5	16	3.5	1	29	
	4xCO2	2	Ice-Free [Jan-Dec]	4xCO2		Ice-Free [Jan-Mar]	11.5	Ice-Free [Aug-Nov]	4xCO2	Ice-Free [Feb-Apr]
CM5-CSM1-0	17.1	7.9	0.5	12	ACCESS-S	14	6	4.5	17	
	4xCO2	12.5	Ice-Free [Jan-Mar]	4xCO2		Ice-Free [Jan-Mar]	7	Ice-Free [Jul-Nov]	4xCO2	1
CanESM5	15	7	6	20	GFDL-ESM2M	14	7	0	9	
	4xCO2	1.2	Ice-Free [Mar-Dec]	4xCO2		Ice-Free [Feb-Mar]	9	Ice-Free [Aug-Sept]	4xCO2	Ice-Free [Feb-Mar]
CMCC-CM2-SR5	15.5	1.5	0.6	14	IPSL-CM5-LRM	13	9	1	17	
	4xCO2	1	Ice-Free [Mar-Dec]	4xCO2		Ice-Free [Mar-Apr]	0.5	Ice-Free [Jul-Oct]	4xCO2	Ice-Free [Jan-Mar]
CNRM-ESM2-1	15.2	8	0.5	16	MIROC-ESM	13	8	1	15	
	4xCO2	7.1	Ice-Free [Jul-Dec]	4xCO2		Ice-Free [Jan-Apr]	0.8	Ice-Free [May-Dec]	4xCO2	Ice-Free
ACCESS-ESM1-5	14.7	5.5	0.5	14	MPI-ESM-LR	12	8	1	15	
	4xCO2	4.7	Ice-Free [Jul-Dec]	4xCO2		Ice-Free [Feb-Mar]	3.5	Ice-Free [Jan-Dec]	4xCO2	Ice-Free [Jan-Mar]
ESM1-0	12	17	2	17	NCAR-CCSM4	13	17	1	22	
	4xCO2	Ice-Free [All Year]	4xCO2	Ice-Free		10	Ice-Free [Aug-Oct]	4xCO2	1	
EC-Earth3-Veg	15	7.8	0.2	10.5	CanESM2	15	5.8	4	27	
	4xCO2	0.5	Ice-Free [Mar-Dec]	4xCO2		Ice-Free	5	Ice-Free [Jul-Nov]	4xCO2	0.5
FGOALS-G3	15	8	2.0	19	FGOALS-s2	12	7	6	22	
	4xCO2	16.3	0.5	4xCO2		1	12	Ice-Free [Apr-Oct]	4xCO2	1
GISS-E2-1-H	22	21	2.5	14	GFDL-ESM2G	18	2.8	0	11.5	
	4xCO2	15.5	Ice-Free [Aug-Sept]	4xCO2		Ice-Free	12	Ice-Free [Aug-Sept]	4xCO2	Ice-Free [Jan-Apr]
INM-CM4-8	14	7	0.5	10	GISS-E2-H	15	2.8	0	7.5	
	4xCO2	9	3.5	4xCO2		Ice-Free [Jan-Mar]	9.5	Ice-Free [Aug-Oct]	4xCO2	Ice-Free
MIROC6	11	3	0.5	17	HadGEM2-ES	16	4.8	1.2	14.2	
	4xCO2	6.5	Ice-Free [Jul-Dec]	4xCO2		Ice-Free	2.3	Ice-Free [Jan-Dec]	4xCO2	Ice-Free
MIROC-ES2L	12	5	3	3	MIROC5	13	7	0	7	
	4xCO2	8	Ice-Free [Jul-Nov]	4xCO2		Ice-Free [Jan-Mar]	7	Ice-Free [Jul-Nov]	4xCO2	Ice-Free [Jan-Mar]
MPI-ESM1-2-LR	12	5.5	0.5	11	MPI-ESM-P	12	5.5	0.9	12	
	4xCO2	7	Ice-Free [Jul-Dec]	4xCO2		Ice-Free [Jan-Mar]	6.5	Ice-Free [Jul-Dec]	4xCO2	Ice-Free [Jan-Mar]
MRI-ESM2-0	15	5	3	20.5	MRI-CGCM3	21	7	3	19	
	4xCO2	9	Ice-Free [Jul-Dec]	4xCO2		Ice-Free [Feb]	13	Ice-Free [Jul-Dec]	4xCO2	Ice-Free [Feb]

390

**Table 1. Climatological maximum and minimum Sea ice area (million square kilometers) for the last 30 years of the abrupt 4xCO<sub>2</sub> numerical experiment and the last 30 years of the piControl run for CMIP5 and CMIP6 models.**

---

## 4 Conclusion

Polar amplification is possibly one of the most important sensitive indicators of climate change. Robust patterns of near-surface temperature response to global warming at high latitudes have been identified in recent studies (Smith et al., 2019; Stuecker et al., 2018; Pithan and Mauritsen, 2014). For northern high latitudes, the shrinkage of sea ice as response to increase of GHG is one of the most cited reasons (Serreze and Barry, 2011; Kumar et al., 2010; Screen and Simmonds, 2010). Here we analyzed the seasonality of polar amplification using some CMIP5 coupled climate models in a quadrupling CO<sub>2</sub> numerical experiment for both, North and South Hemispheres. Our results showed that the Polar Regions are much more vulnerable to a large warming due to an increase in atmospheric CO<sub>2</sub> forcing, than the rest of the world, particularly during the cold season. For northern high latitudes, the feedback albedo sea ice contributes to decrease in sea ice cover, exposing new expanses of ocean and land surfaces (leading to greater solar absorption), thus amplifying the accelerated warming and driving future melting. Despite of the asymmetry in warming between Arctic and Antarctic, both poles show systematically polar amplification in all climate models. Different physical processes acts to explain the sensibilities between poles. While in Northern high latitudes the warming is closely related to sea ice albedo feedback, in southern high latitudes the amplification is related to thermal inertia, combination of changes in winds and ozone depletion. We detected three climate models as having high amplification, in cold season for Arctic: IPSL-CM6A-LR (CMIP6), HadGEM2-ES (CMIP5) and CanESM5 (CMIP6). For South Hemisphere, in the cold season (JJA), the Climate Model identified as having high polar amplification were: IPSL-CM6A-LR (CMIP6), CanESM5(CMIP6) and FGOALS-s2 (CMIP5). For high Northern Hemisphere (high southern Hemisphere) the warming ranged from 10 K to 39 K (-0.5 K – 13 K), INM-CM4 (CMIP5) presents the lowest warming, close from 10 K for Northern high latitudes. For Antarctica, the maximum warming, close to 14 K is presented by FGOALS-s2, close to 70° S. The vertical profiles of air temperature showed stronger warming at the surface, particularly for northern high latitudes, indicating the effectiveness of the albedo-sea ice feedback. Furthermore, we evaluated the linkage between sea ice changes and Polar Amplification from different CMIP5 models. We found that large decreases in sea ice concentration are more evident in models with great Polar Amplification, and for the same range of latitude (75° N – 90° N). We suggest, according to our results,

---

420 that the large difference between models might be related to sea ice initial conditions. Therefore, those  
differences are also related to the parameterizations used to represent changes in clouds and energy  
balance. The coupled ocean-atmosphere-cryosphere physical processes involved in high-latitudes  
climate changes are fully inter-dependent with complicated structures contending with each other at  
425 a leak in reproducibility by the numerical climate models, especially at southern regions. The sparse and  
short data record does not help also. Nevertheless, even with inherent limitations and uncertainties, the  
Global Climate Models are the most powerful tools available for simulating the climatic response to  
GHG forcing and to providing future scenarios to community.

## References

430

Alexeev, V. A., Langen, P. L. and Bates, J. R.: Polar amplification of surface warming on an aquaplanet  
in “ghost forcing” experiments without sea ice feedbacks, *Climate Dynamics*, 24(7–8), 655–666,  
doi:10.1007/s00382-005-0018-3, 2005.

435 Ambaum, M. H. P., Hoskins, B. J. and Stephenson, D. B.: Arctic Oscillation or North Atlantic  
Oscillation?, *Journal of Climate*, 14(13), 3495-3507, doi: 10.1175/1520-  
0442(2001)014<3495:AOONAO>2.0.CO;2, 2001.

Bader, David C.; Leung, Ruby; Taylor, Mark; McCoy, Renata B. E3SM-Project E3SM1.0 model output  
440 prepared for CMIP6 CMIP abrupt-4xCO2. Version YYYYMMDD<sup>[1]</sup>.Earth System Grid  
Federation. <https://doi.org/10.22033/ESGF/CMIP6.4491>, 2019.

Bao, Qing et al. The flexible global ocean-atmosphere-land system model, spectral version 2:  
FGOALS-s2. *Advances in Atmospheric Sciences*, v. 30, n. 3, p. 561-576, 2013.

445

---

Bekryaev, R. V., Polyakov, I. V. and Alexeev, V. A.: Role of Polar Amplification in Long-Term Surface Air Temperature Variations and Modern Arctic Warming, *J. Climate*, 23(14), 3888–3906, doi:10.1175/2010JCLI3297.1, 2010.

450

Bi, D., Dix, M., Marsland, S., O’Farrell, S., Rashid, H., Uotila, P., Hirst, A., Kowalczyk, E., Golebiewski, M., Sullivan, A., Yan, H., Hannah, N., Franklin, C., Sun, Z., Vohralik, P., Watterson, I., Zhou, X., Fiedler, R., Collier, M., Ma, Y., Noonan, J., Stevens, L., Uhe, P., Zhu, H., Griffies, S., Hill, R., Harris, C. and Puri, K.: The ACCESS coupled model: description, control climate and evaluation, *a*, 455 63(1), 41–64, doi:10.22499/2.6301.004, 2013.

Bintanja, R., van Oldenborgh, G. J. and Katsman, C. A.: The effect of increased fresh water from Antarctic ice shelves on future trends in Antarctic sea ice, *Ann. Glaciol.*, 56(69), 120–126, doi:10.3189/2015AoG69A001, 2015.

460

Bintanja, R., van Oldenborgh, G. J., Drijfhout, S. S., Wouters, B. and Katsman, C. A.: Important role for ocean warming and increased ice-shelf melt in Antarctic sea-ice expansion, *Nature Geosci*, 6(5), 376–379, doi:10.1038/ngeo1767, 2013.

465 Capistrano, Vinicius et al. Assessing the performance of climate change simulation results from BESM-OA2. 5 compared with a CMIP5 model ensemble. *Geoscientific Model Development*, v. 13, n. 5, p. 2277-2296, 2020.

470 Casagrande, F., Nobre, P., de Souza, R. B., Marquez, A. L., Tourigny, E., Capistrano, V. and Mello, R. L.: Arctic Sea Ice: Decadal Simulations and Future Scenarios Using BESM-OA, *ACS*, 06(02), 351–366, doi:10.4236/acs.2016.62029, 2016.

Casagrande, F.: Sea ice study and Arctic Polar Amplification using BESM model, PhD, <http://mtcm21b.sid.inpe.br/col/sid.inpe.br/mtcm21b/2016/05.12.04.17/doc/publicacao.pdf>, 2016.

---

Chylek, P. et al. Observed and model simulated 20th century Arctic temperature variability: Canadian earth system model CanESM2. *Atmos. Chem. Phys. Discuss*, v. 11, n. 22, p. 893-22, 2011.

Cohen J L , Furtado J C , Barlow M , et al. Asymmetric seasonal temperature trends[J]. *Geophysical Research Letter*, 2012.

Collier, M. and Uhe, P.: CMIP5 datasets from the ACCESS1.0 and ACCESS1.3 coupled climate models, CAWCR Technical Report, 2012.

485 Collins, W. J., and Coauthors : Evaluation of the HadGEM2 model. Tech. Note HCTN 74, Met Off. Hadley Cent., Exeter, U. K, 2008.

Coumou, D., Di Capua, G., Vavrus, S., Wang, L., & Wang, S. The influence of Arctic amplification on mid-latitude summer circulation. *Nature Communications*, 9(1), 1-12, 2018.

490

Cvijanovic, I., Caldeira, K., & MacMartin, D. G: Impacts of ocean albedo alteration on Arctic sea ice restoration and Northern Hemisphere climate. *Environmental Research Letters*, 10(4), 044020, 2015.

Delworth, T. L., and Coauthors : GFDL's CM2 global coupled climate models. Part I: Formulation and simulation characteristics. *J. Climate*, 19, 643–674, 2006.

495

Dethloff, K., Handorf, D., Jaiser, R., Rinke, A. and Klinghammer, P.: Dynamical mechanisms of Arctic amplification: Dynamical mechanisms of Arctic amplification, *Ann. N.Y. Acad. Sci.*, 1436(1), 184–194, doi:10.1111/nyas.13698, 2019.

500

Dufresne, J.-L., Foujols, M.-A., Denvil, S., Caubel, A., Marti, O., Aumont, O., Balkanski, Y., Bekki, S., Bellenger, H., Benschila, R., Bony, S., Bopp, L., Braconnot, P., Brockmann, P., Cadule, P., Cheruy, F.,

---

Codron, F., Cozic, A., Cugnet, D., de Noblet, N., Duvel, J.-P., Ethé, C., Fairhead, L., Fichet, T., Flavoni, S., Friedlingstein, P., Grandpeix, J.-Y., Guez, L., Guilyardi, E., Hauglustaine, D., Hourdin, F.,  
505 Idelkadi, A., Ghattas, J., Joussaume, S., Kageyama, M., Krinner, G., Labetoulle, S., Lahellec, A.,  
Lefebvre, M.-P., Lefevre, F., Levy, C., Li, Z. X., Lloyd, J., Lott, F., Madec, G., Mancip, M., Marchand,  
M., Masson, S., Meurdesoif, Y., Mignot, J., Musat, I., Parouty, S., Polcher, J., Rio, C., Schulz, M.,  
Swingedouw, D., Szopa, S., Talandier, C., Terray, P., Viovy, N. and Vuichard, N.: Climate change  
510 2123–2165, doi:10.1007/s00382-012-1636-1, 2013.

Eyring, Veronika et al. Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6)  
experimental design and organization. *Geoscientific Model Development*, v. 9, n.5, p. 1937-1958, 2016.

515 Ferrier, B. S., Jin, Y., Lin, Y., Black, T., Rogers, E., and DiMego, G.: Implementation of a 527 new  
grid-scale cloud and precipitation scheme in the NCEP Eta model, American Meteor Society, 19th  
Conf. on weather Analysis and Forecasting/15th Conf. on Numerical Weather Prediction, 280–283,  
2002.

520 Fiedler, Stephanie; Stevens, Bjorn; Wieners, Karl-Hermann; Giorgetta, Marco; Reick, Christian;  
Jungclaus, Johann; Esch, Monika; Bittner, Matthias; Legutke, Stephanie; Schupfner, Martin;  
Wachsmann, Fabian; Gayler, Veronika; Haak, Helmuth; de Vrese, Philipp; Lorenz, Stephan; Raddatz,  
Thomas; Mauritsen, Thorsten; von Storch, Jin-Song; Mikolajewicz, Uwe; Behrens, Jörg; Brovkin,  
Victor; Claussen, Martin; Crueger, Traute; Fast, Irina; Hagemann, Stefan; Hohenegger, Cathy; Jahns,  
525 Thomas; Kloster, Silvia; Kinne, Stefan; Lasslop, Gitta; Kornblueh, Luis; Marotzke, Jochem; Matei,  
Daniela; Meraner, Katharina; Modali, Kameswarrao; Müller, Wolfgang; Nabel, Julia; Notz, Dirk;  
Peters, Karsten; Pincus, Robert; Pohlmann, Holger; Pongratz, Julia; Rast, Sebastian; Schmidt, Hauke;  
Schnur, Reiner; Schulzweida, Uwe; Six, Katharina; Voigt, Aiko; Roeckner, Erich (2019). *MPI-M MPI-  
ESM1.2-LR model output prepared for CMIP6 RFMIP piClim-control*. Version YYYYMMDD<sup>[1]</sup>. Earth  
530 System Grid Federation. <https://doi.org/10.22033/ESGF/CMIP6.6662>

---

535 Figuroa, S. N., Bonatti, J. P., Kubota, P. Y., Grell, G. A., Morrison, H., Barros, S. R. M., Fernandez, J. P. R., Ramirez, E., Siqueira, L., Luzia, G., Silva, J., Silva, J. R., Pendharkar, J., Capistrano, V. B., Alvim, D. S., Enoré, D. P., Diniz, F. L. R., Satyamurti, P., Cavalcanti, I. F. A., Nobre, P., Barbosa, H. M. J., Mendes, C. L. and Panetta, J.: The Brazilian Global Atmospheric Model (BAM): Performance for Tropical Rainfall Forecasting and Sensitivity to Convective Scheme and Horizontal Resolution, *Wea. Forecasting*, 31(5), 1547–1572, doi:10.1175/WAF-D-16-0062.1, 2016.

540 Fogli, Pier Giuseppe; Iovino, Doroteaciro; Lovato, Tomas. *CMCC CMCC-CM2-SR5 model output prepared for CMIP6 OMIP omip2*. Version YYYYMMDD<sup>[1]</sup>. Earth System Grid Federation. <https://doi.org/10.22033/ESGF/CMIP6.13236>, 2019.

545 Gent, P. R., Danabasoglu, G., Donner, L. J., Holland, M. M., Hunke, E. C., Jayne, S. R., Lawrence, D. M., Neale, R. B., Rasch, P. J., Vertenstein, M., Worley, P. H., Yang, Z.-L. and Zhang, M.: The Community Climate System Model Version 4, *J. Climate*, 24(19), 4973–4991, doi:10.1175/2011JCLI4083.1, 2011.

550 Giarolla, E., Siqueira, L. S. P., Bottino, M. J., Malagutti, M., Capistrano, V. B. and Nobre, P.: Equatorial Atlantic Ocean dynamics in a coupled ocean–atmosphere model simulation, *Ocean Dynamics*, 65(6), 831–843, doi:10.1007/s10236-015-0836-8, 2015.

Giorgetta, Marco A. et al. Climate and carbon cycle changes from 1850 to 2100 in MPI-ESM simulations for the Coupled Model Intercomparison Project phase 5. *Journal of Advances in Modeling Earth Systems*, v. 5, n. 3, p. 572-597, 2013.

555

Goosse, H. and Renssen, H.: A two-phase response of the Southern Ocean to an increase in greenhouse gas concentrations, *Geophys. Res. Lett.*, 28(18), 3469–3472, doi:10.1029/2001GL013525, 2001.



- 
- 560 Graversen, R. G. and Wang, M.: Polar amplification in a coupled climate model with locked albedo, *Clim Dyn*, 33(5), 629–643, doi:10.1007/s00382-009-0535-6, 2009.
- Graversen, R. G., Langen, P. L. and Mauritsen, T.: Polar Amplification in CCSM4: Contributions from the Lapse Rate and Surface Albedo Feedbacks, *J. Climate*, 27(12), 4433–4450, doi:10.1175/JCLI-D-13-00551.1, 2014.
- 565 Graversen, R. G., Mauritsen, T., Tjernström, M., Källén, E. and Svensson, G.: Vertical structure of recent Arctic warming, *Nature*, 451(7174), 53–56, doi:10.1038/nature06502, 2008.
- Griffies, S. M. (2012). Elements of the modular ocean model (MOM). *GFDL Ocean Group Tech. Rep*, 7, 620.
- 570 Griffies, S. M.: Elements of MOM4p1. NOAA/Geophysical Fluid Dynamics Laboratory Ocean Group Tech. Rep. 6, 444 pp., 2009.
- 575 Haima, T. et al. Description of the MIROC-ES2L Earth system model and evaluation of its climate—Biogeochemical processes and feedbacks. *Geoscientific Model Development Discussions*, v. 2019, p. 1-73, 2019.
- Holland, M. M. and Bitz, C. M.: Polar amplification of climate change in coupled models, *Climate Dynamics*, 21(3–4), 221–232, doi:10.1007/s00382-003-0332-6, 2003.
- 580 Honda M , Inoue J , Yamane S .Influence of low Arctic sea-ice minima on anomalously cold Eurasian winters[J]. *Geophysical Research Letters*, 36(36):262-275, 2009.
- 585 Hunke, E. C. and Dukowicz, J. K.: An Elastic–Viscous–Plastic Model for Sea Ice Dynamics, *J. of Physical Oceanography*, 27, 19, 1997.

- 
- Jiménez, P. A., Dudhia, J., González-Rouco, J. F., Navarro, J., Mon- távez, J. P., and García-Bustamante, E.: A Revised Scheme for the WRF Surface Layer Formulation, *Mon. Weather Rev.*, 140, 590 898–918, <https://doi.org/10.1175/MWR-D-11-00056.1>, 2012.
- Kumar, A., Perlwitz, J., Eischeid, J., Quan, X., Xu, T., Zhang, T., Hoerling, M., Jha, B. and Wang, W.: Contribution of sea ice loss to Arctic amplification: Sea ice loss and Arctic Amplification, *Geophys. Res. Lett.*, 37(21), n/a-n/a, doi:10.1029/2010GL045022, 2010.
- 595 LI, Lijuan et al. The Flexible Global Ocean–Atmosphere–Land System Model Grid□Point Version 3 (FGOALS-g3): Description and Evaluation. *Journal of Advances in Modeling Earth Systems*, 2020.
- Lu, J. and Cai, M.: Seasonality of polar surface warming amplification in climate simulations, *Geophys. Res. Lett.*, 36(16), L16704, doi:10.1029/2009GL040133, 2009.
- 600 Manabe, S., Wetherald, R. T., Milly, P. C. D., Delworth, T. L. and Stouffer, R. J.: Century-Scale Change in Water Availability: CO<sub>2</sub> -Quadrupling Experiment, *Climatic Change*, 64(1/2), 59–76, 2004.
- 605 Mann, M. E., G. A. Schmidt, S. K. Miller, and A. N. LeGrande: Potential biases in inferring Holocene temperature trends from long-term borehole information, *Geophys. Res. Lett.*, 36, L05708, doi:10.1029/2008GL036354.2009.
- Marshall, J., Armour, K. C., Scott, J. R., Kostov, Y., Hausmann, U., Ferreira, D., Shepherd, T. G. and 610 Bitz, C. M.: The ocean’s role in polar climate change: asymmetric Arctic and Antarctic responses to greenhouse gas and ozone forcing, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 372(2019), 20130040–20130040, doi:10.1098/rsta.2013.0040, 2014.

- 
- 615 Masson-Delmotte, V., Kageyama, M., Braconnot, P., Charbit, S., Krinner, G., Ritz, C., ... & Gladstone,  
R. M.: Past and future polar amplification of climate change: climate model intercomparisons and ice-  
core constraints. *Climate Dynamics*, 26(5), 513-529, 2006.
- Mori M , Watanabe M , Shiogama H , et al. Robust Arctic sea-ice influence on the frequent Eurasian  
620 cold winters in past decades[J]. *Nature Geoscience*, 7(12):869-873, 2014.
- Nobre, P., Siqueira, L. S. P., de Almeida, R. A. F., Malagutti, M., Giarolla, E., Castelão, G. P., Bottino,  
M. J., Kubota, P., Figueroa, S. N., Costa, M. C., Baptista, M., Irber, L. and Marcondes, G. G.: Climate  
Simulation and Change in the Brazilian Climate Model, *J. Climate*, 26(17), 6716–6732,  
625 doi:10.1175/JCLI-D-12-00580.1, 2013.
- O'ishi, R., Abe-Ouchi, I. C. Prentice, and S. Sitch: Vegetation dynamics and plant CO2 responses as  
positive feedbacks in a greenhouse world, *Geophys. Res. Lett.*, 36, L11706,  
doi:10.1029/2009GL038217, 2009.
- 630 Pedersen, R. A., Cvijanovic, I., Langen, P. L., & Vinther, B. M : The impact of regional Arctic sea ice  
loss on atmospheric circulation and the NAO. *Journal of Climate*, 29(2), 889-902,doi: 10.1175/JCLI-D-  
15-0315.1, 2019.
- 635 Pithan, F. and Mauritsen, T.: Arctic amplification dominated by temperature feedbacks in contemporary  
climate models, *Nature Geosci*, 7(3), 181–184, doi:10.1038/ngeo2071, 2014.
- Polyakov, I. V., Alexeev, V. A., Belchansky, G. I., Dmitrenko, I. A., Ivanov, V. V., Kirillov, S. A.,  
Korablev, A. A., Steele, M., Timokhov, L. A. and Yashayaev, I.: Arctic Ocean Freshwater Changes  
640 over the Past 100 Years and Their Causes, *J. Climate*, 21(2), 364–384, doi:10.1175/2007JCLI1748.1,  
2008.

- 
- 645 Polyakov, I. V., Pnyushkov, A. V., Alkire, M. B., Ashik, I. M., Baumann, T. M., Carmack, E. C., Goszczko, I., Guthrie, J., Ivanov, V. V., Kanzow, T., Krishfield, R., Kwok, R., Sundfjord, A., Morison, J., Rember, R. and Yulin, A.: Greater role for Atlantic inflows on sea-ice loss in the Eurasian Basin of the Arctic Ocean, *Science*, 356(6335), 285–291, doi:10.1126/science.aai8204, 2017.
- 650 Polyakov, I. V., Timokhov, L. A., Alexeev, V. A., Bacon, S., Dmitrenko, I. A., Fortier, L., Frolov, I. E., Gascard, J.-C., Hansen, E., Ivanov, V. V., Laxon, S., Mauritzen, C., Perovich, D., Shimada, K., Simmons, H. L., Sokolov, V. T., Steele, M. and Toole, J.: Arctic Ocean Warming Contributes to Reduced Polar Ice Cap, *J. Phys. Oceanogr.*, 40(12), 2743–2756, doi:10.1175/2010JPO4339.1, 2010.
- Rigor, I. G.: Response of Sea Ice to the Arctic Oscillation, *Journal of Climate*, 15(18), 2648-2663, 2002.
- 655 Rong, Xinyao. *CAMS CAMS\_CSM1.0 model output prepared for CMIP6 CMIP*  
*IpctCO2*. Version YYYYMMDD<sup>[1]</sup>.Earth System Grid  
Federation. <https://doi.org/10.22033/ESGF/CMIP6.9701> , 2019.
- 660 Salzmann, M.: The polar amplification asymmetry: role of Antarctic surface height, *Earth Syst. Dynam.*, 8(2), 323–336, doi:10.5194/esd-8-323-2017, 2017.
- Schmidt, Gavin A. et al. Present-day atmospheric simulations using GISS ModelE: Comparison to in situ, satellite, and reanalysis data. *Journal of Climate*, v. 19, n. 2, p. 153-192, 2006.
- 665 Screen, J. A. and Simmonds, I.: The central role of diminishing sea ice in recent Arctic temperature amplification, *Nature*, 464(7293), 1334–1337, doi:10.1038/nature09051, 2010.
- Screen, J. A. Climate science: far-flung effects of Arctic warming. *Nat. Geosci.* 10, 253–254 (2017).
- 670 Semtner, A.J.: A Model for the Thermodynamic Growth of Sea Ice I Numerical Investigations of

Serreze, M. C. and Barry, R. G.: Processes and impacts of Arctic amplification: A research synthesis, *Global and Planetary Change*, 77(1–2), 85–96, doi:10.1016/j.gloplacha.2011.03.004, 2011.

675

Serreze, M. C., Barrett, A. P., Stroeve, J. C., Kindig, D. N. and Holland, M. M.: The emergence of surface-based Arctic amplification, *The Cryosphere*, 9, 2009.

Seferian, Roland. CNRM-CERFACS CNRM-ESM2-1 model output prepared for CMIP6 CMIP amip. Version YYYYMMDD[1].Earth System Grid Federation. <https://doi.org/10.22033/ESGF/CMIP6.3924>, **2019**.

Shu, Q., Song, Z. and Qiao, F.: Assessment of sea ice simulations in the CMIP5 models, *The Cryosphere*, 9(1), 399–409, doi:10.5194/tc-9-399-2015, 2015.

685

SMITH, Doug M. et al. The Polar Amplification Model Intercomparison Project (PAMIP) contribution to CMIP6: Investigating the causes and consequences of polar amplification. *Geoscientific Model Development*, v. 12, n. 3, p. 1139-1164, 2019.

690 Stevens, B., Giorgetta, M., Esch, M., Mauritsen, T., Crueger, T., Rast, S., Salzmann, M., Schmidt, H., Bader, J., Block, K., Brokopf, R., Fast, I., Kinne, S., Kornblueh, L., Lohmann, U., Pincus, R., Reichler, T. and Roeckner, E.: Atmospheric component of the MPI-M Earth System Model: ECHAM6: ECHAM6, *J. Adv. Model. Earth Syst.*, 5(2), 146–172, doi:10.1002/jame.20015, 2013.

695 Stuecker, M. F., Bitz, C. M., Armour, K. C., Proistosescu, C., Kang, S. M., Xie, S.-P., Kim, D., McGregor, S., Zhang, W., Zhao, S., Cai, W., Dong, Y. and Jin, F.-F.: Polar amplification dominated by local forcing and feedbacks, *Nature Clim Change*, 8(12), 1076–1081, doi:10.1038/s41558-018-0339-y, 2018.

---

700 Stocker, T. F., Qin, D., Plattner, G. K., Tignor, M. M., Allen, S. K., Boschung, J., ... & Midgley, P. M.:  
Climate Change 2013: The physical science basis. Contribution of working group I to the fifth  
assessment report of IPCC the intergovernmental panel on climate change. Cambridge: Cambridge  
University Press, doi:  
10.1017/CBO9781107415324, 2014.

705

Sundqvist, H. S., Zhang, Q., Moberg, A., Holmgren, K., Körnich, H., Nilsson, J., and Brattström, G.:  
Climate change between the mid and late Holocene in northern high latitudes – Part 1: Survey of  
temperature and precipitation proxy data, *Clim. Past*, 6, 591–608, [https://doi.org/10.5194/cp-6-591-](https://doi.org/10.5194/cp-6-591-2010)  
2010, 2010.

710

Swart, Neil Cameron; Cole, Jason N.S.; Kharin, Viatcheslav V.; Lazare, Mike; Scinocca, John F.;  
Gillett, Nathan P.; Anstey, James; Arora, Vivek; Christian, James R.; Jiao, Yanjun; Lee, Warren G.;  
Majaess, Fouad; Saenko, Oleg A.; Seiler, Christian; Seinen, Clint; Shao, Andrew; Solheim, Larry; von  
Salzen, Knut; Yang, Duo; Winter, Barbara; Sigmund, Michael. *CCCma CanESM5 model output*  
715 *prepared for CMIP6 ScenarioMIP ssp126*. Version YYYYMMDD<sup>[1]</sup>. Earth System Grid  
Federation. <https://doi.org/10.22033/ESGF/CMIP6.3683>, 2019.

Swart, N. C. and Fyfe, J. C.: The influence of recent Antarctic ice sheet retreat on simulated sea ice area  
trends: Antarctic Sea ice trends, *Geophys. Res. Lett.*, 40(16), 4328–4332, doi:10.1002/grl.50820, 2013.

720

Tatebe, Hiroaki; Watanabe, Masahiro. *MIROC MIROC6 model output prepared for CMIP6 CMIP*  
*historical*. Version YYYYMMDD<sup>[1]</sup>. Earth System Grid  
Federation. <https://doi.org/10.22033/ESGF/CMIP6.5603> 2018.

725 Taylor, K. E., Stouffer, R. J. and Meehl, G. A.: An Overview of CMIP5 and the Experiment Design,  
*Bull. Amer. Meteor. Soc.*, 93(4), 485–498, doi:10.1175/BAMS-D-11-00094.1, 2012.

- 
- Thompson, D. W. J., Solomon, S., Kushner, P. J., England, M. H., Grise, K. M. and Karoly, D. J.: Signatures of the Antarctic ozone hole in Southern Hemisphere surface climate change, *Nature Geosci*, 730 4(11), 741–749, doi:10.1038/ngeo1296, 2011.
- Thompson, David WJ, and Susan Solomon.: Interpretation of recent Southern Hemisphere climate change, *Science* 296 (5569), 895-899, doi: 10.1126/science.1069270, 2002.
- Turner, J., Hosking, J. S., Bracegirdle, T. J., Marshall, G. J. and Phillips, T.: Recent changes in 735 Antarctic Sea Ice, *Phil. Trans. R. Soc. A*, 373(2045), 20140163, doi:10.1098/rsta.2014.0163, 2015.
- Turner, John et al. Unprecedented springtime retreat of Antarctic sea ice in 2016. *Geophysical Research Letters*, v. 44, n. 13, p. 6868-6875, 2017.
- 740 Van der Linden, E. C., Le Bars, D., Bintanja, R, and Hazeleger, W : Oceanic heat transport into the Arctic under high and low CO2 forcing. *Climate Dyn.*, <https://doi.org/10.1007/S00382-019-04824-Y>, 2019.
- Vaughan, D. G., Marshall, G. J., Connolley, W. M., Parkinson, C., Mulvaney, R., Hodgson, D. A., 745 King, J. C., Pudsey, C. J. and Turner, J.: Recent Rapid Regional Climate Warming on the Antarctic Peninsula, *Climate Change*, 60(3), 243-274, doi: 10.1023/A:1026021217991, 2013.
- Veiga, S. F., Nobre, P., Giarolla, E., Capistrano, V., Baptista Jr., M., Marquez, A. L., Figueroa, S. N., Bonatti, J. P., Kubota, P. and Nobre, C. A.: The Brazilian Earth System Model ocean–atmosphere 750 (BESM-OA) version 2.5: evaluation of its CMIP5 historical simulation, *Geosci. Model Dev.*, 12(4), 1613–1642, doi:10.5194/gmd-12-1613-2019, 2019.
- Volodin, Evgeny; Mortikov, Evgeny; Gritsun, Andrey; Lykossov, Vasily; Galin, Vener; Diansky, Nikolay; Gusev, Anatoly; Kostykin, Sergey; Iakovlev, Nikolay; Shestakova, Anna; Emelina,

---

755 Svetlana. *INM INM-CM4-8 model output prepared for CMIP6 CMIP*  
*piControl*. Version YYYYMMDD<sup>[1]</sup>. Earth System Grid  
Federation. <https://doi.org/10.22033/ESGF/CMIP6.5080> , 2019.

Walsh, J. E.: Intensified warming of the Arctic: Causes and impacts on middle latitudes, *Global and*  
760 *Planetary Change*, 117, 52–63, doi:10.1016/j.gloplacha.2014.03.003, 2014.

Walsh, J. E.: Intensified warming of the Arctic: Causes and impacts on middle latitudes, *Global and*  
*Planetary Change*, 117, 52–63, doi:10.1016/j.gloplacha.2014.03.003, 2014.

765 Watanabe, Masahiro et al. Improved climate simulation by MIROC5: Mean states, variability, and  
climate sensitivity. *Journal of Climate*, v. 23, n. 23, p. 6312-6335, 2010.

Watanabe, S., Hajima, T., Sudo, K., Nagashima, T., Takemura, T., Okajima, H., Nozawa, T., Kawase,  
H., Abe, M., Yokohata, T., Ise, T., Sato, H., Kato, E., Takata, K., Emori, S. and Kawamiya, M.:  
770 MIROC-ESM 2010: model description and basic results of CMIP5-20c3m experiments, *Geosci. Model*  
*Dev.*, 4(4), 845–872, doi:10.5194/gmd-4-845-2011, 2011.

Winton, Michael. A reformulated three-layer sea ice model. *Journal of atmospheric and oceanic*  
*technology*, v. 17, n. 4, p. 525-531, 2000.

775

Winton, M.: Amplified Arctic climate change: What does surface albedo feedback have to do with it?,  
*Geophys. Res. Lett.*, 33 (03701), doi:10.1029/2005GL025244, 2006.

Yang, X. Y., Fyfe, J. C., & Flato, G. M.: The role of poleward energy transport in Arctic temperature  
780 evolution. *Geophysical Research Letters*, 37(14). doi: 10.1029/2010GL042487, 2010.



---

785 Yukimoto, Seiji; Koshiro, Tsuyoshi; Kawai, Hideaki; Oshima, Naga; Yoshida, Kohei; Urakawa, Shogo;  
Tsujino, Hiroyuki; Deushi, Makoto; Tanaka, Taichu; Hosaka, Masahiro; Yoshimura, Hiromasa; Shindo,  
Eiki; Mizuta, Ryo; Ishii, Masayoshi; Obata, Atsushi; Adachi, Yukimasa. *MRI MRI-ESM2.0 model*  
output prepared for CMIP6 CMIP esm-hist. Version YYYYMMDD<sup>[1]</sup>. Earth System Grid  
Federation. <https://doi.org/10.22033/ESGF/CMIP6.6807>, 2019.

790 Yukimoto, Seiji et al. A new global climate model of the Meteorological Research Institute: MRI-  
CGCM3—Model description and basic performance—. *Journal of the Meteorological Society of Japan*.  
Ser. II, v. 90, p. 23-64, 2012.

Zhang J , Tian W , Chipperfield M P , et al. Persistent shift of the Arctic polar vortex towards the  
Eurasian continent in recent decades[J]. *Nature Climate Change*, 2016.

795 Ziehn, Tilo; Chamberlain, Matthew; Lenton, Andrew; Law, Rachel; Bodman, Roger; Dix, Martin;  
Wang, Yingping; Dobrohotoff, Peter; Srbinovsky, Jhan; Stevens, Lauren; Vohralik, Peter; Mackallah,  
Chloe; Sullivan, Arnold; O'Farrell, Siobhan; Druken, Kelsey :CSIRO ACCESS-ESM1.5 model output  
prepared for CMIP6 CMIP. Version YYYYMMDD[1]. Earth System Grid Federation, 2019.