#### Response to anonymous referee #1

The manuscript by Battarbee et al. has discussed the proton injection issue with results obtained by global hybrid-Vlasov and test-particle simulations. I think the quality of the paper is more than enough for publication. I have a few comments and suggestions that the author might want to address before the paper should be published.

We thank the referee for the review and improvement suggestions.

The word "non-locality" is a little bit confusing. I think it is more or less similar to the thickness of the shock (although not necessarily the same). It might be better to add some explanation for this as it is not in the standard terminology.

We agree that coming up with terminology for a new concept is challenging, and acknowledge how, in some respects, non-locality is similar to a shock thickness. We propose that a thickness is really only valid when the shock has a well-defined upstream and downstream and a clear transition between them, e.g. in the context of the quasi-perpendicular shock when the shock profile is clear and unambiguous (well localized). In the quasi-parallel region there are challenges associated with finding the shock profile, in particular as the shock reforms, as shown in Figure 2. We will add a comparison to shock thickness to the terminology subsection in the introduction.

According to the description of the simulation parameters, the spatial resolution (228km) is larger than the ion inertial length (125 km). It may not be so bad for modelling global phenomena, but one must be careful for doing accurate simulations of collision-less shocks. In particular, since the authors followed test particle trajectories on top of their simulation results to discuss the particle interaction with the shock, the resolution can be an issue. I guess that it is not easy to perform a convergence study for this particular application in a reasonable amount of computational resources. However, the authors may caution to the readers that there is potentially a numerical resolution issue.

We appreciate the reviewer's concern regarding the ion inertial length. A convergence test is indeed unrealistically expensive to perform. We do, however, intend to investigate this issue in the future.

We would also like to note that there exists a trade-off in simulations which focus on the small scales. For example, the mesoscale reformation features shown in our Figure 2 panel a) can have spatial extents of up to 2 RE or 100 di. These arise from the interaction of the curved bow shock with incident ULF wave fronts. With a given set of simulation resources, one needs to either run a local simulation, perform system re-scaling (e.g. Tóth et al 2017) which will negatively impact the global dynamics, or have a spatial resolution which does not resolve effects at or below ion inertial length scales. Our approach aims to investigate effects arising from the global scale whilst maintaining appropriate scale separation. We intend to elaborate this approach and our motivation in our manuscript.

We also note that the qualitative bow shock effects and reformation seen in these simulations are in agreement with other Vlasiator simulations (see the web site, <u>http://www.physics.helsinki.fi/vlasiator</u>) where the 30 degree IMF simulation cell size was set to the ion inertial length. We will investigate this run in the future, but wanted to utilize the quasi-parallel IMF for this initial study.

### The disagreement between the Vlasiator and test-particle results in table 1 may also arise from the same reason.

Since the test-particles and the Vlasiator distribution functions both are acted upon by fields of identical spatial resolution, we did not consider this a likely cause for the discrepancy. Test-particle fields are interpolated on the subgrid level in a linear fashion whereas the Vlasov distributions use volumetric-reconstructed fields.

The wave fields in the two runs are very different (Turc et al. 2018), resulting in differing trapping dynamics. We plan to elaborate this point of discussion accordingly.

#### It is no surprise to me that the non-locality is not an important factor to affect the injection as the ions have long interaction time with the shock and can travel for a long distance along the shock surface before being reflected or transmitted. The fate of the particles should be determined by the integral of electromagnetic fields as seen by them.

We agree that the electromagnetic fields are indeed the key to evaluating particle injection. As particle injection time scales are indeed significant, and close to reformation time scales, we felt it important to investigate a possible connection. There does not yet exist consensus in the field for injection and shock physics, but hopefully future studies will find convergence in our understanding of the quasi-parallel plasma shock.

#### Response to anonymous referee #2

We wish to thank the referee for the helpful review.

The manuscript analyses Vlasov numerical simulations including test particle runs of the quasi-parallel Earth's bow shock and ion acceleration there. The concept of "shock non-locality" is introduced and it is shown that the non-locality has little direct effect on particle injection. Instead the injection takes place in a larger region surrounding the shock but at the same time local magnetic field distortions at the shock are important for the injection.

#### Abstract promises a novel method for spacecraft data analysis. It is not clear what is the novel method, why and how it should be applied and what would be the outcome. This needs to be clarified.

Thank you, we will clarify that the proposed new analysis is calculating nonlocality from a combination of three plasma measurements.

#### Simulation initialisation: - Why 5∘tilt in the magnetic field?

In this study, we chose to focus on the quasi-parallel bow shock. The 5 degree run was chosen as it provided a large region where global curvature effects were captured within the quasi-parallel bow shock.

### - Is 43eV solar wind plasma temperature motivated by the velocity resolution?

This is correct. In order to ensure the shock dynamics are properly modelled, the incoming solar wind distribution must be adequately resolved by the velocity grid. This is verified by ensuring there is no numerical heating as the distribution propagates from the inflow boundary to the shock.

- There should be a proper discussion why simulating the system with a simulation having the spatial resolution larger than the characteristic ion inertial and ion gyroradius scales is appropriate to address the problem of ion injection where most of the ion reflection can occur on ion kinetic scales. This points needs to be clearly addressed as it may affect the general conclusions of the paper. We thank the referee for the feedback. We intend to improve this discussion in the paper with the following reasoning. Our simulations choose to emphasize the global dynamics due to, e.g. curved bow shock reformation as ULF waves impinge upon it. We acknowledge that there may be additional ion effects at smaller kinetic scales, pending further study and resource expenditure. A convergence study would be a very expensive undertaking, but we intend to further analyse these effects in future studies. To our knowledge, there doesn't exist a study yet which would invalidate the dynamics seen with this resolution.

#### It is not clear why authors have chosen to do the test particle approach if the Vlasov code is supposed to follow the full distribution function. This point needs to be clearly explained.

On line 210-213 we state: "Following the evolution of distribution functions does not allow for tracing of particle histories. In order to evaluate injection probabilities, particles need to be tracked as they meet the bow shock and interact with it, ultimately either returning to the upstream or being transmitted to the downstream." We will reword this for added clarity, highlighting how we actually use these test-particles as a method of tracking the evolution of a small portion of the VDF.

The concept of non-locality is introduced which does not include magnetic field. The motivation is that it provides poor results while at the same time paper mentions that magnetic field structures are very important for the injection. All this makes the motivation for the nonlocality concept very unsatisfactory and it is not clear what authors mean motivating their selection by having poor vs good results. Magnetic field data is one of the primary datasets in the shock analysis and it is unclear why one would want to exclude it from the shock definition. In general, it is not clear why authors want to introduce a new concept.

We acknowledge that magnetic field measurements are often used in spacecraft. We will clarify this section, stating that (in agreement with analytical studies of quasi-parallel shocks resulting in little magnetic field compression), the magnitude of magnetic field at the quasi-parallel shock showed multiple successive enhancements and rarefications. As our proposed method depended upon the measurement reaching a conclusive downstream state, the magnetic field magnitude as such was insufficient. We do, however note that the magnetic field does have a role in the calculation of the shock-normal magnetosonic Mach number, so the magnetic field is not ignored.

#### In Figure 4 100eV case of test particles is shown. It should be motivated why this particular case is shown and not for example the case of Maxwellian distributed particles.

The Maxwellian case appears very similar to the presented 100 eV case, so we shall replace the figure. We also note that the evolution of all performed test-particle distributions can be examined in supplementary movies B and C.

Figure 5 results and discussion are not fully consistent and should be significantly improved. For example, showing injected particle results (column 1 and 2) one makes conclusion that particles with energies below solar wind drift energy are loosing energy on average and particles above are gaining. This results is inconsistent with that the figures shows most of the injected particles have high energy. Such high energy particles if they start at solar wind energy and then during some part of the orbit have energies below the solar wind energy then on average at low energies the energy gain and loss should be equal. If the statement made in the manuscript is true then why there are no low energy injected particles (while there are still a lot of low energy particles at r<0 and they all show negative energy gain.

We wish to thank the referee for pointing out this error. We did additional analysis of our results, and found that when binning the particle energization changes, the script made an erroneous assumption that energy changes per time step would be very small. Due to each energy change being recorded at the end value, the plot emphasized energy gains at high energies and losses at low energies.

We reformulated the analysis to plot the initial energy on the y-axis, and we altered the energy calculations to be done in the incoming solar wind frame. This gave the additional benefit of being able to track the particle populations on the energization map as they approach the shock-associated region, and after the impact. We also implemented a more detailed method of instead of evaluating the position of particles as radial distance from the bow shock fit, we actually calculated the x-directional distance to the closest position where the shock fulfils the solar wind core heating criterion. There was some difference in the analysis of the results, but we can still posit that heating takes place over larger scales – low energy changes in the region in front of the shock, and large energy changes over an extended distance in the downstream.

Similarly, it is not clear how the current simulations results contradict the results from the Johlander et al. Firstly, it is not clear if SLAMS are observed in the current simulation and if they are do they have similar properties as in the observations? Secondly, when comparing with Johlander et al., it would be good to do the comparison in an adequate way, so that one understand how one should translate the results from the Vlasiator case to another cases such as Johlander et al. For comparison with those results one would need to look at solar wind ions that have different kinetic energy in the shock frame and see the differences in the injection rate. The authors should guide the reader where and how this can be seen.

We acknowledge that the comparison was very brief. In Johlander et al, Fig. 5, low-energy particles were found to be likely to reflect from SLAMS, whereas fast particles passed through them. In the quasi-parallel region, structures such as SLAMS merge into the bow shock, and thus, there is some merit in comparing how particles interact with SLAMS vs how they interact with the bow shock. Our results showed that particles with a large energy in the solar wind frame were more likely to be injected, but we acknowledge that a high solar wind frame speed can result in both faster and significantly slower shock-frame particles.

We revised figure 6 to replace the nose angle plot with shock-frame energies, which helps to showcase both how large solar wind frame energies result in a large spread of energies in the shock frame, and also how even at a given shock-frame energy, larger solar wind frame energies (i.e. larger tangential velocities) assist injection.

Figure 6 requires several clarifications. The largest structuring of the injection probabilities is seen in the dependance on the impact position angle. Instead of trying to resolve the physics of the large injection rate variations authors suggest how to smooth these variations which suggests that authors themselves maybe do not trust the numbers. This needs to be clarified.

We indeed expect that panel to not depict any underlying erratic dependence on shock-normal angle, but rather, to be indicative of how our particle injection time window was not long enough to encompass a sufficient amount of shock reformation cycles. Due to the large spatial extent and the large amount of test-particles, we still believe our statistics are sufficient to probe other properties of particle-shock-interactions.

In an ideal world, we would use a longer period of time for our test-particle study, but such a simulation set is not available at this time. However, in

the interest of clarity, we have removed this panel whilst adding a panel depicting shock-frame velocities of particles.

# Another unclear point is how shock non-locality is defined for a particle that starts at one position and gets injected at another position (in general valid for all particles). From which time and position are the given shock non-locality values. Similarly, it is not clear at which time instant is measured the bow-normal angle.

We acknowledge that the statement on lines 289-290: "For each testparticle, we evaluate these properties at the first time the particle reaches a point in the simulation space that fulfills the solar wind core heating (Tcore > 4Tsw ) criterion." was somewhat hidden. We shall clarify this point in the text and the caption.

#### *Minor things: L.8 fix the language of the sentence.*

Thank you for this correction.

## *I.45 The work of Johlander et al. 2016 does not make the mentioned assumptions in the manuscript but shows that SLAMS can contribute to the injection.*

The referee is correct, the assumptions of the test-particle study in Johlander et al (2016) differ in that they investigate a SLAMS instead of a planar shock front. We have clarified the introduction.

# *I.59 It is a bit confusing in which reference frame particle gains energy in the definition of the energization. For example, a particle reflected from a shock can have lower energy in the shock reference frame than the solar wind particles (e.g. when reflected from SLAMS) but it would not be "part of the incident thermal distribution".*

We thank the referee for making this important point, and we will add it to the manuscript.

#### *I.165 the division of the core distribution is unclearly described.*

We will reword this to the following:

"The Vlasiator distribution function is separated into core and suprathermal parts ( $n_{p,core}$  and  $n_{p,st}$ ). Each velocity space cell is evaluated as belonging to the core distribution, if it is inside a sphere centred at  $u_{sw} = (-600, 0, 0)$ 

km/s and with a radius of 690 km/s. Cells outside this sphere are considered as belonging to the suprathermal distribution."

#### *I.368 this should be illustrated and quantified, adding by the figure*

We will reword this for clarity – this in fact was referring exactly to Figure 5, which shows changes in particle energy as measured in the simulation frame. We will also amend the caption of Figure to this effect, and if necessary, change the text according to our re-done analysis.

#### I.411 What do you mean by "high-fidelity"?

We will replace "high-fidelity" with "noise-free".

## Figure 1: please use slightly thinner lines, the structure of the pink line cannot be resolved in the figure due to the thickness of the line.'

We did not intend Figure 1 to be used for evaluating the mesoscale bow shock shape, instead highlighting this in Figure 2. Nevertheless, we will redo Figure 1 with a smaller line width to show the details already in this image.

## Table 1: Why comparison is done with suprathermal densities? Iassume that from Vlasiator one can estimate the flux of reflectedparticles and thus have a good estimate of the injection rate.

Within the foreshock region, the suprathermal density indeed is a close measure of reflected particles, with only possible very minor contamination from the solar wind core during flow deflection events. The datasets of Vlasiator are too large (several terabytes) to store all data at every time step, thus we use reduced measurements such as the described split into core and suprathermal portions of the distribution function. We have added a description to explain that the suprathermal density is a good measure of reflected particles, adding the note that it includes all particles which have been reflected, even a long time ago, and are currently in the upstream.

## Non-locality of the Earth's quasi-parallel bow shock: injection of thermal protons in a hybrid-Vlasov simulation

Markus Battarbee<sup>1</sup>, Urs Ganse<sup>1</sup>, Yann Pfau-Kempf<sup>1</sup>, Lucile Turc<sup>1</sup>, Thiago Brito<sup>1</sup>, Maxime Grandin<sup>1</sup>, Tuomas Koskela<sup>1,2</sup>, and Minna Palmroth<sup>1,3</sup>

<sup>1</sup>Department of Physics, University of Helsinki, Helsinki, Finland

<sup>2</sup>Department of Physics and Astronomy, University of Turku, Turku, Finland

<sup>3</sup>Finnish Meteorological Institute, Helsinki, Finland

Correspondence: Markus Battarbee (markus.battarbee@helsinki.fi)

**Abstract.** We study the interaction of solar wind protons with the Earth's quasi-parallel bow shock using a hybrid-Vlasov simulation. We employ the high-fidelity- global hybrid model Vlasiator to include effects due to bow shock curvature, tenuous upstream populations, and foreshock waves. We investigate the local uncertainty of the position of the quasi-parallel bow shock as a function of several plasma properties, and find that for a significant portion of time, the local bow shock position

5 is challenging to define. Our results support the notion of upstream structures causing patchwork reconstruction of the quasiparallel shock front in a non-uniform manner. We propose a novel method for spacecraft data to be used to analyze this quasi-parallel reformation.

We combine our hybrid-Vlasov results with test-particle studies and show that shock non-locality appears to have only limited effect little direct efficient on particle injection. We show that proton energization, which is required for injection, takes

10 place throughout a larger shock transition zone. Non-local energization of particles is found regardless of the instantaneous non-locality of the shock front. Distortion of magnetic fields in front of and at the shock is shown to have a significant effect on proton injection.

We additionally show that the density of suprathermal reflected particles upstream of the shock may not be a useful metric for the probability of injection at the shock, as foreshock dynamics and particle trapping appear to have a greater significant

15 effect on energetic particle accumulation at a given position in space. Our results have implications for statistical and spacecraft studies of the shock injection problem.

Copyright statement. TEXT

#### 1 Introduction

Collisionless plasma shocks are an ubiquitous source of plasma acceleration, common within stellar, planetary, and interplan-20 etary environments. They-Shock dynamics have been studied in great detail at the Earth's bow shock. In regions of shock geometry where the angle  $\theta_{Bn}$  between the shock-normal direction  $\hat{n}$  and the upstream interplanetary magnetic field (IMF) direction **B** is small ( $\leq 45^{\circ}$ ), the shock is considered quasi-parallel (see, e.g., Burgess et al., 2005). In this region, if the shock is a strong fast-mode supercritical shock, a fraction of thermal incident ions are reflected, streaming away from the shock along the magnetic field lines, forming the foreshock region (Fairfield, 1969; Eastwood et al., 2005). The streaming energized

- 25 particles excite instabilities such as a right-hand ion-ion beam instability, building a wave field of ultra-low frequency (ULF) waves (Hoppe et al., 1981) with periods around ~30s, which further interact with the particles themselves and are convected toward the bow shock. As the waves are convected with the supersonic solar wind flow, they appear mostly left-handed in the spacecraft frame. The incident ULF waves can experience nonlinear steepening, possibly forming shocklets (Hada et al., 1987; Wilson III, 2016) or short large amplitude magnetic structures (SLAMS; Schwartz et al., 1992; Burgess, 1995; Lucek et al.,
- 30 2008), eventually causing patchwork reformation of the bow shock (Scholer and Terasawa, 1990; Thomas and Winske, 1990; Schwartz and Burgess, 1991; Burgess, 1995) as incoming structures proceed to build a new shock front periodically (Burgess, 1989). The complicated structure of the shock-associated transition region was linked with local reconnection in Gingell et al. (2019). As the location of the shock front is challenging to define due to movement i.e. nonstationarity of a well-defined shock front, the formation and convection of a new shock front, and even the disappearance of the old front, we now discuss this
- 35 uncertainty of the shock position which we designate the "non-locality" of the shock. As plasma parameters across a quasiparallel shock can be non-monotonic, non-locality encompasses more than mere thickness of a well-defined shock front. Our definition of non-locality can also be measured using spacecraft, providing a novel metric for quantifying space plasma observations. In this study, we limit our analysis to ion scales and assume the reformation of the quasi-parallel bow shock to happen on temporal and spatial scales similar to those of steepened ULF waves and associated transient structures.
- 40 An important open question for space physics and particle acceleration is the shock injection problem (see, e.g., Zank et al., 2001), or how exactly thermal particles are reflected at a super-critical quasi-parallel shock. Injection from a thermal population is a necessary step in efficient diffusive shock acceleration (DSA; Axford et al., 1977; Blandford and Ostriker, 1978; Bell, 1978; Krymsky et al., 1979), which is a major source of energetic particles throughout the universe. The injection problem has been studied extensively during the past decades with, amongst others, observations (Sckopke et al., 1983; Thomsen et al., 1983;
- 45 Gosling et al., 1989; Johlander et al., 2016), analytical work (Schwartz et al., 1983; Malkov et al., 2016), test-particle modeling (Gedalin, 2001; Battarbee et al., 2011; Gedalin, 2016; Johlander et al., 2016), and particle-in-cell simulations (Caprioli et al., 2015; Liseykina et al., 2015; Sundberg et al., 2016; Hao et al., 2016; Caprioli et al., 2017). Significant historical work using 1-D or 2-D local hybrid simulations can be found in, e.g., Burgess (1989); Scholer (1990) and Kucharek and Scholer (1991). Previous studies have suggested three methods for injection: Specular reflection (Gosling et al., 1982), shock drift acceleration
- 50 (SDA; Giacalone, 1992; Lever et al., 2001; Burgess, 1987) and associated shock surfing (Lever et al., 2001), and thermal leakage from the downstream (Ellison, 1981; Edmiston et al., 1982; Lyu and Kan, 1990; Malkov, 1998). All these- These three methods were derived from assumptions of macroscopic, planar, and stationary shock fronts and are thus limited, but an important first step towards understanding the concept. Magnetic mirroring as described through quasi-linear theory and conservation of the first adiabatic invariant is usually excluded, as changes to magnetic fields may occur on scales much smaller
- 55 and faster than those of ion gyromotion.

In this paper, we investigate the complex structure and non-locality of the Earth's quasi-parallel bow shock as well as the injection problem both through hybrid-Vlasov simulations and test-particle runs. In section 2 we present our hybrid-Vlasov simulations. In section 3 we present results from two different hybrid-Vlasov datasets. In section 4 we introduce our test-particle simulation method, and in section 5 we present results of test-particle injection and energization. Section 6 presents analysis and discussion on our findings, and we present our conclusions in section 7.

60

65

80

Throughout this study, we use the following terminology:

- An *injected* particle has interacted with the bow shock and returned to the upstream. This may also be called *reflection*.
   During this process, particles are energized gain energy in the solar wind frame.
- A *transmitted* particle has passed through the bow shock to the far downstream. The particle may or may not be energized during this process.
- *Energization* is when during a single shock encounter, a particle gains energy in the solar wind frame so that it is no longer part of the incident plasma thermal distribution.
- Acceleration is when injected particles continue to gain energy through continuous and/or repeated shock interactions, such as DSA. This takes place over longer temporal and spatial scales, and is outside the scope of this study.
- *Non-locality* of the quasi-parallel bow shock is a measure of the disagreement between different measurements of where the bow shock is locally estimated to be. This could also be referred to as the uncertainty of the shock position.
  - The *shock-normal direction*  $\hat{\mathbf{n}}$  is normal to the local, reforming shock front. This direction is highly variable.
  - The *bow-normal direction*  $\hat{\mathbf{n}}'$  is the normal direction for a parabola, estimating the global shape of the shock front. This direction is very stable.
- The *shock-normal angle*  $\theta_{Bn}$  is the angle between the upstream magnetic field and the shock-normal direction. The shock-normal direction or a vector antiparallel to it is chosen in order to constrain the value to  $\theta_{Bn} \in [0^\circ, 90^\circ]$ . Due to fluctuations of both the upstream field and the local shock front, this angle is very unpredictable.
  - The *bow-normal angle*  $\theta_{Bn'}$  is the angle between the upstream magnetic field and the bow-normal direction. Like  $\theta_{Bn}$ , it is usually limited to  $\theta_{Bn'} \in [0^\circ, 90^\circ]$ , but in regions of significant mangetic field deformation, is allowed to have values
  - $> 90^{\circ}$ . This measure allows analysis of shock interaction due to upstream magnetic field fluctuations while smoothing out the local reformation effects of the quasi-parallel shock front.

#### 2 Vlasiator simulation

In modeling the Earth's bow shock, we employ Vlasiator (von Alfthan et al., 2014; Pfau-Kempf, 2016; Palmroth et al., 2018), a hybrid-Vlasov code designed to simulate the Earth's magnetosphere and the surrounding space environment. Vlasiator mod-

els kinetic proton-scale plasma physics by calculating the evolution of the proton distribution function on a Cartesian 3-dimensional velocity grid within each cell of a Cartesian spatial grid. In the presented runs, the spatial simulation domain is 2-dimensional. Modeling distribution functions directly instead of using a particle-in-cell method allows for accurate analysis of even the tenuous portions of non-thermal populations in the foreshock, and gives us a realistic model of foreshock and bow shock evolution. The noise-free distribution function formalism further allows using the magnetic field B and electric field E
values as input to test-particle studies without a need for low-pass filtering.

Vlasiator models ions as distribution functions, solving the Vlasov equation for the ion (proton) distribution with electrons modeled as a cold massless charge-neutralizing fluid. Closure is provided via Ohm's law, including the Hall term. We assume that effects due to the electron pressure gradient can be neglected. Vlasiator is capable of modeling a number of ion kinetic effects even without resolving ion kinetic scales spatially the ion skin depth (Pfau-Kempf et al., 2018), and has been used for a number of several interesting foreshock and bow shock studies (Palmroth et al., 2015; Pfau-Kempf et al., 2016; Turc et al.,

2018; Blanco-Cano et al., 2018; Turc et al., 2019). Our choice of simulation parameters do not quite resolve the ion inertial length, but instead ensure correct scale separation between global and local dynamics and a noise-free representation of both thermal and non-thermal plasma. Tóth et al. (2017) have investigated how reconnection physics were affected by overresolving the inertial length (at the expense of scale separation), but they did not study the consequences of underresolving it.

95

- In this paper, we use two datasets (simulations S1 and S2) modeling two different bow shock strengths and interplanetary magnetic field intensities. Results from these simulations have previously been published in Palmroth et al. (2015), and Turc et al. (2018), and Turc et al. (2019). They are ecliptic plane (x - y) 2D–3V simulations (2D in the spatial domain, 3D in the velocity domain) parametrized using the Geocentric Solar Ecliptic (GSE) coordinate system with no tilt for the Earth's dipole. The *x*-coordinate is along the Earth-Sun axis, the *z*-coordinate is aligned with the Earth's magnetic axis, and the *y*-
- 105 coordinate completes the right-handed system. We save variables such as field values and distribution function moments every 0.5 s. The simulation extent is  $2000 \times 1750$  spatial cells, covering the ranges  $x \in [-7.7, 63.6] r_E$  and  $y \in [-31.3, 31.3] r_E$  where  $r_E = 6371$  km is the Earth radius. The simulation domain extent in the z-direction is only one cell thick with periodic boundary conditions. Each spatial cell is a cube of length 228 km along each edge. Our velocity domain employs a sparse representation (von Alfthan et al., 2014) and has a resolution of 30 km s<sup>-1</sup>. The simulation domain is initialized with a somewhat fast and hot
- 110 solar wind inflow of  $n_{p,sw} = 3.3 \times 10^6 \text{ m}^{-3}$ , T = 0.5 MK,  $\mathbf{V} = (-600, 0, 0) \text{ km/s}$ , and either. The magnetic field in simulation S1 is  $\mathbf{B}(S1) = (-5\cos 5^\circ, 5\sin 5^\circ, 0) \text{ nT}$  or whereas in simulation S2 it is  $\mathbf{B}(S2) = 2\mathbf{B}(S1) = (-10\cos 5^\circ, 10\sin 5^\circ, 0) \text{ nT}$ . The quasi-radial IMF in these runs allows us to focus on the quasi-parallel bow shock. The somewhat hot solar wind ensures the inflow Maxwellian distribution is resolved adequately. The Earth's magnetic dipole is implemented at a realistic value of  $8.0 \times 10^{22} \text{ Am}^2$ , and the simulation domain inner boundary is a perfectly conducting sphere located at r = 31800 km or
- 115 about  $5r_{\rm E}$ . The simulation set-up results in solar wind Alfvénic Mach numbers of  $M_{\rm A,1} \sim 10$  and  $M_{\rm A,2} \sim 5$  and magnetosonic Mach numbers of  $M_{\rm ms,1} \sim 5.4$  and  $M_{\rm ms,2} \sim 3.8$  in front of the bow shock nose, and thus, strong fast-mode supercritical shocks. The simulations were run for  $t_{\rm max,1} = 685$  s and  $t_{\rm max,2} = 539$  s, respectively. To facilitate comparison with existing numerical studies, we note that for both simulation runs the solar wind ion inertial length is 125.4 km =  $0.020 r_{\rm E}$ , and for S1, the solar wind plasma beta  $\beta_1 = 2.3$ , and for S2,  $\beta_2 = 0.57$



Figure 1. Overview of the Vlasiator simulation S1 ( $B_{IMF} = 5 \text{ nT}$ ,  $M_A = 10$ ) at time t = 500 s, with proton number density (colormap) overlaid with an estimate of the bow shock position according to plasma compression (thick fuchsia curve,  $n_p > 2n_{p,sw}$ ). Also shown are magnetic field lines (thin black curves) and two white overlapping rectangles indicating zoom-in regions used for analysis of local bow shock structure (smaller rectangle) and test-particle studies (larger rectangle).

- Figure 1 depicts the Vlasiator simulation domain for simulation S1. The color map depicts proton densities, showing a dense magnetosheath between the bow shock and the magnetosphere, as well as variations in the upstream plasma density within the proton foreshock region. A thick fuchsia contour depicts where plasma density has increased two-fold over solar wind values, providing a rough estimate of the bow shock position. Thin b Black lines illustrate magnetic field lines, showing how the foreshock is permeated by fluctuations, as well as visualizing the complicated nature of magnetic flux compression and deflection at the quasi-parallel bow shock. The white circle indicates the simulation inner boundary, and two overlapping white
- rectangles indicate our regions of interest within the simulation. The larger white rectangle is used for visualizing test-particle studies of proton injection, whereas the smaller rectangle is used for analysis of quasi-parallel bow shock non-locality.

#### 3 Vlasiator results

In this section, we present results of hybrid-Vlasov simulations. First, we fit the global position of the bow shock using a

130 quartic estimation and calculate the bow-normal angle to estimate the general direction of the shock normal. As our fit is so close to a parabola, we will henceforth for simplicity refer to it as a parabola. Then, we use several local measurements of plasma properties to estimate the rapidly moving and varying local position of the shock, and use their disagreement to define a non-locality of the shock.

#### 3.1 Bow shock location and the shock-normal angle

- 135 In previous hybrid-method investigations into ion injection at kinetic plasma shocks, the shock descriptions have been usually either 1-D (see, e.g., Lyu and Kan, 1990; Scholer, 1990; Scholer and Terasawa, 1990; Onsager et al., 1991; Su et al., 2012) or if 2-D or 3-D, limited to local geometries (Guo and Giacalone, 2013; Caprioli et al., 2015; Hao et al., 2016; Sundberg et al., 2016; Caprioli et al., 2017). In a local planar shock, it is feasible to simply define the shock-normal direction from simulation box parameters and evaluate 1-D cuts along this line for defining the shock shape. However, as seen in Figure 1,
- 140 in a global 2-D simulation, the curved bow shock has a bow-normal direction dependent on the nose angle  $\phi = \arctan(y/x)$ , which complicates evaluating the shock-normal direction (Thomas and Winske, 1990). Shock and injection investigations within global simulations have recently been published in, e.g., Savoini et al. (2010, 2013); Karimabadi et al. (2014); Savoini and Lembège (2015).

We now determine a rough estimate of the global bow shock shape. We do this by finding the contour where plasma density
increases two-fold over the solar wind value (n<sub>p</sub> > 2n<sub>p,sw</sub>). The value of 2n<sub>p,sw</sub> was chosen based on visual inspection. We then fit a 4th order polynomial

$$r_s(\phi) = a_0 + a_1\phi + a_2\phi^2 + a_3\phi^3 + a_4\phi^4 \tag{1}$$

using the nose angle and the radial distance  $r = \sqrt{y^2 + x^2}$  at each contour position. This fit is performed at times  $t_0 = 438$  s and  $t_f = 538$  s. We found that intermediate time steps are described well by performing linear interpolation in time of the polynomial coefficients.

One of the most commonly used criteria for defining the dynamics and injection characteristics of a shock is the shocknormal angle  $\theta_{Bn}$ , i.e., the angle between the shock-normal direction and the upstream magnetic field. The upstream magnetic field direction in the quasi-parallel shock region varies greatly due to upstream fluctuations (Greenstadt and Mellott, 1985). Thus, even within the quasi-parallel regime, the shock may exhibit a wide variety of shock-normal angles.

155

150

As the shock front evolves, reforms, and fluctuates, the local shock-normal direction also evolves. The local instantaneous shock-normal direction can end up being perpendicular or even reversed to the mean bow shock direction, and is thus challenging to evaluate in a meaningful manner. In this study, we define an alternative measure, the *bow-normal direction*  $\hat{\mathbf{n}}'$ , which is the normal direction for the parabolic fit to the mean shape of the global shape of the shock front. This is calculated as

$$\mathbf{n}' = \left(-\frac{\mathrm{d}r(\phi)}{\mathrm{d}\phi}\cos\phi + r(\phi)\sin\phi, \frac{\mathrm{d}r(\phi)}{\mathrm{d}\phi}\sin\phi + r(\phi)\cos\phi, 0\right) \tag{2}$$

and accordingly  $\hat{\mathbf{n}'} = \mathbf{n'}/n'$ . We use this bow-normal direction both for defining the bow-normal plasma bulk velocity component, used for calculating the magnetosonic Mach number of the shock, and for defining a bow-normal angle  $\theta_{Bn'}$ , describing the angle between the local wave-distorted magnetic field and the bow-normal direction.

#### 3.2 Shock non-locality

- The locations of quasi-perpendicular and subcritical collisionless plasma shocks can, for the most part, be estimated well due to the upstream remaining undisturbed. However, at supercritical quasi-parallel shocks, the upstream is characterized by magnetic and density fluctuations and an abundance of suprathermal particles. This can make defining the exact position of the quasi-parallel shock challenging. This localization is further hindered by the fact that the position of the shock changes locally at timescales related to shock reformation. Additionally, the global position of the shock changes at larger timescales due to variation in solar wind driving conditions. This non-stationarity of the shock is observed as, e.g., spacecraft encountering the shock multiple times during what is expected to be a single crossing (see, e.g., Lucek et al., 2002; Sundberg et al., 2016;
- Gingell et al., 2017). In order to investigate the injection problem, we now attempt to define the local quasi-parallel shock position within a larger shock transition zone (Burgess, 1995) on reformation-related timescales. We also present a novel method for quantifying the difficulty of defining-uncertainty of the shock position, suitable for use in spacecraft observations.

We evaluate the location of the shock as a transition between the upstream and downstream conditions using three plasma

- 175 properties. The first is plasma compression, using the previously introduced criterion of  $n_p > 2n_{p,sw}$ . The second is heating of the solar wind core population,  $T_{core} > 4T_{sw}$ , similar to the method of Wilson III et al. (2014b, a), with the value  $4T_{sw}$  selected based on visual inspection. The Vlasiator distribution function is split into core and suprathermal parts ( $n_{p,core}$  and  $n_{p,st}$ ). by evaluating whether the solar wind frame ( $u_{sw,x} = -600 \text{ km/s}$ ) velocity of particles is above or below  $v_{core,max} = 690 \text{ km/s}$ . The plasma contained in each velocity space cell is evaluated as belonging to the core distribution if it is inside a sphere
- 180 centered at  $u_{sw} = (-600, 0, 0) \,\mathrm{km \, s^{-1}}$  and with a radius of  $690 \,\mathrm{km \, s^{-1}}$ . Cells outside this sphere are considered as belonging to the suprathermal distribution. The third criterion is when the plasma magnetosonic Mach number, calculated using the local fast magnetosonic mode speed and the bow-normal plasma bulk velocity, falls below 1. We do not include any further criteria based on the magnetic field direction or magnitude, as they were found to provide poor results at the quasi-parallel bow shock.magnetic field compression at a quasi-parallel shock is sporadic and limited, and the transition region has a wide range of local
- 185 field orientations (see, e.g., Figure 1 of Gingell et al., 2019). We emphasize that the presented methods will potentially register shocklets and SLAMS as they take part in the reformation process.

In Figure 2 we present in panels (a) and (b) snapshots of plasma density from simulations S1 and S2, respectively, at time t = 500 s, zoomed in on the nose of the quasi-parallel bow shock (indicated by the smaller white rectangle in Figure 1). We have plotted the plasma density with overlaid contours representing the bow shock positions according to criteria for plasma density (fuchsia), solar wind core heating (green), and magnetosonic Mach number (pale blue).

190

The three contours are highly variable and in agreement with agree on the position of the quasi-parallel shock only on the order of 50% of the time. We have selected four positions for profile cuts, depicted by black dashed lines in panel (a), showcasing different kinds of shock crossings. These simulate what a spacecraft might observe, except that they are spatial instead of temporal profiles. Line profiles for the three plasma properties used to gauge the shock position are shown in panels

(c), (d), (e), and (f). Graphed quantities are scaled so that a value of 1 is where the shock is estimated to be. The distance between the positions of bow shock parametrization closest and farthest from the Earth can be considered the disagreement between the three parametrizations, and is shown as shaded gray regions. This distance estimates the uncertainty of the shock position, or the extent of the shock transition region within which the three plasma properties estimate the shock to be. We define this distance as the shock *non-locality*. It is defined in units of Earth radii instead of, e.g., upstream gyroscales in order to facilitate comparison of bow shock structure sizes between different IMF conditions.

The cut shown in panel (c), at  $Y = 3.8 r_E$ , shows regions of low plasma density in what would appear to be the downstream, likely a result of a new shock front forming at  $X \approx 11 r_E$ , with the old shock position closer to  $X \approx 10.5 r_E$ . Panel (d), at  $Y = 2.8 r_E$ , shows active reformation of the quasi-parallel bow shock, with the first and last estimated shock positions disagreeing by over  $1.0 r_E$ , as a new front is forming at  $X \approx 11.7 r_E$ . The cut in panel (e), at  $Y = 1.2 r_E$ , is an example of a well-defined shock front where all criteria agree, and panel (f) shows an intermediate case where the three criteria disagree somewhat and the shock transition seems to extend radially over a distance of several hundred kilometers. An animation

Quantifying the non-locality of the quasi-parallel bow shock using spacecraft data will be more challenging than for simulations. Simulations allow us to directly measure spatial scales, whereas spacecraft motion in relation to quasi-parallel refor-210 mation is slow, and thus, use of constellation spacecraft and multipoint techniques are usually needed in order to infer spatial scales. At Mercury this reformation has been studied through mainly magnetic field measurements in Sundberg et al. (2013).

depicting time evolution of Figure 2 is available as Supplementary Video A.

We now describe how we evaluate the non-locality of the quasi-parallel bow shock in Vlasiator simulations. At one degree nose angle intervals, we draw a profile across the shock in the bow-normal direction, and measure where along the profile each of our three shock criteria (plasma density  $n_{\rm p} = 2n_{\rm p,sw}$ , solar wind core heating  $T_{\rm core} = 4T_{\rm sw}$ , and magnetosonic Mach

- 215 number  $M_{\rm ms} = 1$ ) indicate the local position of the shock is. Then, for each profile, we calculate the distance between the positions of bow shock parametrization closest and farthest from the Earth. This distance estimates the extent of the shock transition region, i.e., the non-locality of the shock. In Figure 3, panels (a) and (b), we plot stacked profiles displaying the temporal evolution of shock non-locality for simulations S1 and S2, respectively. Regions of enhanced shock non-locality appear to move along the shock front away from the nose region (indicated with a dashed line), as shown by the diagonal
- 220 ridges. S1 shows significantly larger and clearer non-locality structures than S2. Still, there exists a qualitative similarity to the structures seen for both simulations. We note that the motion of structures away from the nose might be due to either deflected plasma flow carrying structures along the front, or due to foreshock wave fronts convecting in and interacting with a curved bow shock at increasing nose angle positions. In panels (c) and (d), we show logarithmic histograms of accumulated shock non-locality measurements, showing that a well-defined shock is the most common occurrence, and increasing enhanced values
- of non-locality is-are increasingly rare. This also confirms that S2 has, on average, lower measurements of shock non-locality than S1 does.



Figure 2. Proton number density overlaid with bow shock positions according to criteria for plasma density (fuchsia,  $n_p = 2n_{p,sw}$ ), solar wind core heating (green,  $T_{core} = 4T_{sw}$ ), and magnetosonic Mach number (pale blue,  $M_{ms} = 1$ ). Panel (a) is for S1 ( $B_{sw} = 5 \text{ nT}$ ), panel (b) for S2 ( $B_{sw} = 10 \text{ nT}$ ), both at t = 500 s. Panels (c–f) show line profiles of the three bow shock criteria along the dashed black lines shown in panel (a), corresponding with differing amounts of shock non-locality.



**Figure 3.** Profiles of measured shock non-locality as a function of nose angle. The *y*-axis lists simulation time, used as the base level corresponding with a well-defined shock with a non-locality measure of zero. Regions of enhanced non-locality are shown as colored peaks of the curve, as presented in the color bar. A dashed vertical line indicates nose angle  $0^{\circ}$ . (a): S1. (b): S2. Both plots show chains of enhanced non-locality regions, which move away from the nose region and decrease in intensity as they approach the flanks. Under each stacked profile plot we show a histogram depicting the occurrence rate of different non-locality levels, with panel (c) depicting S1 and panel (d) depicting S2.

#### 4 Test-particle simulations

The Vlasiator model tracks the evolution of distribution functions as volume averages on a Cartesian mesh. Thus, particle trajectories are not a direct output of the code, and tracing particle histories requires the use of a post-processing tracer.

- Following the evolution of distribution functions does not allow for tracing of particle histories. In order to evaluate injection probabilities, particles need to be tracked as they meet the bow shock and interact with it, ultimately either returning to the upstream or being transmitted to the downstream. Thus, we chose to use a test-particle method to track the motion of single protons within the evolving, locally interpolated electric and magnetic fields output from the Vlasiator simulation. The particle propagation uses a Boris-push algorithm (Boris, 1970) with a conservative time step of  $\Delta t = 0.005$  s. This time step is not limited by particle gyrotimes, but rather, ensures that particles up to  $10^5$  eV travel less than  $1/10^{\text{th}}$  of a simulation cell per time
- step.  $\mathbf{E}$  and  $\mathbf{B}$  field values for each particle step are acquired from the Vlasiator output files using linear interpolation in both time and space. Thus, the test-particles act as tracers for an infinitesimal element of the distribution function.

Our goal is to use test-particle simulations to investigate proton injection at the quasi-parallel bow shock. For this purpose, we initialize our particles from the thermal solar wind core population, evenly distributed along a smooth curve a short distance

- 240 in front of the bow shock. We follow the particles as they approach the shock region and interact with it. If a particle reaches again a boundary well in front of the shock, it is considered injected, and if it passes far into the downstream, it is considered transmitted. Once a particle has been flagged as injected or transmitted, it is no longer propagated. A significant portion of test-particles spend so much time within the shock structure that they are not flagged as either injected or transmitted at the end of the run, and their fate remains inconclusive.
- The particle initialization curve is placed  $0.9r_{\rm E}$  outward of the parabolic bow shock fit, extending between nose angles  $\pm 40^{\circ}$ . This is visible in panel (a) of Figure 4 as the location of the first test-particles. An injection flagging boundary is placed  $0.1r_{\rm E}$  beyond the injection curve, and a transmission flagging boundary is placed  $1.5r_{\rm E}$  inward of the parabolic bow shock fit. These values were chosen so that the majority of changes to local quasi-parallel bow shock structure due to reformation fall within this region. We specifically note that the solar wind core heating criterion triggers always within this region.
- Each test-particle run consists of  $N = 10^5$  protons, initially isotropic in the frame co-moving with the inflow plasma, which results in a mean simulation frame energy of 1.9 keV. For each test run, particle velocities were chosen so they were as monoenergetic (10, 20, 50, 100, 200, or 500 eV) in the inflow plasma frame and randomly distributed in direction. Additionally a Maxwellian test run was performed, with particles picked randomly from a Maxwellian 0.5 MK distribution centered in the inflow plasma frame. Particles were placed into the simulation as groups of 25000 particles every 0.5 s for 10 seconds, starting at  $t_0 = 438$  s. Particle propagation was halted at time  $t_f = 538$  s.

#### 5 Test-particle results

In Figure 4, we display snapshots of test-particle propagation for simulation S1 and a plasma frame initialization energy of 100 eV. an initially maxwellian distribution of 0.5 MK in the solar wind frame. The grayscale region shows a logarithmic test-particle density, with black indicating single particles and white indicating over 100 particles per cell. We display contours

260 parametrizing the shock position on top, and also plot two black parabolas which act as the injection and transmission flagging boundaries. Animations depicting the evolution of test-particle populations for all initialization parameters and simulations S1 and S2 are available in Supplementary Videos B and C, respectively.

The panels in Figure 4 show how solar wind protons start as an even curve (a), are launched into the simulation over 10 seconds, after which the first ones have already accumulated as white regions at the shock front (b). We note how the steepened

- structure at  $Y \approx 2r_E$  in panel (b) causes an accumulation of test-particles at its -Y edge, and that the regions of plasma depletion (fuchsia contour at, e.g.,  $Y \approx 6r_E$ ,  $Y \approx 2r_E$ , and  $Y \approx -3r_E$ ) remain void of test-particles at this time. By the time of panel (c), all test-particles have reached the shock transition region, the white regions of test-particle accumulation follow shock ripples, and many of the previously void regions have been filled with <del>plasmatest-particles</del>. In panel (d) we see regions of efficient reflection causing particles to be returned to the upstream direction, but several regions also allow particles to move
- 270 past the shock front-and form into-, reminiscent of magnetosheath jets (Němeček et al., 1998; Hietala et al., 2009; Palmroth et al., 2018). By the time of panel (e), particles have spread to most of the magnetosheath all the way to the transmission boundary. Panel (f) displays how both transmission and injection can be slow processes, with 20–40% of particles still within the simulation after 90 100s of test-particle propagation, both in the upstream and in the downstream of the shock. For these particles, their ultimate fate of being injected or transmitted could not be evaluated from these simulations. Judging from panel 275 (f) of Figure 4, a portion of these particles would likely be injected.

Evaluation of test-particle interactions with the shock structure as seen in Figure 4 did not provide a clear answer as to where within the shock transition region particles truly feel the impact of the shock. As a particle injected into the upstream necessarily will experience energization in the solar wind frame, we tracked the simulation- solar wind frame energies of transmitted and injected particles and measured the regions where particles gained or lost the most energy. In Figure 5 we plot 2D-histograms of mean particle energy rate of change  $\langle \Delta E / \Delta t \rangle$ , which was calculated by gathering all particle energy changes over the whole test-particle simulation and normalizing the result with the amount of test-particles measured at each position in parameter space. As energy gains and losses can be significant near strong electric fields (up to 1 keV per measurement interval), we use the initial energy of each change as the y-coordinate. The black contours depict logarithmic counts of measurements, starting

- from a single particle with the thin dotted line. The colormap of mean energization is only plotted where a minimum of 100 measurements were counted. We note that the energization colormap is a symmetric logarithmic plot, with a small linear region between  $\pm 10 \,\mathrm{eV \, s^{-1}}$ . The presented initialization energies of 10 and 100 eV correspond to  $44 \,\mathrm{km \, s^{-1}}$  and  $138 \,\mathrm{km \, s^{-1}}$ plasma frame velocities, respectively. We show energization plots for only those particles which were registered as transmitted or injected by the end of the test-particle simulation. A grey band indicates the mean energy of incoming solar wind particles in the simulation frame solar wind ram energy, which is the minimum energy required for a particle to travel
- 290

sunwards, and thus the minimum energy for injection  $(1.9 \text{ keV for } 600 \text{ km s}^{-1})$ . In the first two rows of Figure 5, the x-axis shows the distance  $\Delta x$  from the closest position where the solar wind core heating shock criterion is met. The last two rows plot the instantaneous shock non-locality for the measurement, extracted from the nose angle bins calculated in Section 3.2.

The top half of Figure 5 clearly shows how particles start at the bottom right corner of each panel at initialization energies and upstream of the shock, and how they on average gain energy as they approach the shock. In the downstream, energization



Figure 4. Test-particle propagation for simulation S1, monoenergetic 100 eV maxwellian 0.5 MK initialization at 6 different times. Vlasiator simulation proton number density is overlaid with the logarithmic density of test-particles in greyscale, with white indicating over 100 particles in a cell. Two black parabolas are the transmission boundary (left) and the injection boundary (right). Three contours indicate estimates of the local shock position: plasma compression (fuchsia,  $n_{\rm p} > 2n_{\rm p,sw}$ ), solar wind core heating (green,  $T_{\rm core} > 4T_{\rm sw}$ ), and the magnetosonic Mach number (pale blue,  $M_{\rm ms} < 1$ 



**Figure 5.** Mean energization experienced by test-particles over their shock interaction. Energization tracking is performed separately for injected (columns 1 and 2) and transmitted (columns 3 and 4) particles. The top two rows track energization as a function of current particle simulation – solar wind frame energy and position relative to the parabolic bow shock fit  $\Delta x$  from the closest position where the solar wind core heating shock criterion is met, and the bottom two rows as a function of current particle simulation – solar wind frame energy and shock non-locality. Rows 1 and 3 are from Simulation S1, rows 2 and 4 are from S2. Black logarithmic contours indicate the counts of measurements used for evaluating mean energization., with values under 100 counts excluded. A grey band indicates the mean energy of incoming solar wind particles in the simulation frame minimum energy required for propagating upstream against the solar wind flow  $(1.9 \text{ keV for } 600 \text{ km s}^{-1})$ . It is important to note that large values of shock non-locality can indicate signals of shock structure downstream as well as upstream of the parabolic shock fit position.

- 295 of injected particles is very efficient up to about 10 keV and takes place over an extended distance. Injected particles continue to gain energy in the downstream, but begin to lose energy once back in the upstream. It is noteworthy The single most clear result seen in Figure 5 is that in general, particles above the energy corresponding with the solar wind inflow speed (1.9 keV for 600 km s<sup>-1</sup>) gain further energy, whereas particles with energies below that tend to lose energy. This is seen for both simulations S1 and S2. The first two rows, plotting energization as a function of simulation frame particle energy versus
- 300 distance from the parabolic shock fit position  $r_s(\phi)$ , show- that particles which end up injected can penetrate up to almost  $1.5r_E$  into the downstream before returning upstream, but that those particles are a minority, and at high energies and thus large gyroradii. These particles could perhaps be considered to be experiencing thermal leakage. The black contours depicting measurement counts show enhancement close to  $\Delta x = 0$  and  $E \gtrsim 1.9 \text{ keV}$ , consistent with those particles dwelling and being energized at the shock front. That that particles dwell in the vicinity of the shock, and that area is also where injected particles
- 305 may have their lowest simulation frame energies. We also see that on average, injected particles gain energy throughout the upstream region, not only at the shock front itself. On average, injected particles decelerate in the downstream. Evaluating the particle count contours, we see that injected particles gain energy in the upstream as they approach the shock, but are not accelerated above the solar wind ram energy. The final required acceleration takes place in the downstream over a distance of up to  $1.5 r_{\rm E}$ .
- 310 The behaviour of transmitted particles seen in Figure 5 is slightly different. They also start at the bottom right corner, at low energies and upstream of the shock, and experience energization already as they approach the shock. Throughout the downstream, these particles have a wide spread in energy and the dominant mechanism is to cool particles in the downstream rest frame, energizing (solar wind frame) low-energy particles and decreasing the energy of high-energy outliers. It should be noted that a small number of particles in the transmitted particles group are actually able to enter the upstream after exceeding
- 315 the solar wind kinetic energy of 1.9 keV, but the efficient deceleration there returns them to the downstream and, ultimately, the transmission boundary. Both transmitted and injected particles are able to reach energies of up to  $\sim$ 50 keV. also reach lower simulation frame energies after passing the shock fit position, but they are found throughout the downstream region (by necessity, as they must reach the transmission boundary). Particles which end up transmitted appear to gain energy also in the downstream at energies E > 1.9 keV, although the mean energization rate in that region is smaller. Similarly, the mean
- 320 deceleration rate at energies E < 1.9 keV is smaller in the downstream than in the upstream. Again, the region of efficient energization extends well upstream (up to  $1.0r_{\rm E}$ ) of the mean shock position.

The two bottom rows of Figure 5 evaluate mean energization of test-particles as a function of energy and shock non-locality. Particle count contours show that the majority of measurements are made at regions where the shock is well defined, i.e., the non-locality measure is low. However, comparing these counts with the statistics shown in of Figure 3 shows that there is

325 little to no preference for particles spending time in regions of high or low shock non-locality. Interestingly, it appears that the non-locality of the shock front does not affect mean energization rates either, as no clear preference for energization at low or high non-locality values can be seen. We do note that for simulation S1, at non-locality values of  $> 1.5 r_{\rm E}$ , we see energization also at lower energies, though this is mostly seen only for transmitted particles. Although panels i, j, m, and n do not exhibit drastic energization preference for any single non-locality value, there are a number of conclusions to draw from them. At low

- 330 energies ( $E \lesssim 1 \,\text{keV}$ ), S1 shows an energization feature at non-locality values of  $\sim 1.2 \,r_{\text{E}}$ , whereas S2 indicates more efficient energization at non-localities at around  $0.5 \,r_{\text{E}}$ . This would indicate a connection with the inherent size of foreshock structures in the two runs, respectively (Turc et al., 2018). The majority of energization of injected particles happens once particles have reached energies of  $E \gtrsim 1.9 \,\text{keV}$ , allowing them to dwell in the vicinity of the shock. Finally, at very high energies  $E \gtrsim 10 \,\text{keV}$ , a preference can be detected for energization at small values of non-locality and deceleration at large values of nonlocality,
- **335** as indicated by the predominantly red and blue regions, respectively. For transmitted particles, there appears to be no clear indication of preferential energization parameter regions.

Finally, we calculate injection probabilities  $n_{inj}/(n_{inj} + n_{tra})$  for test-particles in runs S1 and S2 as functions of a selection of parameters (detailed below) describing the first detected particle-shock-interaction. For each test-particle, we evaluate these properties at the first time- moment the particle reaches a point in the simulation space that fulfills the solar wind core heating  $(T_{core} > 4T_{sw})$  criterion for the shock. Due to the non-locality of the quasi-parallel shock front, estimating when the particle-

340  $(T_{core} > 4T_{sw})$  criterion for the shock. Due to the non-locality of the quasi-parallel shock front, estimating when the particleshock interaction is most significant is challenging, so we simply chose one of our shock criteria. but we selected the one of our three methods which we visually estimated to be most meaningful (see also panels c–f of Figure 2).

In Figure 6, we plot the estimated injection probabilities for test-particle runs using S1 and S2, using six different solar wind frame initialization energies and a Maxwellian initialization. The first two-three rows use properties of particles in 345 the simulation frame, namely the pitch-cosine  $\mu = \cos(\alpha)$  (where  $\alpha$  is the angle between the particle velocity and the local magnetic field direction), and the incidence angle (the angle between the particle direction of travel and the opposite of the bow-normal direction  $\angle(\mathbf{v}, -\hat{\mathbf{n}})$ ), and the shock-frame kinetic energy *E*.

The last three-two rows of Figure 6 use shock properties, namely the local bow-normal angle  $\theta_{Bn'}$ , and the local shock position non-locality, and the impact position noise angle. Again, these values were measured at the moment the particle first

350 encountered the shock, according to the solar wind core heating criterion. Error bars are provided by the Agresti-Coull method with a 95% confidence interval.

The first row of Figure 6 indicates that if the particle encounters the shock with negative pitch-cosine, it is likely to be injected. In our simulation set-up, most particles travel roughly in the  $-v_x$  direction, and with the IMF pointing roughly antisunward, most particles have pitch-cosines close to 1. Significant deviation from this suggests local magnetic field directions which have changed significantly due to foreshock wave effects. Our results indicate that these magnetic field deflections can enhance injection probabilities.

According to the second row, if the particle has a large incidence angle (the bow-normal velocity component is positive or small compared to the bow-perpendicular velocity component), injection is again likely. Incidence angles above 90° in fact suggest the particle was travelling away from the bow shock when it first met a shock structure. This could perhaps happen due to the particle suggest a deflected magnetic field line with a pitch angle close to zero, so that its perpendicular velocity

360

355

to the particle gyrating along a deflected magnetic field line with a pitch-angle close to zero, so that its perpendicular velocity causes it to encounter a shock peninsula such as the one seen at  $Y = 2.8 r_{\rm E}$  in Figure 2 from behind. We note that these plots show on average larger injection probabilities for higher plasma frame particle initialisation energies. This is as expected, as higher plasma frame initialization energies enable greater maximum energies when transforming into the spacecraft or simulation frame.

- 365 In the third row, we plot injection probabilities as a function of simulation frame energy, which corresponds very well with shock-frame energy due to the shock being mostly stationary on a global scale. This panel shows clearly how particles with greater initialization energies in the solar wind frame have a much larger spread in energy in the shock frame. Both very small and very large energies in the shock frame can lead to efficient injection. Small energies result in the particle spending much time at the shock, possibly then being accelerated in the shock frame with an upstream-directed velocity. Very large energies
- on the other hand mean that the particle does not need to be energized, it is enough to bend its path to the upstream in order to 370 inject it. What we also see is that particles with a higher solar wind frame initialization energy tend to have a greater chance of being injected at a given shock-frame energy. These particles have a larger velocity component tangential to the shock, which suggests that being able to perform gyromotion in the fields at the shock is important for the injection and energization process.

375

The third-fourth row shows injection probability as a function of the local bow-normal angle  $\theta_{Bn'}$ . For S1, we see a small bump for low initialization energies at  $\sim 70^{\circ}$ , and a significant increase at all energies at  $\sim 85^{\circ}$ . Considering bow-normal angles above  $90^{\circ}$  may seem odd, but these regions are where foreshock fluctuations and shock effects have caused the local magnetic field to twist back on itself. For simulation S2, with a lower Mach number, these situations are not registered.

The fourth fifth row indicates injection probability as a function of the shock non-locality measure. Both simulations S1 and S2 show a peak in injection probability at a non-locality value of  $\sim 0.4 r_{\rm E}$ , with even the lowest initialisation energies having a  $\sim 10\%$  probability in S1. For simulation S1, there is a decline in injection probability as the non-locality value increases beyond 380  $\sim 0.8 r_{\rm E}$ , with an additional peak of injection at energies > 100 eV at  $1.5 r_{\rm E}$ . These peak positions are in rough agreement with the results of Figure 5, except for the  $\sim 0.4 r_{\rm E}$  peak for S1. As that signal is very strong at all initialization energies, it may be related to a particularly strong local magnetic field twist or some other transient. Simulation S2 did not exhibit large values on non-locality so the peak at  $1.5r_{\rm E}$  cannot be verified, but there is a decrease in injection probability for 10 and 20 eV 385 initialization energies when going to non-locality values of  $\geq 0.5 r_{\rm E}$ . The fifth row, showing injection probability as a function of global position (nose angle) indicates that despite particles being energized over large radial distances, and there being no strong indication of shock non-locality affecting injection, there are regions along the shock front where local deformed shock

and magnetic field structures enhance particle injection significantly. We attempted to smooth out some of these effects by

390

launching test-particles into the simulation over a duration of 10 seconds, but an even longer test-particle initialisation and simulation extent would likely be required to smooth out all of these effects. In this study, we were limited by the fact that

simulation S2 ended at time  $t_{\rm f} = 537 \, {\rm s.}$ 

As a final step, in Table 1 we display the overall calculated injection probabilities  $N_{\rm ini}/(N_{\rm ini}+N_{\rm tra})$  per test-particle run for six test-particle initialization energies and a Maxwellian initialization. Due to the limited time period of test-particle propaga-

395

tion, at the end of the run a portion of particles were still within the shock transition zone. This is indicated by the completion ratio  $(N_{\rm ini} + N_{\rm tra})/N_{\rm init}$ . We find that the completion rate-ratio for S1 rises somewhat with increasing initialization energy, but is very stable for S2. In agreement with expectations, the injection rate increases monotonically with greater initialization energies. The injection rates for Maxwellian distributions are located between the values for 50 eV and 100 eV initializations, where the thermal speed for the 0.5 MK Maxwellian distribution is approximately 100 eV. As a point of comparison, we also extracted the Vlasiator simulation suprathermal particle densities at positions  $0.5 r_{\rm E}$  and  $1.0 r_{\rm E}$  upstream of the shock,



Figure 6. Test-particle injection probabilities for six different solar wind frame initialization energies and a 0.5 MK Maxwellian initialization and five different parameters. Left column: S1. Right column: S2. Rows 1 and 2-through 3 show properties of particles, namely the pitchcosine  $\mu = \cos(\alpha)$  and, the incidence angle, and the shock-frame energy. Rows 3 through 4 and 5 show shock properties, namely the local bow-normal angle  $\theta_{Bn'}$ , and the local shock porosity, and the impact position nose angle. non-locality. Displayed values were taken at the first encounter of each particle with the condition  $T_{\rm core} > 4T_{\rm core,sw}$ . Error bars are provided by the Agresti-Coull method with a 95% 18 confidence interval.

**Table 1.** Test-particle proton statistics using simulations S1 ( $M_A \approx 10$ ) and S2 ( $M_A \approx 5$ ) with six different solar wind frame initialization energies  $E_{init}$  and also a Maxwellian initialization distribution with a temperature of 0.5 MK. Columns list the estimated injection probability  $N_{inj}/(N_{inj} + N_{tra})$  and the completion ratio ( $N_{inj} + N_{tra}$ )/ $N_{init}$ . Also shown is the ratio of injection probabilities for S2 and S1. The final two rows show suprathermal proton density measurements  $\langle n_{p,st} \rangle$  extracted from Vlasiator simulations S1 and S2, at positions  $0.5 r_E$  and  $1.0 r_E$  upstream of the mean bow shock position, averaged over nose angles between  $\pm 45^{\circ}$  and the test-particle run time extent.

Test-particle	S1	S1	S2	S2	S2/S1
$E_{\mathrm{init}}$	injection	completion	injection	completion	injection ratio
10 eV	0.011	0.58	0.0058	0.79	0.53
20 eV	0.013	0.59	0.0063	0.79	0.48
50 eV	0.018	0.59	0.0086	0.79	0.48
100 eV	0.027	0.60	0.013	0.78	0.48
200 eV	0.047	0.62	0.027	0.77	0.75
500 eV	0.13	0.67	0.085	0.77	0.65
Maxwellian	0.021	0.59	0.010	0.78	0.48
Vlasiator	$\langle n_{\rm p,st}({\rm S1}) \rangle$		$\langle n_{\rm p,st}({\rm S2}) \rangle$		$\langle n_{\rm p,st}({\rm S2}) \rangle$
suprathermals	$n_{\rm p,sw}$		n <sub>p,sw</sub>		$\overline{\langle n_{\rm p,st}({ m S1}) \rangle}$
at $r_{\rm shock} + 0.5 r_{\rm E}$	0.042		0.061		1.45
at $r_{\rm shock} + 1.0 r_{\rm E}$	0.027		0.037		1.37

- averaged over nose angles between ±45° and between simulation times t<sub>0</sub> = 438s and t<sub>f</sub> = 538s. To facilitate comparison of these Vlasiator suprathermal particle densities (n<sub>p,st</sub>) with test-particle injection probabilities, the values are given in units of solar wind density and included as the final two rows of Table 1. We note that altough the suprathermal particle density derives from the injection probability, it measures both freshly injected protons and those protons which have spent longer in the upstream. The order of Vlasiator S1 and S2 upstream suprathermal particle densities as a function of Mach number is thus opposite to that of test-particle injection probabilities. This effect is likely not an artefact of the test-particle method, but rather results from energetic particles being trapped in the upstream, interacting with the ULF waves. Although S2 is less efficient at injection, the foreshock wave-particle trapping interactions can cause reflected particles to spend extended periods of time in the upstream before returning to the shock. Supplementary videos B and C visualize the different dynamics between simulations S1 and S2. may be caused by the differing scales of foreshock structures, with S2 foreshock dynamics causing
  elumped enhancements of reflected particles in the vicinity of the shock, and the strong ULF and SLAMS signatures in S1
- 410 champed enhancements of reflected particles in the vicinity of the shock, and the strong OEP and SEANS signatures in SP causing corresponding rarefications as well as enhancements. This phenomenon was The suprathermal particle dynamics of S1 and S2 were investigated in Turc et al. (2018), as shown in their Figure 2, panels b through d.

#### 6 Discussion

We now discuss our results presented in sections 3 and 5, attempting to clarify questions related to the non-locality of the

- 415 quasi-parallel bow shock and thermal particle injection at the Earth's quasi-parallel bow shock. We note that our approach has a number of differences compared with previous shock injection studies. We make no pre-selection that particles must encounter the shock with only a single big energization like, e.g., Sundberg et al. (2016) do. We track particle injection based on a spatial boundary, instead of requiring the ion to achieve a given energy. In our simulation the mean solar wind energy or the shock ram energy is  $E_{\rm ram} = \frac{m_i}{2} (M_{\rm A} v_{\rm A})^2 \approx 1.9 \,\rm keV$ , and a requirement of 5–10 times this energy for particle injection
- 420 (such as required by Caprioli et al., 2015) is met by approximately 40%-50% of our injected particles. We additionally note that the complicated global shock geometry used in our study prevents use of simple injection measures such as a positive  $v_x$ component (Sundberg et al., 2016). We note that in modeling the cross-shock potential we neglect the electron pressure gradient term. The majority of the potential difference at the shock is, however, included in the Lorentz and Hall terms (Eastwood et al., 2007; Yang et al., 2009).
- 425 Examination of Figure 2 shows that the spatial structure of bow shock non-locality depends on the magnitude of the upstream magnetic field, and thus, the spatial scale of foreshock structures. In Figure 3, it is evident that S1 shows clearer structures and stronger peaks of non-locality. The fine structure seen in S2 is as expected due to the increased magnetic field strength, which gives rise to smaller-scale structures in the foreshock and higher frequencies for the ULF waves (Turc et al., 2018), which in turn are expected to drive shock reformation. We suggest that spacecraft measurements of bow shock crossings could be
- 430 evaluated using our definition of non-locality, inferring tendencies for the non-locality of the bow shock versus, e.g., IMF conditions and position nose angle. Although our method was defined as a function of radial distance, it should be applicable for spacecraft time series as well.

We also investigated the energization taking place during the first shock encounter of particles, before diffusive acceleration per se. We found that all examined particles had their efficient increases and losses of energy taking place in the larger shock

- 435 transition region and extending up to  $1r_{\rm E}$  into the upstream, not limited to a narrow shock front position. particles were weakly energized over a large distance as they approached the shock, that strong energization took place at the shock and over a distance of up to  $1r_{\rm E}$  in the downstream, and that those protons which returned to the upstream experienced solar wind frame energy losses over the whole upstream region. Particles did, however, dwell for longer at the mean shock front position (panels a, b, e, and f of Figure 5). We found that the majority of reflected injected particles did not penetrate far into the
- 440 downstream, but a few did, and as they had achieved high energies, they might constitute injection through thermal leakage from the downstream. As we initialized our particles isotropic in the upstream plasma frame, we could see that particles which had simulation frame energies below the mean solar wind energy were preferentially decelerated, and particles above this energy were preferentially energized. This is opposite to what citetJohlander2016 reported, as they saw slow particles reflected at SLAMS, with fast particles passing through them. This suggest that the ubiquitous energization we see is not associated with
- 445 only SLAMS, but also other foreshock processes. A new finding which may be related is that of localized reconnection found in the quasi-parallel shock transition region citepGingell2019. well below the solar wind energy were actually preferentially

injected, similar to the SLAMS reflection test-particle studies of Johlander et al. (2016) (see panels e and f of Figure 6). Protons with shock-frame particle energies close to the solar wind ram energy were more likely to be transmitted.

- Interestingly, our result of energization taking place over a large area somewhat contradicts the results of, e.g., Guo and 450 Giacalone (2013), who in simulations of a  $M_{\rm A} = 4$  shock saw initial energization very close to the shock (within  $\sim 10 c/\omega_{ci}$  of the shock, or in our nomenclature,  $\sim 0.2 r_{\rm E}$ ). The difference may be caused by our integral energization tracking method differing from their method. The size of bow shock reformation in our simulation is (at  $\sim 50 c/\omega_{pi}$ ) in agreement with the results of, e.g., Omidi et al. (2013) and Caprioli and Spitkovsky (2013).
- We also evaluated particle energization as a function of shock non-locality , and found little dependence. For the most part, energization rates appear to be equal at all non-locality values, although at low energies each simulation showed increased energization at a non-locality length scale which appears related to the spatial scale of foreshock structures. Particles are expected to be efficiently energized between the existing shock and incoming shocklets/SLAMS or steepened ULF waves, and this is a likely explanation for this link. a weak signature of energization of slower particles could be seen if the associated shock non-locality value reached  $\sim 1.5 r_{\rm E}$ . This might be associated with bow shock bulges or erosion, as reported in citeBlanco-Cano2018.

Statistical analysis of correlations between shock and particle properties and injection probability is presented in Figure 6. The most obvious result is that there are very few particles at large incidence angles, especially at lower initialization energies. For S1, there appears to be a connection between enhanced injection probability and incidence angles close to zero. A small incidence angle will likely correlate with greater-than-average inertial simulation frame initialization energy, and higher energy

465 is known to increase injection probability. We also reported on an increase in injection probability both with increasing solar wind frame energy and with shock frame energy diverging from the solar wind ram energy.

The third-fourth row of Figure 6 highlights the importance of magnetic field deflections upstream and at the shock for efficient particle injection. Simulation S1 is much more efficient at forming strong deflections, resulting in bow-normal angles of above  $80^{\circ}$ , whereas they are absent in S2. We emphasize that these measurements were performed within the globally

- quasi-parallel region of the bow shock, between nose angles ~± 40°. We also note that in S1, there is an increase in injection at low initialization energies for bow-normal angles ≤ 15°. This is likely the same effect as what Sundberg et al. (2016) described as injected ions encountering a locally quasi-perpendicular field downstream of the shock. This also warrants further investigation. Strong deformation of magnetic fields can also lead to other forms of energization such as localized reconnection found in the quasi-parallel shock transition region (Gingell et al., 2019). Resolving these effects appears to require higher
  resolution simulations.
- 475 resolution simulations.

480

The fourth- fifth row of Figure 6 evaluates the link between shock front non-locality and proton injection. Both simulations S1 and S2 exhibit a peculiar peak in injection probability at  $\sim 0.4r_{\rm E}$ , with the peak especially strong in S1. S1 exhibits a peculiar peak in injection probability at  $\sim 0.4r_{\rm E}$ , which we presume to be due to a reformation-associated transient. S2 does not exhibit large non-locality values, but for S1, injection probability seems to fall past values  $\sim 0.9r_{\rm E}$ , with another peak at  $\sim 1.5r_{\rm E}$ . At low initialization energies, injection probabilities appear to fall off faster with increasing non-locality of the shock.

Similar to Figure 5, slight enhancements in injection can be seen at non-locality values which appear related to the size of foreshock structures in the vicinity of the shock front.

The final row of Figure 6 shows We finally note that on time scales represented in our test-particle simulations, local structures of the quasi-parallel bow shock do have a significant effect on particle injection at all initialization energies. This

485

is likely akin to what, e.g., Hao et al. (2017) and Sundberg et al. (2016) reported on, with rippled shapes of the shock front and advected magnetic fluctuations resulting in regions of localized injection. The non-locality of the quasi-parallel bow shock and its influence on the injection problem will thus need additional study and more comprehensive analysis of local shock and magnetic field structures.

The overall injection probabilities inferred from our test-particle studies agree with the strength of the shock (and the 490 Alfvénic Mach number) indicating the overall injection probability of the shock. However, we note that the suprathermal particle density registered in the upstream of the shock did not agree with this result, indicating that the evolution of suprathermal particle populations throughout the foreshock is a complicated process and not a simple indicator of local shock reflectivity. One important effect to note is that of particle trapping between foreshock waves, as reported by Wu et al. (2015). We suggest that when performing studies of shock reflectivity using spacecraft measurements, extra care should be taken to differentiate 495 freshly injected particles from an evolved foreshock population.

7 Conclusions

We have investigated the dynamics of the reforming quasi-parallel bow shock of the Earth in connection with the injection of thermal solar wind protons, using both hybrid-Vlasov and test-particle studies. Our high-fidelity- noise-free hybrid-Vlasov simulations have allowed us to probe the reforming quasi-parallel bow shock dynamics in greater detail than previously possible, accounting for correct scale separation, the global dynamics of bow shock curvature, and for effects stemming from tenuous upstream particle distributions. Our results have shown that the energization and injection of solar wind ions within this region are not local effects taking place at a single shock location, but rather, are spread out over a larger shock transition region spanning at least  $1.0 r_{\rm E}$ . We confirm enhanced particle injection with higher Alfvénic shock Mach number, and plasma

505

particle injection is enhanced. A weak enhancement could also be seen in one of our simulations at very small bow-normal angles  $\theta_{Bn'}$ , so the interaction of magnetic field directions just upstream and downstream of the shock requires further study.

frame particle energy, as expected. We also find that whenever the shock-associated magnetic field is deflected a great deal,

In our investigation, we defined a new metric for the bow shock, indicating the magnitude of non-locality of the shock front, associated with reformation. This metric was seen to correlate with the parameters of the foreshock and associated fluctuations, and also thus the shock Alfvénic Mach number. We found little to no correlation between solar wind ion injection

510 probability- only a weak link between the energization of cool solar wind frame particles and the shock non-locality, which is in agreement of with our finding of particle energization within the quasi-parallel bow shock region taking place over a large upstream extent, not only at the local or non-local shock front. At very high energies  $E \gtrsim 10 \text{ keV}$ , some preference was seen for particle energization at small values of non-locality. Although the metric was defined as a spatial measurement, it can be applied to spacecraft measurements and used to investigate the effect of shock reformation on energization of injected particles,

515 particularly at high energies.

530

Our study concentrated on two bow shock simulations, so additional studies into the locality of injection and energization of solar wind particles is warranted, using a more extensive simulation database are warranted.

We further note that the local density of suprathermal particles may be a poor indicator of injection efficiency of the shock due to large-scale dynamics of the foreshock region, such as particle trapping. This is an important factor when using either simulation results or spacecraft observations for estimating injection efficiencies at the bow shock.

*Code and data availability.* Vlasiator (http://www.physics.helsinki.fi/vlasiator/, Palmroth, 2020) is distributed under the GPL-2 open source license at https://github.com/fmihpc/vlasiator/ (Palmroth and the Vlasiator team, 2019). Vlasiator uses a data structure developed in-house (https://github.com/fmihpc/vlsv/, Sandroos, 2019), which is compatible with the VisIt visualization software (Childs et al., 2012) using a plugin available at the VLSV repository. The Analysator software (https://github.com/fmihpc/analysator/, Hannuksela and the Vlasiator team, 2020) was used to produce the presented figures. The run described here takes several terabytes of disk space and is kept in storage maintained within the CSC – IT Center for Science. Data presented in this paper can be accessed by following the data policy on the Vlasiator web site.

*Video supplement.* The Supplementary Videos A, B, and C provide movie extensions of Figures 2 and 4, showcasing the evolution of the quasi-parallel shock front profiles and the associated non-locality (Video A) and the evolution, transmission, and injection of test-particle populations of various initialization parameters for simulations S1 (Video B) and S2 (Video C).

Movie A. Movie extension of Figure 2. Animation of proton number density overlaid with bow shock positions according to criteria for plasma density (fuchsia,  $n_p = 2n_{p,sw}$ ), solar wind core heating (green,  $T_{core} = 4T_{sw}$ ), and magnetosonic Mach number (pale blue,  $M_{ms} = 1$ ). Panel (a) is for S1 ( $B_{sw} = 5 \text{ nT}$ ), panel (b) for S2 ( $B_{sw} = 10 \text{ nT}$ ), both at t = 500 s. Panels (c–f) show line profiles of the three bow shock criteria along the dashed black lines shown in panel (a), corresponding with differing amounts of shock non-locality.

- 535 Movie B. Movie extension of Figure 4. Test-particle propagation for simulation S1 ( $B_{sw} = 5 \text{ nT}$ ), with 6 different monoenergetic initialization as well as a Maxwellian 0.5 MK initialization. Vlasiator simulation proton number density is overlaid with the logarithmic density of test-particles in greyscale, with white indicating over 100 particles in a cell. Two black parabolas are the transmission boundary (left) and the injection boundary (right). Three contours indicate estimates of the local shock position: plasma compression (fuchsia,  $n_p > 2n_{p,sw}$ ), solar wind core heating (green,  $T_{core} > 4T_{sw}$ ), and the magnetosonic Mach number (pale blue,  $M_{ms} < 1$ .
- 540 Movie C. Movie extension of Figure 4. Test-particle propagation for simulation S2 ( $B_{sw} = 10 \,\mathrm{nT}$ ), with 6 different monoenergetic initialization as well as a Maxwellian 0.5 MK initialization. Vlasiator simulation proton number density is overlaid with the logarithmic density of test-particles in greyscale, with white indicating over 100 particles in a cell. Two black parabolas are the transmission boundary (left) and the injection boundary (right). Three contours indicate estimates of the local shock position: plasma compression (fuchsia,  $n_p > 2n_{p,sw}$ ), solar wind core heating (green,  $T_{core} > 4T_{sw}$ ), and the magnetosonic Mach number (pale blue,  $M_{ms} < 1$ .

545 *Author contributions*. MB carried out the analysis and wrote the manuscript. UG and YPK have made significant contributions to the simulation methods and analysis. All co-authors helped in the interpretation and vizualisation of the results, read the manuscript, and commented on it.

Competing interests. The authors declare that they have no conflict of interest.

- Acknowledgements. We acknowledge the European Research Council for Starting grant 200141-QuESpace, with which Vlasiator (Palmroth, 2020) was developed, and Consolidator grant 682068-PRESTISSIMO awarded to further develop Vlasiator and use it for scientific investigations. The Finnish Centre of Excellence in Research of Sustainable Space, funded through the Academy of Finland grant number 312351, supports Vlasiator development and science as well. We also gratefully acknowledge the Academy of Finland (grant number 267144). The CSC IT Center for Science in Finland is acknowledged for the Sisu supercomputer usage and Grand Challenge award leading to the results presented here. The work of L. Turc was supported by a Marie Sklodowska-Curie Individual Fellowship (#704681). We wish to thank Lynn
- 555 B. Wilson III and Andreas Johlander for fruitful commentary on this topic.

#### References

575

Axford, W. I., Leer, E., and Skadron, G.: The Acceleration of Cosmic Rays by Shock Waves, in: International Cosmic Ray Conference, vol. 11, 1977.

Battarbee, M., Laitinen, T., and Vainio, R.: Heavy-ion acceleration and self-generated waves in coronal shocks, Astronomy & Astrophysics, 535, https://doi.org/10.1051/0004-6361/201117507, 2011.

Bell, A. R.: The acceleration of cosmic rays in shock fronts. I, Monthly Notices of the Royal Astronomical Society, 182, 1978.

Blanco-Cano, X., Battarbee, M., Turc, L., Dimmock, A. P., Kilpua, E. K. J., Hoilijoki, S., Ganse, U., Sibeck, D. G., Cassak, P. A., Fear, R. C., Jarvinen, R., Juusola, L., Pfau-Kempf, Y., Vainio, R., and Palmroth, M.: Cavitons and spontaneous hot flow anomalies in a hybrid-Vlasov global magnetospheric simulation, Annales Geophysicae, 36, https://doi.org/10.5194/angeo-36-1081-2018, 2018.

- 565 Blandford, R. D. and Ostriker, J. P.: Particle acceleration by astrophysical shocks, The Astrophysical Journal, 221, https://doi.org/10.1086/182658, 1978.
  - Boris, J.: Acceleration calculation from a scalar potential, Tech. rep., Princeton Univ., N. J. Plasma Physics Lab., MATT-769, 1970.

Burgess, D.: Shock drift acceleration at low energies, Journal of Geophysical Research, 92, https://doi.org/10.1029/JA092iA02p01119, 1987.

- Burgess, D.: Cyclic behavior at quasi-parallel collisionless shocks, Geophysical Research Letters, 16, 570 https://doi.org/10.1029/GL016i005p00345, 1989.
  - Burgess, D.: Foreshock-shock interaction at collisionless quasi-parallel shocks, Advances in Space Research, 15, https://doi.org/10.1016/0273-1177(94)00098-L, 1995.
    - Burgess, D., Lucek, E. A., Scholer, M., Bale, S. D., Balikhin, M. A., Balogh, A., Horbury, T. S., Krasnoselskikh, V. V., Kucharek, H., Lembège, B., Möbius, E., Schwartz, S. J., Thomsen, M. F., and Walker, S. N.: Quasi-parallel Shock Structure and Processes, https://doi.org/10.1007/s11214-005-3832-3, 2005.
  - Caprioli, D. and Spitkovsky, A.: Cosmic-ray-induced filamentation instability in collisionless shocks, The Astrophysical Journal, 765, https://doi.org/10.1088/2041-8205/765/1/L20, 2013.
    - Caprioli, D., Pop, A.-R., and Spitkovsky, A.: Simulations and Theory of Ion Injection at Non-relativistic Collisionless Shocks, The Astrophysical Journal, 798, https://doi.org/10.1088/2041-8205/798/2/L28, 2015.
- 580 Caprioli, D., Yi, D. T., and Spitkovsky, A.: Chemical Enhancements in Shock-Accelerated Particles: Ab initio Simulations, Physical Review Letters, 119, https://doi.org/10.1103/PhysRevLett.119.171101, 2017.
  - Childs, H., Brugger, E., Whitlock, B., Meredith, J., Ahern, S., Pugmire, D., Biagas, K., Miller, M., Harrison, C., Weber, G. H., Krishnan, H.,
     Fogal, T., Sanderson, A., Garth, C., Bethel, E. W., Camp, D., Rübel, O., Durant, M., Favre, J. M., and Navrátil, P.: VisIt: An End-User
     Tool For Visualizing and Analyzing Very Large Data, in: High Performance Visualization–Enabling Extreme-Scale Scientific Insight, pp.
- 585 357–372, Chapman & Hall / CRC, 2012.
  - Eastwood, J. P., Lucek, E. A., Mazelle, C., Meziane, K., Narita, Y., Pickett, J., and Treumann, R. A.: The foreshock, https://doi.org/10.1007/s11214-005-3824-3, 2005.
  - Eastwood, J. P., Bale, S. D., Mozer, F. S., and Hull, A. J.: Contributions to the cross shock electric field at a quasiperpendicular collisionless shock, Geophysical Research Letters, 34, https://doi.org/10.1029/2007GL030610, 2007.
- 590 Edmiston, J. P., Kennel, C. F., and Eichler, D.: Escape of heated ions upstream of quasi-parallel shocks, Geophysical Research Letters, 9, https://doi.org/10.1029/GL009i005p00531, 1982.

Ellison, D. C.: Monte Carlo Simulation of charged particles upstream of the Earth's bow shock, Geophysical Research Letters, 8, https://doi.org/10.1029/GL008i009p00991, 1981.

Fairfield, D. H.: Bow shock associated waves observed in the far upstream interplanetary medium, Journal of Geophysical Research, 74,

595 https://doi.org/10.1029/JA074i014p03541, 1969.

625

- Gedalin, M.: Influence of the rippling on the collisionless ion and electron motion in the shock front: A model study, Journal of Geophysical Research: Space Physics, 106, https://doi.org/10.1029/2000JA000185, 2001.
  - Gedalin, M.: Transmitted, reflected, quasi-reflected, and multiply reflected ions in low-Mach number shocks, Journal of Geophysical Research: Space Physics, 121, https://doi.org/10.1002/2016JA023395, 2016.
- 600 Giacalone, J.: Shock drift acceleration of energetic protons at a planetary bow shock, Journal of Geophysical Research, 97, https://doi.org/10.1029/92JA00313, 1992.
- Gingell, I., Schwartz, S. J., Burgess, D., Johlander, A., Russell, C. T., Burch, J. L., Ergun, R. E., Fuselier, S., Gershman, D. J., Giles, B. L., Goodrich, K. A., Khotyaintsev, Y. V., Lavraud, B., Lindqvist, A., Strangeway, R. J., Trattner, K., Torbert, R. B., Wei, H., and Wilder, F.: MMS Observations and Hybrid Simulations of Surface Ripples at a Marginally Quasi-Parallel Shock, Journal of Geophysical Research:
   Space Physics, 122, https://doi.org/10.1002/2017JA024538, 2017.
- Gingell, I., Schwartz, S., Eastwood, J., Stawarz, J., Burch, J., Ergun, R., Fuselier, S., Gershman, D., Giles, B., Khotyaintsev, Y., Lavraud, B., Lindqvist, P., Paterson, W., Phan, T., Russell, C., Strangeway, R., Torbert, R., and Wilder, F.: Statistics of Reconnecting Current Sheets in the Transition Region of Earth's Bow Shock, Journal of Geophysical Research: Space Physics, https://doi.org/10.1029/2019JA027119, 2019.
- 610 Gosling, J. T., Thomsen, M. F., Bame, S. J., Feldman, W. C., Paschmann, G., and Sckopke, N.: Evidence for specularly reflected ions upstream from the quasi-parallel bow shock, Geophysical Research Letters, 9, https://doi.org/10.1029/GL009i012p01333, 1982.
  - Gosling, J. T., Thomsen, M. F., Bame, S. J., and Russell, C. T.: Ion reflection and downstream thermalization at the quasi-parallel bow shock, Journal of Geophysical Research: Space Physics, 94, https://doi.org/10.1029/JA094iA08p10027, 1989.

Greenstadt, E. W. and Mellott, M. M.: Variable field-to-normal angles in the shock foreshock boundary observed by ISEE 1 and 2, Geophys-

- 615 ical Research Letters, 12, https://doi.org/10.1029/GL012i003p00129, 1985.
  - Guo, F. and Giacalone, J.: The acceleration of thermal protons at parallel collisionless shocks: three-dimensional hybrid simulations, The Astrophysical Journal, 773, https://doi.org/10.1088/0004-637X/773/2/158, 2013.
  - Hada, T., Kennel, C. F., and Terasawa, T.: Excitation of compressional waves and the formation of shocklets in the Earth's foreshock, Journal of Geophysical Research, 92, https://doi.org/10.1029/JA092iA05p04423, 1987.
- 620 Hannuksela, O. and the Vlasiator team: Analysator: python analysis toolkit, Github repository, https://github.com/fmihpc/analysator/, last access: 23.01.2020, 2020.
  - Hao, Y., Lu, Q., Gao, X., and Wang, S.: Ion dynamics at a rippled quasi-parallel shock: 2D hybrid simulations, The Astrophysical Journal, 823, https://doi.org/10.3847/0004-637X/823/1/7, 2016.
  - Hao, Y., Gao, X., Lu, Q., Huang, C., Wang, R., and Wang, S.: Reformation of rippled quasi-parallel shocks: 2-D hybrid simulations, Journal of Geophysical Research: Space Physics, https://doi.org/10.1002/2017JA024234, 2017.
- Hietala, H., Laitinen, T. V., Andréeová, K., Vainio, R., Vaivads, A., Palmroth, M., Pulkkinen, T. I., Koskinen, H. E. J., Lucek, E. A., and Rème, H.: Supermagnetosonic Jets behind a Collisionless Quasiparallel Shock, Physical Review Letters, 103, https://doi.org/10.1103/PhysRevLett.103.245001, 2009.

- Hoppe, M. M., Russell, C. T., Frank, L. A., Eastman, T. E., and Greenstadt, E. W.: Upstream hydromagnetic waves and their association with backstreaming ion populations: ISEE 1 and 2 observations, Journal of Geophysical Research: Space Physics, 86, https://doi.org/10.1029/JA086iA06p04471, 1981.
  - Johlander, A., Vaivads, A., Khotyaintsev, Y. V., Retinò, A., and Dandouras, I.: Ion Injection at Quasi-parallel Shocks seen by the Cluster Spacecraft, The Astrophysical Journal, 817, https://doi.org/10.3847/2041-8205/817/1/L4, 2016.

Karimabadi, H., Roytershteyn, V., Vu, H. X., Omelchenko, Y. A., Scudder, J., Daughton, W., Dimmock, A., Nykyri, K., Wan, M., Sibeck, D.,

- 635 Tatineni, M., Majumdar, A., Loring, B., and Geveci, B.: The link between shocks, turbulence, and magnetic reconnection in collisionless plasmas, Physics of Plasmas, 21, https://doi.org/10.1063/1.4882875, 2014.
  - Krymsky, G., Kuzmin, A., Petukhov, S., and Turpanov, A.: Physical Principles of Regular Acceleration Mechanism of Charged Particles, in: International Cosmic Ray Conference, vol. 2 of *International Cosmic Ray Conference*, 1979.

Kucharek, H. and Scholer, M.: Origin of diffuse superthermal ions at quasi-parallel supercritical collisionless shocks, Journal of Geophysical Research, 96, https://doi.org/10.1029/91JA02321, 1991.

640

655

- Lever, E., Quest, K., and Shapiro, V.: Shock surfing vs. shock drift acceleration, Geophysical Research Letters, 28, https://doi.org/10.1029/2000GL012516, 2001.
- Liseykina, T. V., Dudnikova, G. I., Vshivkov, V. A., and Malkov, M. A.: Ion-acoustic shocks with reflected ions: modelling and particle-in-cell simulations, Journal of Plasma Physics, 81, https://doi.org/10.1017/S002237781500077X, 2015.
- 645 Lucek, E. A., Horbury, T. S., Dunlop, M. W., Cargill, P. J., Schwartz, S. J., Balogh, A., Brown, P., Carr, C., Fornacon, K.-H., and Georgescu, E.: Cluster magnetic field observations at a quasi-parallel bow shock, Annales Geophysicae, 20, https://doi.org/10.5194/angeo-20-1699-2002, 2002.
  - Lucek, E. A., Horbury, T. S., Dandouras, I., and Rème, H.: Cluster observations of the Earth's quasi-parallel bow shock, Journal of Geophysical Research: Space Physics, 113, https://doi.org/10.1029/2007JA012756, 2008.
- 650 Lyu, L. H. and Kan, J. R.: Ion leakage, ion reflection, ion heating and shock-front reformation in a simulated supercritical quasi-parallel collisionless shock, Geophysical Research Letters, 17, https://doi.org/10.1029/GL017i008p01041, 1990.
  - Malkov, M. A.: Ion leakage from quasiparallel collisionless shocks: Implications for injection and shock dissipation, Physical Review E, 58, https://doi.org/10.1103/PhysRevE.58.4911, 1998.

- Němeček, Z., Šafránková, J., Přech, L., Sibeck, D. G., Kokubun, S., and Mukai, T.: Transient flux enhancements in the magnetosheath, Geophysical Research Letters, 25, https://doi.org/10.1029/98GL50873, 1998.
  - Omidi, N., Zhang, H., Sibeck, D., and Turner, D.: Spontaneous hot flow anomalies at quasi-parallel shocks: 2. Hybrid simulations, Journal of Geophysical Research: Space Physics, 118, https://doi.org/10.1029/2012JA018099, 2013.
- 660 Onsager, T. G., Winske, D., and Thomsen, M. F.: Ion injection simulations of quasi-parallel shock re-formation, Journal of Geophysical Research, 96, https://doi.org/10.1029/91JA01986, 1991.

Palmroth, M.: Vlasiator, Web site, http://www.physics.helsinki.fi/vlasiator/, last access: 23.01.2020, 2020.

Palmroth, M. and the Vlasiator team: Vlasiator: hybrid-Vlasov simulation code, Github repository, https://github.com/fmihpc/vlasiator/, version 3.0, last access: 23.01.2020, 2019.

Malkov, M. A., Sagdeev, R. Z., Dudnikova, G. I., Liseykina, T. V., Diamond, P. H., Papadopoulos, K., Liu, C.-S., and Su, J. J.: Ion-acoustic shocks with self-regulated ion reflection and acceleration, Physics of Plasmas, 23, https://doi.org/10.1063/1.4945649, 2016.

- 665 Palmroth, M., Archer, M., Vainio, R., Hietala, H., Pfau-Kempf, Y., Hoilijoki, S., Hannuksela, O., Ganse, U., Sandroos, A., von Alfthan, S., and Eastwood, J. P.: ULF foreshock under radial IMF: THEMIS observations and global kinetic simulation Vlasiator results compared, Journal of Geophysical Research: Space Physics, https://doi.org/10.1002/2015JA021526, 2015.
  - Palmroth, M., Ganse, U., Pfau-Kempf, Y., Battarbee, M., Turc, L., Brito, T., Grandin, M., Hoilijoki, S., Sandroos, A., and von Alfthan, S.: Vlasov methods in space physics and astrophysics, Living Reviews in Computational Astrophysics, 4, https://doi.org/10.1007/s41115-018.0003.2 2018
- **670** 018-0003-2, 2018.
  - Pfau-Kempf, Y.: Vlasiator From local to global magnetospheric hybrid-Vlasov simulations, Ph.D. thesis, University of Helsinki, http: //urn.fi/URN:ISBN:978-952-336-001-3, 2016.
  - Pfau-Kempf, Y., Hietala, H., Milan, S. E., Juusola, L., Hoilijoki, S., Ganse, U., von Alfthan, S., and Palmroth, M.: Evidence for transient, local ion foreshocks caused by dayside magnetopause reconnection, Annales Geophysicae, 34, https://doi.org/10.5194/ANGEO-34-943-2016,
- 675 2016.

690

Pfau-Kempf, Y., Battarbee, M., Ganse, U., Hoilijoki, S., Turc, L., von Alfthan, S., Vainio, R., and Palmroth, M.: On the Importance of Spatial and Velocity Resolution in the Hybrid-Vlasov Modeling of Collisionless Shocks, Frontiers in Physics, 6, https://doi.org/10.3389/fphy.2018.00044, 2018.

Sandroos, A.: VLSV: file format and tools, Github repository, https://github.com/fmihpc/vlsv/, last access: 23.01.2020, 2019.

- 680 Savoini, P. and Lembège, B.: Production of nongyrotropic and gyrotropic backstreaming ion distributions in the quasi-perpendicular ion foreshock region, Journal of Geophysical Research: Space Physics, 120, https://doi.org/10.1002/2015JA021018, 2015.
  - Savoini, P., Lembège, B., and Stienlet, J.: Origin of backstreaming electrons within the quasi-perpendicular foreshock region: Twodimensional self-consistent PIC simulation, Journal of Geophysical Research: Space Physics, 115, https://doi.org/10.1029/2010JA015263, 2010.
- 685 Savoini, P., Lembège, B., and Stienlet, J.: On the origin of the quasi-perpendicular ion foreshock: Full-particle simulations, Journal of Geophysical Research: Space Physics, 118, https://doi.org/10.1002/jgra.50158, 2013.
  - Scholer, M.: Diffuse ions at a quasi-parallel collisionless shock: Simulations, Geophysical Research Letters, 17, https://doi.org/10.1029/GL017i011p01821, 1990.

Scholer, M. and Terasawa, T.: Ion reflection and dissipation at quasi-parallel collisionless shocks, Geophysical Research Letters, 17, https://doi.org/10.1029/GL017i002p00119, 1990.

- Schwartz, S. J. and Burgess, D.: Quasi-parallel shocks: A patchwork of three-dimensional structures, Geophysical Research Letters, 18, https://doi.org/10.1029/91GL00138, 1991.
  - Schwartz, S. J., Thomsen, M. F., and Gosling, J. T.: Ions upstream of the Earth's bow shock: A theoretical comparison of alternative source populations, Journal of Geophysical Research, 88, https://doi.org/10.1029/JA088iA03p02039, 1983.
- 695 Schwartz, S. J., Burgess, D., Wilkinson, W. P., Kessel, R. L., Dunlop, M., and Lühr, H.: Observations of short large-amplitude magnetic structures at a quasi-parallel shock, Journal of Geophysical Research, 97, https://doi.org/10.1029/91JA02581, 1992.

Sckopke, N., Paschmann, G., Bame, S. J., Gosling, J. T., and Russell, C. T.: Evolution of ion distributions across the nearly perpendicular bow shock: Specularly and non-specularly reflected-gyrating ions, Journal of Geophysical Research, 88, https://doi.org/10.1029/JA088iA08p06121, 1983.

700 Su, Y., Lu, Q., Gao, X., Huang, C., and Wang, S.: Ion dynamics at supercritical quasi-parallel shocks: Hybrid simulations, Physics of Plasmas, 19, https://doi.org/10.1063/1.4752219, 2012.

- Sundberg, T., Boardsen, S. A., Slavin, J. A., Uritsky, V. M., Anderson, B. J., Korth, H., Gershman, D. J., Raines, J. M., Zurbuchen, T. H., and Solomon, S. C.: Cyclic reformation of a quasi-parallel bow shock at Mercury: MESSENGER observations, Journal of Geophysical Research: Space Physics, 118, https://doi.org/10.1002/jgra.50602, 2013.
- 705 Sundberg, T., Haynes, C. T., Burgess, D., and Mazelle, C. X.: Ion acceleration at the quasi-parallel bow shock: decoding the signature of injection, The Astrophysical Journal, 820, https://doi.org/10.3847/0004-637X/820/1/21, 2016.
  - Thomas, V. A. and Winske, D.: Two dimensional hybrid simulation of a curved bow shock, Geophysical Research Letters, 17, https://doi.org/10.1029/GL017i009p01247, 1990.
- Thomsen, M. F., Schwartz, S. J., and Gosling, J. T.: Observational evidence on the origin of ions upstream of the Earth's bow shock, Journal of Geophysical Research, 88, https://doi.org/10.1029/JA088iA10p07843, 1983.
  - Tóth, G., Chen, Y., Gombosi, T. I., Cassak, P., Markidis, S., and Peng, I. B.: Scaling the Ion Inertial Length and Its Implications for Modeling Reconnection in Global Simulations, Journal of Geophysical Research: Space Physics, 122, https://doi.org/10.1002/2017JA024189, 2017.
    - Turc, L., Ganse, U., Pfau-Kempf, Y., Hoilijoki, S., Battarbee, M., Juusola, L., Jarvinen, R., Brito, T., Grandin, M., and Palmroth, M.: Foreshock Properties at Typical and Enhanced Interplanetary Magnetic Field Strengths: Results From Hybrid-Vlasov Simulations, Journal
- of Geophysical Research: Space Physics, 123, https://doi.org/10.1029/2018JA025466, 2018.
- Turc, L., Roberts, O. W., Archer, M. O., Palmroth, M., Battarbee, M., Brito, T., Ganse, U., Grandin, M., Pfau-Kempf, Y., Escoubet, C. P., and Dandouras, I.: First Observations of the Disruption of the Earth's Foreshock Wave Field During Magnetic Clouds, Geophysical Research Letters, 46, https://doi.org/10.1029/2019GL084437, 2019.
- von Alfthan, S., Pokhotelov, D., Kempf, Y., Hoilijoki, S., Honkonen, I., Sandroos, A., and Palmroth, M.: Vlasiator: First global
- 720 hybrid-Vlasov simulations of Earth's foreshock and magnetosheath, Journal of Atmospheric and Solar-Terrestrial Physics, 120, https://doi.org/10.1016/j.jastp.2014.08.012, 2014.
  - Wilson III, L. B.: Low Frequency Waves at and Upstream of Collisionless Shocks, in: Low-Frequency Waves in Space Plasmas, American Geophysical Union (AGU), https://doi.org/10.1002/9781119055006.ch16, 2016.
  - Wilson III, L. B., Sibeck, D. G., Breneman, A. W., Contel, O. L., Cully, C., Turner, D. L., Angelopoulos, V., and Malaspina, D. M.: Quantified
- 725 energy dissipation rates in the terrestrial bow shock: 2. Waves and dissipation, Journal of Geophysical Research: Space Physics, 119, https://doi.org/10.1002/2014JA019930, 2014a.
  - Wilson III, L. B., Sibeck, D. G., Breneman, A. W., Contel, O. L., Cully, C., Turner, D. L., Angelopoulos, V., and Malaspina, D. M.: Quantified energy dissipation rates in the terrestrial bow shock: 1. Analysis techniques and methodology, Journal of Geophysical Research: Space Physics, 119, https://doi.org/10.1002/2014JA019929, 2014b.
- 730 Wu, M., Hao, Y., Lu, Q., Huang, C., Guo, F., and Wang, S.: The role of large amplitude upstream low-frequency waves in the generation of superthermal ions at a quasi-parallel collisionless shock: CLUSTER observations, The Astrophysical Journal, 808, https://doi.org/10.1088/0004-637X/808/1/2, 2015.

Yang, Z. W., Lu, Q. M., and Wang, S.: The evolution of the electric field at a nonstationary perpendicular shock, Physics of Plasmas, 16, https://doi.org/10.1063/1.3275788, 2009.

735 Zank, G. P., Rice, W. K. M., Le Roux, J., Cairns, I. H., and Webb, G. M.: The injection problem for quasiparallel shocks, Physics of Plasmas, 8, https://doi.org/10.1063/1.1400125, 2001.