Final response for "Measurements of aerosols and charged particles on the BEXUS18 stratospheric ballon"

Dear Editor,

Thank you very much for providing us the opportunity to submit a revised version of our paper to your journal.

Please find below the comments received by two anonymous reviewers after submission to ANGEOD, followed by our replies and finally our changes in the manuscript.

1. Comments from Referees

Anonymous Referee #1

The contribution by Erika Brattich and colleagues reports the measurement and modelling of charged aerosols in the stratosphere. The manuscript is very well written, logically constructed, easy to follow and informative. The manuscript starts with a substantial review of the relevant literature, followed by a thorough theory section that explains the basis for the simulations. Compared to these first two sections, the consecutive section on the experimental results is rather terse and provides little guidance to the reader as to how the individual findings reported in the list of Figures contribute to the key points of the paper. As a result, it would be helpful to expand this section to make the narrative more clear. It is also somewhat surprising that Fig. 5 is not listed in this section, perhaps because it is not considered to be a result of the conducted work. The final sections with the discussion and conclusions emphasise to a large degree the agreement of the findings with previous work. While it is undoubtedly important to put the findings of this study into context, it makes is harder for the reader to appreciate the novelty of the presented work which becomes less clear. It therefore appears to be beneficial for these two sections to distinguish more clearly between known facts and novel findings. Besides this apparent imbalance between the first and second part of the manuscript, I think it is a valuable contribution to the scientific literature as the current knowledge on charged aerosols in the stratosphere and their spatiotemporal variabilities is somewhat limited at present.

Some minor suggestions on how to improve the manuscript are given below.

(1) Fig 2: The concentrations of negative ions appear to be large compared to previous findings. Is there any explanation for this? It is also not clearly explained how the total concentration of aerosols can be smaller than the concentration of negative ions. Is the reader supposed to infer from this that the aerosols <200 nm mainly contribute to the negative ions?

(2) Fig 3: 19 channels are listed in the legend, but only 8 height dependent traces can be distinguished. It is practically impossible to infer any useful information for the PBL.

(3) Fig 4: The x-axis labels are rather sparse and could be more populated.

(4) Fig 5: Again, only 9 curves are shown for 19 channels listed in the legend, as in Fig. 3. Would it not be better to combine some of these channels for the benefit of clarity?

(5) The arrangement in the table appears somewhat unfortunate to me. The first two rows seem to be unrelated to the remainder of the table and the table deserves a heading to state the unit (nm) for the first column and a symbol with unit for the second column.

(6) The acknowledgments have distinct font variations disturbing this reader.

Anonymous Referee #2

General comments:

This study reports the aerosol measurements obtained by the BEXUS18 stratospheric balloon flight. Its distinct feature is being equipped with ion counters. A role of ion chemistry in the stratosphere is still an open question and could be essential for the understanding of the atmospheric impact from the space. This topic is suitable for ANGEO. However, information on the instruments of aerosol and ion measurements is completely lacking in this manuscript, so that it is impossible to evaluate whether their observations are reliable or not. Thus I recommend a rejection of this manuscript.

Detailed comments are given below.

Specific comments:

-Temperature and pressure measurements

The authors showed results of temperature and pressure measurement in Fig. 1, but it is found that their results are not so reliable compared to nearby radiosonde observation. Including a radiosonde in their payload does not look difficult, so that I am wondering why they did not do it. In addition, they used temperature and pressure data in their model calculation. Is it really meaningful to use such unreliable data? They need to show how sensitive their model calculation is to temperature and pressure errors. -Aerosol measurements

Their aerosol instrument, LOAC, has been used in 150 flights, so that its precision, resolution, etc. should be well known. However, they did not give those information at all in the manuscript. Although they mentioned an existence of the thin aerosol layer with a thickness of less than 100m at p.10, 1.25, I cannot judge whether this instrument has a vertical resolution high enough to detect such a layer. -Ion measurements

They mentioned that the performance of their ion measurement was checked by preflight lab experiments, but it is not shown in the manuscript at all. Thus I cannot judge whether their ion measurements are reliable or not.

-Average

In order to show the aerosol data, they often used arithmetic mean/smoothing. Since aerosol density changes by several orders of magnitude, the arithmetic mean strongly depends on the largest value. Geometrical mean or median filter would be better to represent aerosol distributions

-Figs. 3 and 4

What is dN/dlog(D) in Fig. 3? A caption of Fig. 4 does not correspond to Fig. 4 about their y-axis.

-Eq. (3) Units look different between the terms. Probably some variables are missing.

2. Author's response

Anonymous Referee #1 General comments

We thank the reviewer for his/her constructive comments. We have included Fig.5 in the narrative of the experimental results section. In addition, the final sections (Sections 4 Results, 5 Discussion, and 6 Conclusions) were revised in accordance to the guidelines provided by the reviewer.

(1) The detection of high concentrations of negative ions is probably due to the fact we added a separate suitable instrument for measuring ions in this flight. Because of the LOAC lower limit of detection of aerosols at 200 nm, we can expect to have ions concentrations greater than the aerosol detected total concentrations due to the presence of aerosols with aerodynamic diameter less than 200 nm. Since the number of ions is greater than the detected aerosol, this indicates that aerosol smaller than 200 nm are the main contributor to negative ions. This comment was added in the revised version of the manuscript.

(2) The text in the revised version of the manuscript was changed to better describe the vertical profiles of the different particles' sizes presented in the Figure. Information on the fact that the information on the PBL, partially commented but out of the scope of the paper, was also added. The combination of the large size-classes in a few super-size classes can be misleading and potentially losing information on the real size of the biggest particles.

(3) The x-axis is in logarithmic scale; however, ticks were added to have a more populated x-axis.

(4) As previously replied, the combination of the large size-classes in a few super-size classes can be misleading and potentially losing information on the real size of the biggest particles.

(5) The arrangement of the table is rather customary for a correlation table: the table presents the correlation coefficients between the variables presented in each row (here, ions and particles' number detected in each size range) and those presented in the columns (here, positive and negative ions). The units were added to the table.

(6) The font variations in the acknowledgements were removed in the revised version of the paper.

Anonymous Referee #2

General comments:

We thank the reviewer for his/her comments. The revised version of the manuscript now contains information on the instruments of aerosol and ion measurements, as will be detailed more precisely in the following answers.

Specific comments:

-Temperature and pressure measurements

In the revised version of the manuscript, we have addressed and provided the results of the sensitivity tests of the model simulations to changes in the T-p profile. In particular, the change in temperature and pressure profile affect only the ion aerosol attachment (approximately 10% change for 20% change in temperature) and the charge particle coagulation coefficient (approximately 6% change for 20% change in temperature and pressure. This does not affect the final model results drastically, as steady state conditions are reached in a couple of hours. As shown in the next Figure, which presents the results of sensitivity test with 20% difference in the input temperature profile, only 1% change in aerosol concentration is observed for an input temperature profile 20% higher/lower in the model input. Because of this, repeating the model calculations with the Kandalashka T-p profile, no significant differences are observed in the final result. To conclude, the T-p profile only changes the rate of the reaction, but not steady-state concentrations.



-Aerosol measurements

We thank the reviewer for his/her comment. Indeed, this information was missing from the previous version of the manuscript. Instead of applying a smoothing procedure, in the revised version we have integrated the raw measurements over 5 minutes. Thus, we have changed the figures 2 and 3 according to this new procedure. Explanations for this are now provided in the text.

-Ion measurements

Indications of the performance of the ions' measurements as derived by preflight lab experiments are provided in the revised version of the manuscript, in particular in the form of median absolute deviations separately for positive and negative ions measurements at the 200 mbar pressure level.

-Average

We thank the reviewer for his/her comment. As previously replied to the comment related to the aerosol measurements, the aerosols data are now not smoothed, thus not averaged.

-Figs. 3 and 4

dN/dlog(D), or DN/dln(D), represents a commonly used notation in aerosol science to indicate the number concentration of particles in the various size classes (dN) divided by the width of the size classes (dln(D)).

The caption of Figures 3 and 4 was slightly modified to indicate more correctly what is shown in the Figure.

-Eq. (3)

Eq. 3 has been corrected in the revised version of the manuscript.

3. Author's changes in the manuscript

We have expanded the Results section inserting also Figure 5 and better presenting our results as shown later in our replies to the reviewers' comments. We have added in the discussion section, page 13, lines 8-11, a summary of our main findings:

"Summarizing, our observations first of all demonstrate the effectiveness of the adopted instrumental setup in measuring vertical profiles of particles' size distributions and particles' typology together with ions. In addition, they can also provide interesting results in terms of the association between cosmic rays and ions, and further to reveal novel features in terms of the charged fraction, from new stratospheric flights with a similar instrumental setup.

(1) We have added in Section 4, page 10, lines 29-32:

"It is important to note first of all that since the lower limit of detection of LOAC is for particles presenting an optical diameter of 200 nm, we can expect to have ions concentrations greater than the

aerosol detected total concentrations. In particular, since the number of ions is greater than the detected aerosol, we can infer that aerosol smaller than 200 nm are the main contributors to negative ions."

(2) We have added better description of the vertical profiles of the different particles' sizes presented in the Figure. The text at page 11, lines 1-8 is now:

"Fig. 3 reports the vertical profiles of aerosol size distribution for each size bin measured by the LOAC instrument. Most of the particles have size below 1 μ m, as expected in a clean free troposphere and in the stratosphere. Few particles greater than 1 μ m and smaller than 15 μ m and just one 50 μ m particle were detected in the stratosphere. All fine particles (< 1 μ m) presented the same vertical variation, with a global trend of decreasing concentrations at heights higher than the tropopause. Larger particles, besides presenting lower number concentrations as expected, presented a different vertical profile, with the presence of an abrupt increase in the PBL and then at 10 km less evident in finer particles. However, it is important to note that with these measurements it is difficult to derive information on the PBL, which is out of the focus of this paper."

(3) The x-axis of Figure 4 was populated with more ticks.

(4) Since the combination of the large size-classes in a few super-size classes can be misleading and potentially losing information on the real size of the biggest particles, no change was made.(5) We added the units to the parameters in Table 1.

(6) The font variations in the Acknowledgements were removed in the revised version of the paper.

-Temperature and pressure measurements

We have added in the experimental section about numerical simulations (Section 3), page 9, lines 25-26 and page 10, lines 1-6:

"Sensitivity tests to temperature and pressure indicate that the change in temperature and pressure profile affect only the ion aerosol attachment coefficient (approximately 10% change for 20% change in temperature and pressure) and charge particle coagulation coefficient (approximately 6% change for 20% change in temperature and pressure). This does not affect the final model results drastically, as results show that steady state conditions are reached in more or less a couple of hours. Only 1% change in aerosol concentration is observed for an input of 20% higher/lower temperature profile into the model (reported in the Supplementary Material). Overall, no significant differences are observed for 20% change in T-p profile. The T-p profile only changes the rate of the reaction, but not steady state concentrations."

As indicated in the text, a Figure showing the results of sensitivity tests for 20% higher/lower temperature is now provided in the Supplementary Material.

-Aerosol measurements

We have added in the text in Section 2.1, at page 6, lines 19-22:

"The LOAC vertical resolution is linked to the total concentrations of aerosols, as due to the Poisson counting statistics and the instrument capability to detect the smallest particles. A detailed analysis of the raw measurements has shown that the data must be integrated over 5 minutes to remove the oscillations due to the measurements' uncertainty. Considering the balloon ascent speed, this procedure provides a resolution of about 1 km."

And later at page 7, lines 19-20:

"To compare directly with the aerosols' measurements, the ions' measurements data are also integrated with a 1-km vertical resolution."

-Ion measurements

We have added in the text, in Section 2.2, at page 7, lines 7-9:

"Those tests indicate that the average MAD (median absolute deviation) of ions' measurements was equal to 15 ions at 200 mbar for negative ions, and 7 ions for positive ions at the same pressure level."

-Average

Modifications to the text are reported in the previous response to the comment on aerosol measurements -Figs. 3 and 4

No change is made but for the captions.

Figure 3: Vertical profiles of particles size distributions for the 19 size classes of the LOAC particle counter as part of the A5-Unibo experiment on the BEXUS18 stratospheric flight. The notation dN/dlog(D) used in the x-axis stands for the number concentration of particles in the various size classes divided by the width of the size classes

Figure 4: Average size distribution of aerosol particles at various height layers during BEXUS18 stratospheric flight. The five curves are obtained averaging the aerosol number densities as a function of the atmospheric layers pointed out by the temperature profile as follows: 1: 0-404 m (black line); 2: 650-1519 m (red line); 3: 1765-10118 m (blue line); 4: 10650-25044 m (pink line); 5: 25289-27191 m (green line). X- and Y-axes are in log-normal scales.

-Eq. (3)

The Equation (3) was corrected in the revised version of the manuscript:

 $\frac{dN_{i,j}}{dt} = \beta_{i-1,j}^+ N_{i-1,j} n^+ + \beta_{i+1,j}^- N_{i+1,j} n^- - \beta_{i,j}^+ N_{i,j} n^+ - \beta_{i,j}^- N_{i,j} n^- + \frac{1}{2} \sum_{l,m=-p}^{l,m=p} \int_0^v K_{j-\nu,\nu}^{l,m} N_{j-\nu}^m d\nu - \sum_{l,m=-p}^{l,m=p} \int_0^v K_{j-\nu,\nu}^{l,m} N_{j-\nu,\nu}^m d\nu + \sum_{l=0}^{l,m=p} \int_0^v K_{j-\nu,\nu}^m N_{j-\nu}^m d\nu + \sum_{l=0}^{l,m=p} \int_0^v K_{j-\nu,\nu}^m d\nu + \sum_{l=0}^{l,m=p} \int_0^\infty K_{j-\nu,\nu}^m d\nu + \sum_{l$

 $N_{i,j} \sum_{q=-p}^{p} \int_{0}^{v} K_{j,v}^{i,q} N_{v}^{q} dv$ (3)

Measurements of Aerosols and Charged Particles On the BEXUS18

Stratospheric Balloon

Erika Brattich¹, Encarnación Serrano Castillo², Fabrizio Giulietti², Jean-Baptiste Renard³, Sachi N. Tripathi⁴, Kunal Ghosh⁴, Gwenael Berthet³, Damien Vignelles³, Laura Tositti⁵

⁵ ¹Department of Physics and Astronomy DIFA, University of Bologna, 40126 Bologna (BO), Italy ²Industrial Engineering Department, University of Bologna, 47121 Forlì (FC), Italy ³LP2CE-CNRS/Université d'Orléans, 45071 Orléans, France ⁴Department of Civil Engineering & Department of Earth Sciences, Indian Institute of Technology, 208016, Kanpur, India ⁵Department of Chemistry "G. Ciamician", University of Bologna, 40126 Bologna (BO), Italy

10 Correspondence to: Erika Brattich (erika.brattich@unibo.it)

Abstract. This paper describes the aerosol measurements setup and results obtained during the BEXUS18 stratospheric balloon within the "A5-Unibo" (Advanced Atmospheric Aerosol Acquisition and Analysis) experiment performed on October 10th, 2014 in northern Sweden (Kiruna). The experimental setup was designed and developed by the University of Bologna with the aim of collecting and analyzing vertical profiles of atmospheric ions and particles together with

- 15 atmospheric parameters (temperature, relative humidity and pressure) all along the stratospheric ascent of the BEXUS18 stratospheric balloon. Particles size distributions were measured with the MeteoModem Light Optical Aerosol Counter (LOAC) and air ion density was measured with a set of two commercial and portable ion counters. Though the experimental setup was based upon relatively low-cost and light-weight sensors, vertical profiles of all the parameters up to an altitude of about 27 km were successfully collected. The results obtained are useful for elucidating the relationships between aerosols
- 20 and charged particles between ground level and the stratosphere with great potential in collecting and adding useful information in this field, also in the stratosphere where such measurements are rare. In particular, the equipment detected coherent vertical profiles for particles and ions, with a particularly strong correlation between negative ions and fine particles, possibly resulting from proposed associations between cosmic rays and ions as previously suggested. In addition, the detection of charged aerosols in the stratosphere is in agreement with the results obtained by a previous flight and with
- 25 simulations conducted with a stratospheric ion-aerosol model. However, further measurements under stratospheric balloon flights equipped with a similar setup are needed to reach general conclusions on such important issues.

Keywords: 0305 Aerosols and particles; 0394 Instruments and techniques

1 Introduction

It is well-recognized that aerosols play a fundamental role in the lower atmosphere as they may affect climate with both a direct effect on absorption and scattering of solar radiation but also an indirect effect through cloud processing (Yu and

- 5 Turco, 2001; Forster et al., 2007). Aerosols are tightly involved in the atmospheric chemical mass balance, including the stratospheric chemistry through the heterogeneous reactions with nitrogen and halogen species triggering the austral ozone hole through Polar Stratospheric Clouds (PSCs) (Hanson et al., 1994; Deshler, 2008). Aerosol still represents the largest uncertainty in the correct estimate and interpretation of the ongoing change in Earth's energy budget (Boucher et al., 2013, IPCC, 2013; Myhre et al., 2013). In this framework, it is therefore of paramount importance to accurately and systematically
- 10 collect experimental data such as particle number densities, as well as all the properties shedding light on their nature, their size distribution, and their source in order to define both qualitatively and quantitatively their role, in the troposphere as well as stratosphere.

Stratospheric aerosols are contributed by several sources which determine particle size, composition and morphology, as well as their mean residence time. Historically, the first measurements of stratospheric aerosol were carried out by Junge

- 15 (Junge, 1961; Junge et al., 1961); stratospheric aerosols drew the attention of scientists during the Cold War owing to the artificial radioactivity released into the stratosphere and returned to the troposphere through the Stratosphere-to-Troposphere exchange processes (Corcho Alvarado et al., 2014; Feely et al., 1966). The monitoring of radioactive fallout from nuclear weapon testing (and in 1964 from the accident of SNAP9A, a nuclear-fueled satellite which released ²³⁸Pu in the upper atmosphere, upon navigation failure, Eisenbud and Gesell, 1997) not only brought about the understanding of the basic
- 20 dynamic processes coupling the troposphere and stratosphere, but also the discovery of cosmogenic radionuclides which have their maximum production in the stratosphere, mostly in the form of aerosols, still largely employed in the study of vertical exchange between the innermost atmospheric layers (Cristofanelli et al., 2018). Beside radionuclides, the main source of aerosol particles in the stratosphere is through the flux of sulfur bearing molecules into the stratosphere from the troposphere, primarily, OCS (atmospheric carbonyl sulphide), during non-volcanic times, and SO₂ from volcanic eruptions:
- 25 after release, sulfur is oxidized and converted to sulfuric acid which then condenses forming the bulk of the stratospheric aerosol layer (e.g., Kremser et al., 2016). Secondary sources comprehend the outer space, contributing an array of mineral micrometeoritic particles mainly in the solid phase, *in situ* emission from aircrafts (Murphy et al., 1998) and the troposphere itself, through active upward transportation; the troposphere may also act as a source through more localized exchange processes likely mediated by cumulo-nimbus dynamics.
- 30 Tropospheric sources include volcanic eruptions, usually occurring on an event basis and with a strong dependency on the event energy (see for example Deshler, 2008 or Murphy et al., 2014), as well as the transport in the Tropical Tropopause Layer (TTL; ~ 12-18 km) of air and (water-insoluble) gases and particles driven by cumulonimbus cloud and further upward

transport within the TTL (and in the tropical lower stratosphere) due to the Brewer-Dobson circulation (e.g., Buchart, 2014). Fromm et al. (2000) also showed that major forest fires can locally inject large number of carbonaceous and potassium rich aerosols in the lower stratosphere, often internally mixed with meteoritic smoke (Hervig et al., 2009; Neely et al., 2011) and solid grains survived from their atmospheric entry (Cziczo et al. 2001; Renard et al., 2005). In this framework, additional

5 evidence of tropospheric contribution has been shown lately by Yu et al. (2017), who indicated a major role of the Asian summer monsoon in contributing more than 15% of the particles in the stratosphere, potentially emphasizing anthropogenic aerosol contributions.

Overall, while recent progress both in stratospheric observations and research suggests the need for increasing attention and consequently for data collection on stratospheric aerosol, most of the investigations on this topic are still mainly focused on

- 10 volcanic contributions which constitute a sort of baseline to assess behavior and properties of stratospheric aerosols themselves, but also inspire potential as well as very arguable global warming countermeasures under the wide term of geoengineering (Launder and Thompson, 2009). Indeed, volcanic eruptions strongly influence particle populations in the stratosphere, in particular those from El Chichon in 1982 and from Mt. Pinatubo in 1991 (e.g. Russell et al., 1996), whose monitoring in the course of the past decades brought to assess how stratospheric aerosols may be strongly affected by
- 15 extreme events until their slow removal, when they reach background concentration levels. Recently, it was shown that even weaker volcanic events, similar to biomass burning, are capable to access at least the lower stratosphere suggesting the need for further investigation and monitoring (Robock, 2000). As a result, using various measurement techniques such as remote sensing observations from satellites and *in situ* measurements is needed in order to get an updated view of the stratospheric system and its connections with upper and lower layers of the atmosphere (e.g. Steele et al., 1999; Deshler et al., 2003).
- 20 It is commonly assumed that stratospheric aerosols are mostly liquid consisting of sulfuric acid from volcanic eruptions, while the upper stratosphere is free of aerosols except for some instances of residual particles from meteoritic disintegration and for interplanetary grains in very low concentrations (Murphy et al., 2014). Nevertheless, it seems that the stratospheric aerosol content is more complex, both in terms of aerosol concentrations and nature (Renard et al., 2008). Vernier et al. (2011) showed that moderate volcano eruptions can inject a significant number of aerosols into the stratosphere, refilling the
- 25 aerosol layer episodically. Though less investigated than tropospheric aerosols, extensive details on the properties of stratospheric aerosols are provided in the recent review by Murphy et al. (2014). In this framework, another physical property still highly underscored, if not in very specialized science fields, is represented by accord close here here the properties of differently signal inter here here the properties of the property still highly underscored.

by aerosol electrical characteristics. Air conductivity due to the presence of differently sized ions has long been recognized and studied as reviewed in Clement and Harrison (1991), Hirsikko et al. (2011), Harrison and Carslaw (2003). Charged

30 tropospheric aerosols were detected not only in disturbed weather but also in fair weather atmosphere, as resulting from ion diffusion. Charged particles have also been detected in volcanic ashes (Gilbert et al., 1991; Harrison et al., 2010) and in Saharan dust layers (Nicoll et al., 2011). In the mesosphere, smoke and ice particles are part of the plasma in the D-region and carry positive and negative charges (Rapp, 2009). In the stratosphere, electrified aerosols have been detected in situ for the first time during a balloon-borne aerosol counting measurements (Renard et al., 2013). The observations carried on

during that stratospheric flight showed that most of the aerosols are charged in the upper troposphere from altitudes below 10 km and in the stratosphere from altitudes above 20 km, while the aerosols seem to be uncharged between 10 km and 20 km. The electrification of the aerosols could originate from ion clusters produced mainly in the atmosphere by the interaction of galactic cosmic rays with the atmospheric gases especially in the dense regions of the planetary atmospheres where solar extreme ultra violet radiation is absent (Harrison and Carslaw, 2003).

Major sources of ions in the atmosphere include radon isotopes, cosmic rays, and terrestrial gamma radiation, with a variable relative contribution depending on the altitude and latitude (Tinsley, 2008): while ionization from turbulent transport of radon and gamma radiation prevail near the Earth's surface and over the continents, ionization due to cosmic rays dominates far away from the continental surface (i.e., over the oceans and in the upper stratosphere/lower troposphere) (Hirsikko et al.,

5

- 10 2011) and where also the production of cosmogenic radionuclides is highest (Tositti et al., 2014 and references therein). Both primary and secondary ionization may therefore interact with air components to produce air ions. While the overall air ion population is largely responsible for the so-called "atmospheric global electric circuit" (Tinsley, 2008), the detection of charged particles in the stratosphere is extremely important since they might have affected both sprite formation and stratospheric photochemistry (e.g., Belikov and Nikolayshvili, 2016 and references therein).
- 15 In this framework, the detection of ions and charged particles across the atmospheric column is ever increasingly drawing attention due to the experimental evidence linking ions to nucleation mechanism, firstly proposed by Raes and Janssens (1985). Requiring a smaller supersaturation of the involved gases, ion-induced nucleation is thermodynamically advantaged over homogeneous nucleation. Even though experiments still do not agree on the relative contributions of ions and neutral nucleation (e.g., Eisele et al., 2006; Suni et al., 2008; Yu, 2010), this effect is at the basis of the proposed link between the
- 20 flux of ionizing galactic cosmic rays modulated by solar activity and the global cloud cover (Svensmark and Friis-Christensen, 1997; Carslaw, Harrison and Kirkby, 2002) which predicts that an increase in cosmic ray intensity causes an enhancement in CCN (cloud condensation nuclei) abundance and therefore of cloud reflectivity and lifetime (by suppressing rainfall). This hypothesis stems from the observed enhancement of cloud cover during peaks of high energy radiation leading to enhanced particle formation and growth in the presence of ions: depending on the competition between condensation
- 25 growth and processes reducing particle concentrations (i.e., coagulation, surface deposition, and in-cloud scavenging), a fraction of those particles may eventually grow into the size of CCN. This mechanism, unlike the aerosol indirect effect, is only driven by changes in the rates of microphysical processes and acts on a global scale being stronger in regions of low aerosol concentrations.

A second link between galactic cosmic rays and global cloud cover, the ion-aerosol near-cloud mechanism (Tinsley et al.,
 2000), involves the effects of cloud microphysical properties due to the accumulation of space charge on the tops of clouds. This mechanism is less understood than the former one, but is linked to the hypothesis that variations in cosmic ray ionization might modulate the fair-weather current generated by the electrical current flowing into clouds leading to a sequence of both micro and macro-physical responses in cloud processing (Dunne et al., 2012).

Observations conducted to study and quantify these effects (e.g., Laakso et al., 2007; Svensmark et al., 2007; Pierce and Adams, 2009) are often incomplete and non-conclusive leaving model simulations as the most convenient source of information. The search for a link between CRs (Cosmic Rays) and cloud formation is also one of the main drivers for the CLOUD experiment being conducted at CERN since 2009 where a chamber filled with atmospheric gases is crossed by

- 5 charged pions that simulate ionizing CRs. While some preliminary results suggested that indeed IIN (Ion Induced Nucleation) is a relevant factor to determine nucleation rates in the upper troposphere (e.g., Kirkby et al., 2011), recent results show that cosmic ray intensity cannot meaningfully affect climate via nucleation (Dunne et al., 2016) while others indicate that IIN of pure organic particles constitutes a potentially widespread source of aerosol particles in terrestrial environments with low sulfuric acid pollution (Kirkby et al., 2016).
- Stratospheric balloon research devoted to the collection of aerosol profiles vs. height have traditionally been carried out in the last decades with the aim of elucidating properties and processes of this fundamental air component. While as a rule, in situ observations, either onboard stratospheric balloons either onboard aircrafts, are fairly demanding and costly (e.g., Hunton et al., 2005; Curtius et al., 2005; Andersson et al., 2013; Watanabe et al., 2004; Murphy et al, 2014; Sugita et al., 1999; ; Matsumura et al., 2001; Hervig and Deshler, 2002; Kasai et al., 2003; Deshler et al., 2003; Shiraishi et al., 2011), the
- 15 present paper aims at promoting the formation of young researchers in the spirit of the BEXUS initiative (Balloon-borne Experiments for University Students, see below), but also in elaborating effective and "relatively" cheap experiments to fulfill the need of experimental vertical profiles of aerosol data useful for filling knowledge gaps in atmospheric and climatological research. The "A5-Unibo" (Advanced Atmospheric Aerosol Acquisition and Analysis) experiment designed by the University of Bologna has been developed with the purpose of collecting and studying vertical profiles of atmospheric
- 20 ions and particles in addition to atmospheric parameters (temperature, relative humidity and pressure) all along the flight path of the BEXUS18 stratospheric balloon, using a relatively low-cost and low-weight setup compared to the conventional instrumentation onboard stratospheric balloons. This paper describes the setup of the experiment and the measurements obtained during the flight.

25 2 Instrumentation

30

The "A5-UNIBO" experiment was flown from SSC, Esrange Space Center in northern Sweden (Kiruna; 67°53'N, 21°04'E) on October 10th, 2014 with the BEXUS18 stratospheric balloon under the REXUS/BEXUS program. The program was realized under a bilateral Agency Agreement between the German Aerospace Centre (DLR) and the Swedish National Space Board (SNSB). The Swedish share of the payload is available through collaboration with the European Space Agency (ESA).

The balloon was a Zodiac BL-DD-12SF-404-ZIT filled with helium gas with a volume of 12,000 m^3 and a diameter of 14 m. The flight lasted for 3 hours from 08:48 to 12:00, and reached a maximum altitude of 27.2 km with an average ascending

speed during climbing of $3.5m \text{ s}^{-1}$. Due to onboard electric problems, no data was available between the altitude range of 18.5 km and 20 km. The floating time in the stratosphere was 1 hour and 8 minutes. The balloon eventually landed in Finland where it was promptly retrieved and safely brought back to the Esrange Space Center facilities the day after the flight.

5 Apart from the "A5-UNIBO" experiment whose results and setup are herein described, the whole payload of the stratospheric balloon was quite big and included a wide range of different experiments: the AFIS-P (Antiproton Flux In Space-Prototype), ARCA (Advanced Receiver Concepts for ADS-B), COUGAR (Control of Unmanned Ground Vehicle from Higher Altitude in near Real Time), and POLARIS (POLymer-Actuated Radiator with Independent Surfaces)

2.1 Aerosol measurements

- 10 Particles size distribution vertical profile was measured by the Light Optical Aerosol Counter (LOAC) (MeteoModem Inc.), an optical particle counter/sizer (Renard et al., 2016a,b based on scattering measurements at angles of 12° and 60°. The instrument is light (250 g total weight including the pump) and compact enough to perform measurements on board of all kinds of balloons. As described in detail in Renard et al. (2016a,b), the combination of the measurements at two scattering angles provides both the determination of the particle size distribution and an estimation of the typology of particles in 19
- 15 size classes from 0.2 to 100 µm: briefly, while the measurement at 12° scattering angle does not depend on the refractive index of the particles and enables for accurate size determination and counting, the measurement at 60° scattering angle is strongly sensitive to the refractive index of the particles, giving information on the nature of the particles. LOAC has already performed more than 150 flights in the stratosphere since 2013.

The LOAC vertical resolution is linked to the total concentrations of aerosols, as due to the Poisson counting statistics and

20 the capability of the instrument ability to detect the smallest particles. A detailed analysis of the raw measurements has shown that the data must be integrated over 5 minutes to remove the oscillations due to the measurements'29 uncertainty. Considering the balloon ascent speed, this procedure gives provides a resolution of about 1 km.

2.2 Ion measurements

- 25 Air ion densities vertical profile was measured by means of two air ion counters (ALPHALAB Inc.), respectively for positive and negative ions. These instruments are handheld meters designed to measure ion density, i.e., the number of ions per cubic centimetre (ions/cc) in air. The instrument is a ion density meter, based on a Gerdien Tube condenser design, and containing a fan which draws air through the meter at a calibrated rate and is able to count the number of positive/negative ions when the voltage applied to the outer cylinder is positive/negative respectively. The air ion counter can be used for the
- 30 detection of natural and artificial ions. Natural ions include those generated from the decay of radioactive minerals and radon gas, fires, lightning, and evaporating water, and finally ions associated with storm activity. These devices are light and small enough (305 g each; 160 x 100 x 55 mm) to be mounted on balloons; their measurement resolution extends up to 200000

ions cm⁻³. These probes are equipped with a fan to create an air flow of 24 l min⁻¹ throughout the gondola. Air ambient ions are diverted from the flow and collected on a plate which returns a voltage output proportional to the number of ions. The air is then expelled downwards through the bottom plate.

The use of these small commercial probes based on Gerdien tube meter (see for example Aplin and Harrison, 1999) for stratospheric balloon experiments is unprecedented, therefore quality control procedures for ion quantification are not vet

- available; however, the instrumental performance of the air ion counters at stratospheric conditions was tested in pre-flight lab experiments simulating stratospheric conditions. <u>Those tests indicate that the average MAD (median absolute deviation)</u> of ions' measurements was equal to 15 ions at 200 mbar for negative ions, and 7 ions for positive ions at the same pressure level.
- 10 The ion counters were operated for offset values first in a vacuum changing the pressure between 1000 and 5 hPa and subsequently in a thermal chamber between 15°C and -60°C.

The result of these tests was that both of the air ion counters worked properly in low-pressure environments, while the offset was found to be independent of external pressure. It is also worth pointing out that during this experiment the fan flow rate was expected to remain constant during the balloon's ascent phase. Even though the flow rate actually could monotonously

- 15 decrease with increasing altitude, thus leading to underestimations in in on density concentrations with increasing altitude, preliminary tests performed to assess the flow rate dependence on pressure were not conclusive and did not provide a satisfactory working curve. However, even if ion concentration variations might be biased by the pressure dependence of the air flow rate, this bias does not affect the relative variations of concentrations (local strong increases or decreases), which in fact variations seem to be consistent, as shown and discussed later on.
- 20 <u>To make direct comparison</u> compare directly with the aerosols's measurements, the ions measurements data awere integrated with a 1-km vertical resolution.

2.3 Temperature, relative humidity measurements and other instrumentation

5

The BEXUS-18 gondola was also equipped with a Parallax MS5607 altimeter module for pressure readings, which was successfully tested at 120,000 feet. A humidity sensor HIH9120-021 was used to record the vertical profile of relative humidity. A temperature sensor LM35DZ was mounted on the electronic board to control the internal temperature. External temperature and GPS data was instead retrieved by the Esrange Balloon Service System (EBASS), a Telemetry/Telecommand (TM/TC) service system for stratospheric balloons developed by SSC and DLR in 1998.

All the sensors and the instruments onboard (especially the air ion counters and the LOAC) were tested prior to the 30 stratospheric flight in a vacuum chamber to ensure their proper functioning at ambient pressures from 1000 to 5 mbar. In addition, after complete assembling of all the probes, the whole experiment was put into a thermal chamber to ensure its proper working at low temperatures conditions. All the tests confirmed the correct performance of the experiment under stratospheric conditions. For a detailed test related to the LOAC performance, the reader is referred to Renard et al. (2016a). An Arduino MEGA 2560 microcontroller was used for data acquisition from sensors and instruments. The sensors were connected to ARDUINO through a hardware interface and two stacked boards. An Arduino Ethernet Shield was used to connect the Arduino MEGA 2560 board to the BEXUS telemetry system. Two additional electronic boards were designed for the respective control of the heating system which kept the temperature of the key components above 0°C and of a power control unit which fed the probes and sensors with the required voltage and current.

Data acquired by the on-board unit, including ambient data and internal sensors, was collected with a 10 second time resolution; the data was transmitted to the Ground Station and displayed via HID through a graphical interface. The data was then integrated over 60 seconds and only data acquired during the ascent and floating phases was analyzed.

3 Numerical simulations

5

- 10 Model calculations have been used to quantify the electrification of aerosols with a stratospheric ion-aerosol model in the altitude range of LOAC measurements. Ion clusters in the atmosphere are produced primarily by interaction of galactic cosmic- rays with atmospheric gases, especially in the dense regions of planetary atmospheres where extreme solar ultraviolet radiation is absent (Harrison and Carslaw, 2003). A high fraction of the cosmic ray (1 GeV) energy flux is typically carried by particles of high-kinetic energy. The peak ion production rate by this process has been found to be
- 15 generally located at altitudes between 14 and 17 km (Rawal et al., 2013), which is our major study area, and the ion pair production rate is calculated using the statistical model of O'Brien (2005), considering SO_4^{2-} and NH_4^+ as the most abundant ion clusters produced by this process- (Renard et al., 2013). Other sources (radon isotopes and terrestrial gamma radiation) can be included for the further improvement of the model simulation as one of the future scopes of the study.
- This ion pair production rate is calculated using the statistical model developed by O'Brien (2005) with the major ions 20 considered being SO_2^{-4} and NH^{+4} . Electrons are not included in the model as they recombine with positive ions and uncharged molecules very rapidly, and are consequently not available to interact with aerosols. The charging of aerosols is calculated using charge balance equations as described by Michael et al. (2008, 2009) and Tripathi et al. (2008). The charged particles are constantly interacting with each other resulting in changes in initial charge and size distribution with time. As this is a bipolar interaction, it is expected that charge distribution will be wider than initial distribution with time (Ghosh et 25 al., 2017).

$$\frac{dn^{+}}{dt} = q - \alpha n^{+} n^{-} - [n^{+} \sum_{j=rmin}^{rmax} \sum_{l=-p}^{p} \beta_{l,j}^{+} N_{l,j}]$$
(1)
$$\frac{dn^{-}}{dt} = q - \alpha n^{+} n^{-} - [n^{-} \sum_{j=rmax}^{rmax} \sum_{j=1}^{p} \beta_{l,j}^{-} N_{l,j}]$$
(2)

i=rmin i=-p

In equation (1) and (2), n^+ and n^- represent positive and negative ion concentrations, respectively, q is the ion pair production rate, β is the ion-ion recombination coefficient, N is the aerosol concentration, and α is the ion-aerosol attachment rate. Here the radii of the aerosols vary from size rmin to rmax, and the maximum number of elementary charges an aerosol can own is p. The aerosol concentration for any size and charge is calculated by Eq. (3), where *i* represents the number of elementary charges on a particle for *i*, the associated radius bin.

5

25

$$\frac{dN_{i,j}}{dt} = \beta_{i-1,j}^{+} N_{i-1,j} n^{+} + \beta_{i+1,j}^{-} N_{i+1,j} n^{-} - \beta_{i,j}^{+} N_{i,j} n^{+} - \beta_{i,j}^{-} N_{i,j} n^{-} + \frac{1}{2} \sum_{l,m=-p}^{l,m=-p} \int_{0}^{v} K_{j-v,v}^{l,m} N_{j-v}^{l} N_{v}^{m} dv - N_{i,l} \sum_{q=-p}^{p} \int_{0}^{v} K_{i,v}^{l,m} N_{v}^{q} dv$$
(3)

In equation (3), the first two terms on the right-hand side are the probability of interaction between the ions and aerosols of 10 any particular charge and size, and the last term is for growth of that particle due to the charge-particle coagulation process. K is the charge particle coagulation coefficient (probability of collision between two charged particles). v is the aerosol particle volume assuming all particles have a spherical shape, N_s is the number of aerosol particles for any particular size. Full details are provided in Ghosh et al. (2017).

The model is run for amount of charges of any particular size of particle (q) running from +20 to -20. The ion-aerosol 15 attachment coefficients (β) are calculated in different ways depending on the relative size of the particles with the ionic mean free path. The calculation depends in particular from the different regimes, i.e. diffusion, free molecular and transition. Hoppel and Frick (1986) developed a method to calculate in all three different regimes. The major requirements for this calculation are the ionic mobility and mean free path, which are calculated using the expressions given by Borucki et al. (1982). The charge coagulation coefficient K was calculated from diffusional force (including vertical diffusion), turbulent

20 shear force, turbulent inertial force along with electrostatic force due to charge on particles (Ghosh et al., 2017).

A polydispersed distribution of aerosols is used in the model and is obtained from the LOAC observation. The LOAC measured data was used for calculation of different input parameters, like ionic mobility, charge particle mobility, charge particle coagulation, ion-aerosol attachment coefficient and ion-ion recombination, which are the global input parameters for the overall model (Global Electrical Circuit Model (GEC), as described in Rawal et al. (2013)). The model also uses temperature and pressure measured during the experiments. Sensitivity tests to temperature and pressure indicate that the change in the input temperature and pressure profile affects only the ion aerosol attachment coefficient (approximately 10%

change for 20% change in temperature and pressure) and charge particle coagulation coefficient (approximately 6% change for 20% change in temperature and pressure). This does not affect the final model results drastically, as results show that steady state conditions are reached in more or less a couple of hours. Only 1% change in aerosol concentration is observed for an input of 20% higher/lower temperature profile into the model (reported in the Supplementary Material). Overall, no

5 significant differences are observed for 20% change in T-p profile. The T-p profile only changes the rate of the reaction, but not steady state concentrations. We added charged particle coagulation model to the Renard et al. (2013) model, as it is close to accurate simulation scenario. The charge balance equations are solved by implicit numerical method to obtain concentrations of positive ions, negative ions, uncharged aerosols and charged aerosols, for the steady state.

4 Results

- 10 Fig. 1 shows the comparison between the vertical profiles of relative humidity and temperature measured during the flight with those measured with the radiosonde sounding at Kandalashka (67.15 N, 32.35 E, 25m asl; Russia), 12UTC. The comparison of the temperature profiles shows good agreement in the troposphere with a small inversion layer close to the ground and the starting of the inversion typical of the tropopause located at about 11 km up; The comparison of the temperature profile in the stratosphere presents instead major differences: in fact, while onboard the Bexus flight the 15 temperature remains almost constant in the stratosphere up to an altitude of 26 km and presents a sudden and strong increase around 26-27 km, the temperature profile measured at Kandalashka presents a small decrease until about 25-26 km, typical of profiles of this time of the year in the Arctic region. The reason of such discrepancy in the stratosphere might be due to
- solar heating and perhaps to heat/solar light reflection from other instruments/structures of the gondola. The comparison of the relative humidity profiles presents instead major differences already in the troposphere: in particular, the strong dryness in the Planetary Boundary Layer (PBL) detected onboard the Bexus flight (less than 20%) with a further decrease until the altitude of about 5 km is probably due to the slow response of the relative humidity sensors used onboard. Since standard radiosonde measurements of relative humidity are only reliable in the troposphere above temperatures near -40°C, whereas below these temperatures and in the stratosphere special instrumentation for stratospheric water vapor measurements is

needed (Berthet et al., 2013; Tomikawa et al., 2015), measurements of relative humidity above the tropopause are reported

- 25 only for the Kandalaksha profile, which have to be treated with care nevertheless. Fig. 2 reports the vertical profiles for the cumulative aerosol particle number density obtained by summing up the data from all the size bins collected by the LOAC (in black) together with the negative (blue) and positive (red) ions during the ascent. A sliding smoothing (i.e., each point is simply replaced with the average of m adjacent points) is applied to suppress small scale fluctuations. It is important to note first of all that since the lower limit of detection of LOAC is for particles presenting
- 30 an optical aerodynamic diameter of 200 nm, we can expect to have ions concentrations greater than the aerosol detected total concentrations. In particular, since the number of ions is greater than the detected aerosol, we can infer that aerosol smaller than 200 nm are the main contributors to negative ions.

Fig. 3 reports the vertical profiles of aerosol size distribution for each size bin measured by the LOAC instrument. Most of the particles have size below 1 μ m, as expected in a clean free troposphere and in the stratosphere. Few particles greater than 1 μ m and smaller than 15 μ m and just one 50 μ m particle were detected in the stratosphere. , which shows that aAll fine particles (< 1 μ m) presented the same vertical variation, with a global trend of decreasing concentrations at heights higher

5 than the tropopause. L-while-larger particles, besides presenting lower number concentrations as expected, presented a different vertical profile, with the presence of an abrupt increase in the PBL and then at 10 km less evident in finer particles. However, it is important to note that with these measurements it is difficult to derive information on the PBL, which is out of the focus of this paper.

In Fig. 4 we report the average variation of particle size distribution with altitude. The five size distributions depicted have

been determined empirically by averaging the LOAC data over five temperature intervals along the height profile (see Fig. 1) according to roughly coherent atmospheric layers, as obtained during the BEXUS-18 experiment. The intervals chosen were respectively: 0 - 404 m; 650 - 1519 m; 1765 - 10118 m; 10650 - 25044 m; and 25289 - 27191m.

In practice, they correspond respectively to lower, upper PBL, free troposphere, tropopause/lower stratosphere, and mid stratosphere, which as known is characterized by a marked increase in temperature owing to the ozone absorption of longer

15 wavelength UV radiation. However, the marked temperature increase recorded at float in our measurements is probably due to instrumental errors since the relative speed between the balloon and the air is close to 0 there (no ventilation). While there is a steady decrease of particles in all the size bins as the altitude increases, in the intermediate tropospheric groupings an increase in the coarse particle bins around and above 10 µm is observed.

Finally, Fig. 5 reports the vertical profiles of simulated fraction of charged particles, which can be used for comparison with
 measured profiles and with previous simulations and observations presented by Renard et al. (2013) with completely different instrumentation

5 Discussion

The data collected shows that there is a steady decrease in the particle number density with height for all the size bins determined. At all heights sub-micron particles are the most numerous, though coarse particles show a relative, sensitive increase in the upper PBL and free troposphere. At an altitude of 1 km, LOAC typology measurements indicate the presence of a thin layer, less than 100m width, of transparent particles, possibly droplets. At the other altitudes the typology measurements indicate optically absorbing and semi-absorbing particles, probably related to the presence of minerals (dust). In particular, above the tropopause, almost all of the particles detected by LOAC were smaller than 1 µm, and their typology

30 measurements indicate both the occurrence of stratospheric liquid droplets and the presence of optically absorbing material (i.e., internally or externally mixed particles): even though LOAC typology measurements cannot provide precise information on particles' chemical composition, <u>it</u> is in agreement with results from aircraft observations in the lower stratosphere (e.g. Schwartz et al., 2006, Murphy et al., 2014), which showed that while sulfate particles dominate the aerosol composition in the stratosphere, other sources producing absorbing particles also contribute. The vertical profiles of integrated concentration of aerosols > 200 nm and of ions (Figure 2) present interesting features. Firstly, positive charges are only present relatively close to the ground, which is in agreement with previous observations (e.g., Li et al., 2015), even

- 5 though we cannot exclude that the complete absence of positive charges at upper levels derives from a failure of the positive ions counter during the flight. Even though we are performing more flights with a similar instrumental setup in order to compare and provide evidence of our findings, this general behavior is also consistent with previous observations, showing that ionization from turbulent transport of radon (positively charged product ions) and gamma radiation (negative ions) prevail close to the Earth's surface, whereas ionization from cosmic rays (negative ions) dominates away from the
- 10 continental surface (upper troposphere and above) (Hirsikko et al., 2011), both ion distribution playing a basic role in the terrestrial global circuit (Tinsley and Zhou, 2006). Indeed, preliminary results of a stratospheric flight with the ion counters performed on 8 April 2017 in Australia show and confirm the detection of only positive ions in the lower troposphere, while both polarities, with a prevalence of negative ions, were present at upper levels. Secondly, vertical profiles of particles and ions present the same general structure above the tropopause, including an enhancement in the 20-25 km altitude range.
- 15 None of these profiles are similar to the temperature, humidity, and pressure profiles, which can exclude the possibility of instrumental contamination by these atmospheric parameters. Moreover, the ion concentration variations between 10 and 20 km cannot be linked to the decrease in the airflow fan, whose precise dependence on pressure cannot be correctly estimated for the time being as previously pointed out in the material and methods section. Even though the absolute values could be biased from this effect, the relative variations seem to be real. In particular, Spearman's correlation coefficient (a
- 20 nonparametric measure of rank correlation, where nonparametric means not based on parameterized families of probability distributions) (Table 1) indicates a strong negative correlation (i.e., anticorrelation) of negative ions with fine particles, a behavior which might have resulted from ion induced nucleation and in particular from the proposed association between cosmic rays and ions ("ion-aerosol clear-air" mechanism), although we are aware that nucleation concerns particles in the 1-2 nm range, and further growth is governed by condensation. Our results are also in agreement with previous observations
- 25 showing that, in general, negative ions more efficiently promote nucleation than positive ions (Eisele et al., 2006; Suni et al., 2008; Svensmark and Friis-Christensen, 1997).

The weaker, but positive correlation between positive ions and coarse particles might instead arise from their simultaneous detection closer to the Earth's surface.

The number of (negative) charges as well as of particles strongly increases above 20 km. The maximum value around 20 km 30 corresponds to the region of maximum ionization (Regener-Pfotzer maximum (Regener and Pfotzer, 1935)) and was previously observed by Harrison et al. (2014) observing count rates through Geiger counters on standard meteorological balloons. Stratospheric ion-aerosol model simulations can be used to quantify and explain the electrification of the aerosols (Rawal et al., 2013). The simulated profile shows that more than 75% of aerosols are charged above the altitude of 5 km (Figure 5). This result is in agreement with the presence of the charged liquid and/or solid particles detected by ion detectors

and could be used as an estimate of the vertical variability of their percentage. The measurements presented here are also in general good agreement with the unique previous direct detection of charged stratospheric aerosols of Renard et al. (2013). In addition, they reveal a "depletion layer" of poorly charged aerosols from the tropopause to an altitude of about 20 km, where the charged fraction drops at about 1%, similar to the one previously detected by Renard et al. (2013).

5 In particular, as from Figure 5 b, it is clear that as from model simulations, fine particles are the ones contributing to the largest variations in the fraction of the charged fraction, while coarse particles, when present, are mostly charged, confirming the calculations made by Renard et al. (2013) (see their figure 4).

Summarizing, our observations first of all demonstrate the effectiveness of the adopted instrumental setup in measuring vertical profiles of particles' size distributions and particles' typology together with ions. In addition, they can also provide

10 interesting results in terms of the association between cosmic rays and ions, and further to reveal novel features in terms of the charged fraction, from new stratospheric flights with a similar instrumental setup.

6 Conclusions

The A5-UNIBO experiment flown under a stratospheric balloon seems to have confirmed the previous detection of charged aerosols in the stratosphere and a possible vertical variability. In particular, the results show coherent vertical profiles for particles and ions, with a particularly strong correlation between negative ions and fine particles, possibly resulting from proposed associations between cosmic rays and ions as previously suggested. Due to the important implications of charged aerosols on the high-energy phenomena (sprites, blues jet and elves) in the middle stratosphere (Fullekrug et al., 2016) and of ions in nucleation mechanisms, further stratospheric balloon-borne measurements of charged particles are necessary.

- 20 Poly-instrumented gondolas with aerosols counters for the estimate of the percentage of charged particles, positive and negative ions counters, and Geiger counter, will help to better evaluate the direct link between cosmic rays, ions, and the charged aerosols. Furthermore, the addition of small condensation particle counters able to characterize particles in the 1-2 nm range could help to gain precise information on nucleation, which here was only derived and could not directly observed. In particular, since both the present flight and the Renard et al. (2013) flight were performed for altitudes below 27 km, an
- 25 experimental campaign comprehensive of many flights at higher altitude, up to the maximum altitude reachable with stratospheric balloons (around 40 km), is necessary to better document the vertical evolution of the charged aerosols. Also, the analysis of the results obtained during such future flights will be helpful in answering open questions raised in this and previous flights:
 - Are the stratospheric aerosols always charged?
- Is there variability in the percentage of charged aerosols, for example with latitude, season, or solar eruptions?

- Is the percentage of charged aerosols dependent on the nature of the aerosols (liquid droplet, ash from volcanic eruptions, meteoritic mineral material, or carbonaceous particles from Earth and space)?
- Is the "depletion layer" of poorly charged aerosols above the tropopause, not expected from modeling, a transient phenomenon or a permanent feature?
- 5 Such aerosols measurements could have implication on climate and atmospheric chemistry issues, but also on the atmospheric electricity and high energy phenomena such as sprites, blue jets and elves that are not yet well understood.

Acknowledgements

- This work was designed and developed within the collaboration of the Flight Mechanics Laboratory (Prof. Fabrizio Giulietti) and the Environmental Chemistry and Radioactivity Laboratory of the Department of Chemistry "G. Ciamician" (Prof. Laura Tositti) of the 10 University of Bologna as main supporters. In particular, the experiment was flown onboard the BEXUS18 stratospheric balloon under the REXUS/BEXUS program, supported under a bilateral Agency Agreement between the German Aerospace Centre (DLR) and the Swedish National Space Board (SNSB). The Swedish share of the payload is available through collaboration with the European Space Agency (ESA). Every team member of the A5-UNIBO experiment is acknowledged for his/her essential role for the success of the experiment: Encarnación Serrano Castillo (Team Leader and System Engineer), Riccardo Lasagni Manghi (Verification and Testing engineer), Erika Brattich (Data Analysis, Scientific expert), Igor Gai (Ground Station engineer), Danilo Boccadamo (Power engineer), Paolo Lombardi (Mechanics), Alice 15 Zaccone (Software engineer), Abramo Ditaranto (Electronics engineer), Luca Mella (Software engineer), and Marco Didonè (Thermal engineer). We also acknowledge: a) institutional supporters: DLR, Rymdstyrelsen, SSC, ESA Education Office, EuroLaunch, ZARM; b) private companies and associations: AlphaLab Inc., Boxer, Iacobucci HF Aerospace, Icos, CNA Forlì-Cesena, Dogcam, Gruppo SDS, Società Italiana di Medicina Generale, Plastica Panaro, Bustaplast, Bellini Tiziana, Mascherpa. We also thank Sergio Brattich for a thorough revision of the English language of 20 the manuscript. Erika Brattich also thanks the Department of Biological, Geological and Earth Sciences of the University of Bologna for grant
- support during her PhD study, during which the experiment was developed, and the Department of Chemistry "G. Ciamician" of the University of Bologna, for support during her post-doc

Formattato: Tipo di carattere: 9 pt, Inglese (Stati Uniti), Bordo: : (Nessun bordo)

Formattato: Tipo di carattere: 9 pt, Inglese (Stati Uniti), Bordo: : (Nessun bordo)

Formattato: Tipo di carattere: 9 pt, Bordo: : (Nessun bordo)
Formattato: Tipo di carattere: 9 pt, Inglese (Stati Uniti), Bordo: : (Nessun bordo)
Formattato: Tipo di carattere: 9 pt, Bordo: : (Nessun bordo)
Formattato: Tipo di carattere: 9 pt

References

Andersson, S.M., Martinsson, B.G., Freiberg, J., Brenninkmeijier, C.A.M., Rauthe-Schöch, A., Hermann, M., van Velthoven, P.F.J., and Zahn, A.: Composition and evolution of volcanic aerosol from eruptions of Kasatochi, Sarychev and Eyjafjallajökull in 2008-2010 based on CARIBIC observations. Atmos. Chem. Phys., 13, 1781-1796, doi:10.5194/acp-13-1781-2013, 2013.

5

Aplin, K.L., and Harrison, R.G.: The interaction between air ions and aerosol particles in the atmosphere. Inst. Phys. Conf. Ser., 163, 411-414. Paper presented at the 10th Int. Conf., Cambridge, 28-31 March 1999, available at https://arxiv.org/ftp/arxiv/papers/1209/1209.4549.pdf, , 1999.

Arnold, F.: Ion nucleation-a potential source for stratospheric aerosols. Nature, 299, 134-37, 1982

10 Arnold, F.: Atmospheric ions and aerosol formation. Space Sci. Rev., 137, 225-239, doi:10.1007/s11214-008-9390-8, 2008.

Belikov, Y., and Nikolayshvili, S.: The role of the dipole interaction of molecules with charged particles in the polar stratosphere. Journal of Earth Science and Engineering, 6, 115-149: doi:10.17265/2159-581X/2016.03.001, 2016.

Berthet, G., Renard, J.-B., Ghysels, M., Durry, G., Gaubicher, B., and Amarouche, N.: Balloon-borne observations of midlatitude stratospheric water vapour: comparisons with HALOE and MLS satellite data. J. Atmos. Chem., 70, 197-219, doi:10.1007/s10874-013-9264-7, 2013.

15

Boruchi, W.J., Levin, Z., Whitten, R.C., Keesee, R.G., Capone, L.A., Toon, O.B., and Dubach, J.: Predicted electrical conductivity between 0 and 80 km in the Venusian atmosphere. Icarus, 51, 302-321, 1982.

Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S.K., Sherwood, S., Stevens, B., and Zhang, X.Y.: Clouds and aerosols, In: Climate

Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the 20 Intergovernmental Panel on Climate Change. [Stocker T.F., et al. (eds.)] Cambridge University Press, Cambridge, United Kingdom, and New York, USA, pp. 571-657, 2013.

Buchart, N.: The Brewer-Dobson circulation. Rev. Geophys., 52(2), doi:10.1002/2013RG000448, 2014

Carslaw, K.S., Harrison, R.G., and Kirkby, J.: Cosmic rays, clouds and climate. Science, 298, 1732-1737, doi:10.1126/science.1076964, 2002. 25

Clement, C. F. and Harrison, R. G.: Charge distributions on aerosols. Institute of Physics Conference Series, 118, 275-280, 1991.

Corcho Alvarado, J.A., Steinmann, P., Estier, S., Bochud, F., Haldimann, M., and Froidevaux, P.: Anthropogenic« radionuclides in atmospheric air over Switzerland during the last few decades. Nature Communications, 5,

Formattato: Interlinea: 1.5 righe

30 doi:10.1038/ncomms4030, 2014. Formattato: Interlinea: 1.5 righe

- Cristofanelli, P., Brattich, E., Decesari, S., Landi, T.C., Maione, M., Putero, D., Tositti, L., and Bonasoni, P.: Chapter 5. Studies on Environmental Radionuclides at Mt. Cimone. In: High-Mountain Atmospheric Research. The Italian Mt. Cimone WMO/GAW Global Station (2165 m a.s.l.). Springer Briefs in Meteorology, Springer, Cham, ISBN 978-3-319-61126-6, doi:10.1007/978-3-319-61126-6, 2018.
- 5 Curtius, J., Weigel, R., Vössing, H.-J., Wernli, H., Werner, A., Volk, C.-M., Konopka, P., Krebsbach, M., Schiller, C., Roiger, A., Schlager, H., Dreiling, V., and Borrmann, S.: Observations of meteoric material and implications for aerosol nucleation n the winter Arctic lower stratosphere derived from in situ particle measurements. Atmos. Chem. Phys., 5, 3053-3069, 2005.
- Cziczo, D. J., Thomson, D. S., and Murphy, D. M.: Ablation, flux and atmospheric implication of meteors inferred from stratospheric aerosols. Science, 291, 1772-1775, doi:10.1126/science.1057737, 2001.

Deshler, T.: A review of global stratospheric aerosol: measurements, importance, life cycle, and local stratospheric aerosol. Atmos. Res, 90, 2-4, 223-232, doi:10.1016/j.atmosres.2008.03.016, 2008.

Deshler, T., Hervig, M. E., Hofmann, D. J., Rosen, J. M., and Liley, J. B.: Thirty years of in situ stratospheric aerosol size distribution measurements from Laramie, Wyoming (41°N) using balloon-borne instruments. J. Geophys. Res., 108, D5 4167, doi:10.1029/2002JD002514, 2003.

Dunne, E.M., Lee, L.A., Reddington, C.L., and Carslaw, C.S.: No statistically significant effect of a short-term decrease in the nucleation rate on atmospheric aerosols. Atmos. Chem. Phys., 12, 11573-11587, doi:10.5194/acp-12-11573-2012, 2012.

15

Dunne, E.M., Gordon, H., Kürten, A., Almeida, J., Duplissy, J., Williamson, C., Ortega, I.K., Pringle, K.J., Adamov, A., Baltensperger, U., Barmet, P., Benduhnm, F., Bianchi, F., Breitenlechner, M., Clarke, A., Curtius, J., Dommen, J., Donahue,

- 20 N.M., Ehrhart, S., Flagan, R.C., Franchin, A., Guida, R., Hakala, J., Hansel, A., Heinritzi, M., Jokinen, T., Kangasluoma, J., Kirkby, J., Kulmala, M., Kupc, A., Lawler, M.J., Lehtipalo, K., Makhmutov, V., Mann, G., Mathot, S., Merikanto, J., Miettinen, P., Nenes, A., Onnela, A., Rap, A., Reddington, C.L.S., Riccobono, F., Richards, N.A.D., Rissanen, M.P., Rondo, L., Sarnela, N., Schobesberger, S., Sengupta, K., Simon, M.,, Sipilä, M., Smith, J.N., Stozkhov, Y., Tomé, A., Tröstl, J., Wagner, P.E., Wimmer, D., Winkler, P.M., Worsnop, D.R., and Carslaw, K.S.: Global atmospheric particle formation from 25 CERN CLOUD measurements. Science, 354, 6316, 1119-1124, doi:10.1126/science.aaf2649, 2016.
- Eisele, F.L., Lovejoy, E.R., Kosciuch, E., Moore, K.F., Mauldin, III R.L., Smith, J.N., McCurry, P.H., and Iida, K.: Negative atmospheric ions and their potential role in ion-induced nucleation. J. Geophys. Res., 111, D04305, doi:10.1029/2005JD006568, 2006.
- Eisenbud, M., and Gesell, T.: Environmental Radioactivity from Natural, Industrial and Military Sources. 4th Edition.
 30 Academic Press, ISBN 9780080505800, 1997.

Enghoff, M.B., and Svensmark, H.: The role of atmospheric ions in aerosol nucleation – a review. Atmos. Chem. Phys., 8, 4911-4923, doi:10.5194acp-8-4911-2008, 2008

Feely, H. W., Seitz, H., Lagomarsino, R. J., and Biscaye, P. E.: Transport and fallout of stratospheric radioactive debris. Tellus, 18, 2-3, 316-328, doi:10.1111/j.2153-3490.1966.tb00243.x, 1966

- 5 Forster, P., Ramaswamy, V., Artaxo, P., Bernsten, T., Betts, R., Fahey, D.W., Haywood, J., Len, J., Lowe, D.C., Myhre, G., Nganga, J., Prinn, R., Raga, G., Schultz, M., and Van Dorland, R.: Changes in atmospheric constituents and in radiative forcing. In: Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. [Solomon S. et al. (eds.)] Cambridge University Press, Cambridge, United Kingdom and New York, USA, pp 129-134, 2007.
- 10 Fromm, M., Alfred, J., Hoppel, K., Hornstein, J., Bevilacqua, R., Shettle, E., Servranckx, R., Li, Z., and Stocks, B.: Observations of boreal forest fire smoke in the stratosphere by POMA III, SAGE II and lidar in 1995. Geophys. Res. Lett., 27(9), 1407-1410, 2000.
- Füllekrug, M., Diver, D., Pinçon, J.-L., Phelps, A.D.R., Bourdon, A., Helling, C., Blanc, E., Honary, F., Harrison, R.G., Sauvaud, J.-A., Renaard, J.-B., Lester, M., Rycroft, M., Kosch, M., Horne, R.B., Soula, S., and Gaffet, S.: Energetic Charged
 Particles Above Thunderclouds. Surv Geophys, 34(1), 1-41, doi:10.1007/s10712-012-9205-z, 2013.
- Ghosh, K., Tripathi, S.N., Joshi, M., Mayya, Y.S., Khan, A., and Sapra, B.K.: Modeling studies on coagulation of charged particles and comparison with experiments. J. Aerosol Sci., 105, doi:10.1016/j.jaerosci.2016.11.019., 2017.

Gilbert, J.S., Lane, S.J., Sparks, R.S.J., and Koyaguchi, T.: Charge measurements on particle fallout from a volcanic plume. Nature, 349, 598-600, doi:10.1038/349598a0, 1991.

20 Hanson, D. R., Ravishankara, A.R., and Solomon, S.: Heterogeneous reactions in sulphuric acid aerosols: a framework for model calculation. J. Geophys. Res., 99, 3615-3629, doi:10.1029/93JD02932, 1994

Harrison, R.G., and Carslaw, K.S.: Ion-aerosol-cloud processes in the lower atmosphere. Rev. Geophys, 41(3), 2-1 2-26, doi:101029/2002RG000114. 2003.

- Harrison, R. G., Nicoll, K. A., Ulanowski, Z., and Mather, T. A.: Self-charging of the Eyjafjallajökull volcanic ash plume.
 Environ. Res. Lett., 5, 024004, doi:10.1088/1748-9326/5/2/024004., 2010.
 - Harrison, R.G., Nicoll, K.A., and Aplin, K.L.: Vertical profile measurements of lower troposphere ionization. J. Atmos Sol-Terr. Phys., 119, 203-210, doi:10.1016/j.jastp.2014.08.006., 2014.
 - Hervig, M., and Deshler, T.: Evaluation of aerosol measurements from SAGE II, HALOE and balloonborne optical particle counters. J. Geophys. Res., 107, D3, 4031, doi:10.1029/2001JD000703, 2002.

- Hervig, M. E., Gordley, L. L., Deaver, L. E., Siskind, D. E., Stevens, M. H., Russell, III J. M., Bailey, S. M., Megner, L., and Bardeen, C. G.: First satellite observation of meteoric smoke in the middle atmosphere. Geophys. Res. Lett., 36, L18805, doi:10.1029/2009GL039737, 2009.
- Hirsikko, A., Nieminen, T., Gagné, S., Lehtipalo, K., Manninen, H.E., Ehn, M., Hörrak, U., Kerminen, V.-M., Laakso, L.,
 McHurry, P.H., Mirme, S., Petäjä, T., Tammet, H., Vakkari, V., Vana, M., and Kulmala, M.: Atmospheric ions and nucleation: a review of observations. Atmos. Chem. Phys., 11, 767-798, doi:10.5194/acp-11-767-2011, 2011.
 - Hoppel, W., and Frick, G.: Ion-aerosol attachment coefficients and the steady-state charge distribution on aerosols in a bipolar environment. J. Aerosol Sci., 5, 1-21, 1986.
- Hunton, D.E., Viggiano, A.A., Miller, T.M., Ballenthin, J.O., Reeves, J.M., Wilson, J.C., Lee, S.-H., Anderson, B.E., Brune,
 W.H., Harder, H., Simpas, J.B., and Oskarsson, N.,: In-situ aircraft observations of the 2000 Mt. Hekla volcanic cloud: composition and chemical evolution in the Arctic. J Volcanol Geotherm Res, 145, 23-24, doi:10.1016/j.volgeores. 2005.01.005, 2005.
 - IPCC (Intergovernmental Panel on Climate Change): Climate Change 2013, The Physical Science Basis. Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. (Stocker, T.F., Qin, D.,
- 15 Plattner, G.-K., Tignor, M.M.B., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P.M., eds.). Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, ISBN 978-1-107-05799-1, 1535 pp, 2013
 - Junge, C.E.: Stratospheric aerosols studies. J. Geophys. Res. 66(7), 2163-2182, 1961.

Junge, C.E., Chagnon, C.W., Manson, J.E.: Stratospheric aerosols. J Met 18, 81-108, 1961.

- 20 Kanawade, V., and Tripathi, S.N: Evidence for the role of ion-induced particle formation during an atmospheric nucleation event observed in Tropospheric Ozone Production about the Spring Equinox (TOPSE). J. Geophys. Res., 111, D02209, doi:10.1029/2005JD006366., 2006.
 - Kasai, T., Tsuchiya, M., Takami, K., Hayashi, M., and Iwasaka, Y.: Balloon borne optical particle counter for stratospheric observation. Review of Scientific Instruments, 74, 1082, doi:10.1063/1.1533791, 2003.
- 25 Kazil, J., Harrison, R.G., and Lovejoy, E.R.: Tropospheric new particle formation and the role of ions. Space Sci. Rev., 137, 241-255, doi:10.1007/s11214-008-9388-2, 2008.

Kirkby, J., Curtius, J., Almeida, J., Dunne, E., Duplissy, J., Ehrhart, S., Franchin, A., et al.: Role of sulphuric acid, ammonia and galactic cosmic rays in atmospheric aerosol nucleation. Nature, 476(7361), 429-433, doi:10.1038/nature10343, 2011.

Kirkby, J., Duplissy, J., Sengupta, K., Frege, C., Gordon, H., et al.: Ion-induced nucleation of pure biogenic particles.
Nature, 533, 521-526, doi:10.1038/nature17953, 2016.

Kremser, S., Thomason, LW., von Hobe, M., Hermann, M., Deshler, T., et al.-: Stratospheric aerosol-Observations, processes, and impact on climate. Rev Geophys 54, 278-335, doi:10.1002/2015RG00511, 2016.

Laakso, L., Grönholm, T., Kulmala, L., Haapanala, S., Hirsikko, A., et al.: Hot-air balloon as a platform for boundary layer profile measurements during particle formation. Boreal Env. Res., 12, 279-294, 2007.

- 5 Lary, D. J., Shallcross, D. E., and Toumo, R.: Carbonaceous aerosols and their potential role in atmospheric chemistry. J. Geophys. Res., 104, 15,929-15,940, doi:10.1029/1998JD1000091, 1999.
 - Launder, B., and Thompson, J.M.T.: Geo-engineering climate change: environmental necessity or Pandora's box? Cambridge University Press, Cambridge, UK, ISBN 0961408812, 332 pp, 2009
- Li, Y., Guo, X., Wang, T., Zhao, Y., Zhang, H., and Wang, W.: Characteristics of atmospheric small ions and their
 application to assessment of air quality in a typical semi-arid city of Northwest China. Aerosol Air Quality Res, 15, 865-874, doi:10.4209/aaqr.2014.06.0123, 2015.

Matsumura, T., Hayashi, M., and Fujiwara, M.: Observations of stratospheric aerosols by balloon-borne optical particle counter at Bandung, Indonesia. Journal of the Meteorological Society of Japan, 2, 709-718, 2001.

- Michael, M., Barani, M., Tripathi, S.N.: Numerical predictions of aerosol charging and electrical conductivity of the lower 15 atmosphere of Mars. Geophys. Res. Lett., 34, L04201, doi:10.1029/2006GL028434, 2007
- Michael, M., Tripathi, S.N., and Mishra, S.K.: Dust charging and electical conductivity in the day and nighttime atmosphere of Mars. J. Geophys. Res., 113, E07010, doi:10.1029/2007JE003047, 2008

Murphy, D.M., Thomson, D.S., and Mahoney, M.J.: In situ measurements of organics, meteoritic material, mercury, and other elements in aerosols at 5 to 19 kilometres. Science, 282, 1664-1669, doi:10.1126/science.282.5394.1664, 1998.

20 Murphy, D.M., Froyd, K.D., Schwarz, J.P., and Wilson, J.C.: Observations of the chemical composition of stratospheric aerosol particles. Q. J. Roy. Meteor. Soc., 140, 681, 1269-1278, doi:10.1002/qj.2213, 2014

Myhre, G., Shindell, D., Bréon, F.-M., Collins, W., Fuglestvedt, J., Huang, J., et al.: Anthropogenic and natural radiative forcing. Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. [Stocker T.F., et al. (eds.)] Cambridge University Press, Cambridge,

- 25 United Kingdom, and New York, USA, pp. 659-740, 2013.
 - Neely, R. English, R., J. M., Toon, O. B., Solomon, S., Mills, M., and Thayer, J. P.: Implications of extinction due to meteoritic smoke in the upper stratosphere. Geophys. Res. Lett., 38, L24808, doi:10.1029/2011GL049865, 2011.

Nicoll, K. A., Harrison, R. G., and Ulanoswki, Z.: Observations of Saharan dust layer electrification. Environ. Res. Lett., 6, 014001, doi:10.1088/1748-9326/6/1/014001, 2011.

- O'Brien, K.: The theory of cosmic-ray and high-energy solar-particle transport in the atmosphere, in: The Natural Radiation Environment VII, Seventh International Symposium on the Natural Radiation Environment (NRE-VII), edited by: McLaughlin, J.P., Simmonds, S.E., and Steinhäusler, F., Elsevier, Amsterdam, The Netherlands, 29-44, 2005.
- Pawar, S.D., Meena, G.S., and Jadhav, D.B.: Air ion variation at poultry-farm, coastal, mountain, rural and urban sites in 5 India. Aerosol Air Quality Res., 12, 444-455, doi:10.4209/aaqr.2011.09.0148, 2012
 - Pierce, J.R., and Adams, P.J.: Can cosmic rays affect cloud condensation nuclei by altering new particle formation rates? Geophys. Res. Lett., 36, L09820, doi:10.1029/2009GL037946, 2009
- Raes, F., and Janssens, A.: Ion-induced aerosol formation in a H2O2-H2SO4 system-I. Extension of the classical theory and search for experimental evidence. J. Aerosol Sci., 16, 217-227, 1985.
- 10 Rapp, M.: Charging of mesospheric aerosols particles: the role of photodetachment and photoionization from meteoric smoke and ice particles. Ann. Geophys., 27, 2417-2422, doi:10.1016/j.jastp.2011.01.008, 2009.
- Rawal, A., Tripathi, S.N., Michael, M., Srivastava, A. K., and Harrison, R. G.: Quantifying the importance of galactic cosmic rays in cloud microphysical processes. J. Atmos. Sol-Terr. Phys., 102, 243-251, doi:10.1016/j.jastp.2013.05.017, 2013.
- 15 Regener, E, and Pfotzer, G.: Vertical intensity of cosmic rays by threefold coincidences in the stratosphere. Nature, 136, 718-719, 1935.
- Renard, J.-B., Ovarlez, J., Berthet, G., Fusson, D., Vanhellemont, F., Brogniez, C., Hadamcik, E., Chartier, M., and Ovarlez, H.: Optical and physical properties of stratospheric aerosols from balloon measurements in the visible and near-infrared domains. III. Presence of aerosols in the middle stratosphere. Appl. Optics, 44, 19, 4086-4095, doi:10.1364/AO.44.004086, 20 2005.
 - Renard, J.-B., Brogniez, C., Berthet, G., Bourgeois, Q., Gaubicher, B., Chartier, M., et al., 2008: Vertical distribution of the different types of aerosols in the stratosphere, Detection of solid particles and analysis of their spatial variability. J. Geophys. Res., 113, D21303, doi:10.1029/2008JD010150.
- Renard, J.-B., and coauthors, 2013: In situ detection of electrified aerosols in the upper troposphere and stratosphere. Atmos.
 Chem. Phys., 13, 1-8, doi:10.5194/acp-13-11187-2013.
 - Renard, J.-B., Tripathi, S.N., Michael, M., Rawal, M., Berthet, G., Fullekrug, M., Harrison, R.G., Robert, C., Tagger, M., and Gaubicher, B.: LOAC: a small aerosol optical counter/sizer for ground-based and balloon measurements of the size distribution and nature of atmospheric particles – Part 1: Principle of measurements and instrument evaluation. Atmos. Meas. Tech., 9, 1721-1742, doi: 10.5194/amt-9-1721-2016, 2016a.

- Renard, J.-B., Dulac, F., Berthet, G., Lurton, T., Vignelles, D., et al.: LOAC: a small aerosol optical counter/sizer for ground-based and balloon measurements of the size distribution and nature of atmospheric particles Part 2: First results from balloon and unmanned aerial vehicle flights. Atmos. Meas. Tech., 9, 3673-3686, doi:10.5194/amt-9-3673-2016Robock, A., 2000: Volcanic eruptions and climate. Rev. Geophys., 38, 2, 191-219, 2016b.
- 5 Russell, P.B., Livingston, J.M., Pueschel, R.F., Bauman, J.J., Pollack, J.B., et al.: Global to microscale evolution of the Pinatubo volcanic aerosol derived from diverse measurements and analyses. J. Geophys. Res., 101, D13, 18745-18763, doi:10.1029/96JD01162, 1996,
- Schwarz, J. P., Gao, R. S., Fahey, D. W., Thomson, D. S., Watts, L. A., et al.: Coatings and their enhancement of black carbon light absorption in the tropical atmosphere. J. Geophys. Res., 113, D03203, doi:10.1029/2007JD009042, 2008
- 10 Shiraishi, K., Hayashi, M., Fujiwara, M., Shibata, T., Watanabe, M., Iwasaka, Y., Neuber, R., and Yamanouchi, T.: Comparative analysis of measurements of stratospheric aerosol by lidar and aerosol sonde above Ny-Ålesund in the winter of 1995 [Comparative analysis of lidar and OPC observations]. Polar Sci, 5, 399-410, doi:10.1016/j.polar.2011.08.003, 2011.
- Steele, H. M., Lumpe, J. D., Turco, R. P., Bevilacqua, R. M., and Massie, S. T.: Retrieval of aerosol surface area and volume densities from extinction measurements: application to POAM II and SAGE II. J. Geophys. Res, 104, 9325-9336, doi:10.1029/1999JD900032, 1999.
- Sugita, T., Kondo, Y., Koike, M., Kanada, M., Toriyama, N., and Nakajima, H.: Balloon-borne optical counter for in situ aerosol measurements. J. Atmos. Chem., 32, 183-204, 1999.
- Suni, T., Kulmala, M., Hirsikko, A., Bergman, T., Laakso, L., et al.,: Formation characteristics of ions and charged aerosol particles in a native Australian Eucalypt forest. Atmos. Chem. Phys., 8, 129-139, doi:10.5194/acp-8-129-2008, 2008.
- 20 Svensmark, H., and Friis-Christensen, E.: Variation of cosmic ray flux and global cloud coverage-a missing link in solarclimate relationships. J. Atmos. Sol.-Terr. Phys., 59(11), 1225-1232, doi:10.1016/S1364-6826(97)00001-1, 1997
 - Svensmark, H., Pedersen, J.O., Marsh, N.D., Enghoff, M.B., and Uggerhøj, U.I.: Experimental evidence for the role of ions in particle nucleation under atmospheric conditions. P. Roy. Soc. A-Math. Phys., 463, 385-396, doi:10.1098/rspa.2006.1773, 2007
- 25 Tinsley, B., Rohrbaugh, R., Hei, M., and Beard, K.: Effects of image charges on the scavenging of aerosol particles by cloud droplets and on droplet charging and possible ice nucleation processes. J. Atmos. Sci., 57, 2118-2134, doi:10.1175/1520-0469(2000)057, 2000.
 - Tinsley, B.A., and Zhou, L.: Initial results of a global circuit model with variable stratospheric and tropospheric aerosols. J. Geophys. Res. D Atmos., 111, D16205, doi:10.1029/2005JD006988., 2006

Tinsley, B.A.: The global atmospheric electric circuit and its effects on cloud microphysics. Rep. Prog. Phys., 71(6): 066801, doi:10.1088/0034-4885/71/6/066801, 2008.

Tomikawa, Y., Sato, K., Hirasawa, N., Tsutsumi, M., and Nakamura, T.: Balloon-borne observations of lower stratospheric water vapor at Syowa Station, Antarctica in 2013. Polar Sci., 9, 4, 345-353, doi:10.1016/j.polar.2015.08.003, 2015.

5 Tositti, L., Brattich, E., Cinelli, G., and Baldacci, D.: 12 years of ⁷Be and ²¹⁰Pb in Mt. Cimone, and their correlation with meteorological parameters. Atmos. Environ., 87, 108-122, doi:10.1016/j.atmosenv.2014.01.014, 2014.

Tripathi, S.N., Michael, M., and Harrison, R.G.: Profiles of ion and aerosol interactions in planetary atmospheres. Space Sci. Rev., 137(1-4), 193-211, doi:10.1007/s11214-008-9367-7, 2008.

Vernier, J.-P., Thomason, L. W., Pommereau, J.-P., Bourassa, A., Pelon, J., Garnier, A., Hauchecorne, A., Blanot, L., Trepte,

10 C., Degenstein, D., and Vargas, F.: Major influence of tropical volcanic eruptions on the stratosphere aerosol layer during the last decade. Geophys. Res. Lett., 38, L12807, doi:10.1029/2011GL047563. 2011.

Watanabe, M., Iwasaka, Y., Shibata, T., Hayashi, M., Fujiwara, M., and Neuber, R.: The evolution of Pinatubo aerosols in the Arctic stratosphere during 1994-2000. Atmos. Res., 69, 199-215, doi:10.1016/j.atmosres.2003.09.006, 2004.

Williams, E., and Mareev, E.: Recent progress on the global electric circuit. Atmos. Res., 135-136, 208-227, 15 doi:10.1016/j.atmosres.2013.05.015, 2014.

Yu, F., and Turco, R.P.: Ultrafine aerosol formation via ion-mediated nucleation. Geophys. Res. Lett., 27, 6, 883-886, doi:10.1029/1999GL011151, 2000.

Yu, F. and Turco, R.P.: From molecular clusters to nanoparticles: The role of ambient ionization in tropospheric aerosol formation. J. Geophys. Res., 106, 4797-4814, doi:10.1029/2000JD900539, 2001.

20 Yu, F., Wang, Z., Luo, G., Turco, R.: Ion-mediated nucleation as an important global source of tropospheric aerosols. Atmos. Chem. Phys., 8, 2537-2554, doi:10.5194/acp-8-2537-2008, 2008.

Yu F.: Ion-mediated nucleation in the atmosphere: Key controlling parameters, implications, and look-up table. J. Geophys Res, 115, D03206, doi:10.1029/2009JD012630, 2010.

Yu, P., Rosenlof, K.H., Liu, S., Telq, H., Thornberry, T.D., et al.: Efficient transport of tropospheric aerosol into the stratosphere via the Asian summer monsoon anticyclone. Proc Natl Acad Sci USA, 114, 27, 6972-6977, doi:10.1073/pnas.1701170114, 2017.



Figure 1: Vertical profiles of ambient external temperature (Temp) and relative humidity (RH) as measured along the BEXUS18 stratospheric flight on 10 October 2014 and by the radiosounding from the Kandalashka station (67.15N, 32.35E) on 11 October 2014 at 00UTC.





Figure 2: Vertical profiles of integrated aerosols concentration, for aerosols greater than 200 nm (Black line), and of positive (red line) and negative ions (blue line).





Figure 3: Vertical profiles of particles size distributions for the 19 size classes_of the LOAC particle counter as part of the A5-Unibo experiment on the BEXUS18 stratospheric flight. The notation dN/dlog(D) used in the x-axis stands for the number concentration of particles in the various size classes divided by the width of the size classes





Figure 4: Average size distribution of aerosol particles as a function of a various height lavers height—during BEXUS18 stratospheric flight. The five curves are obtained averaging the aerosol number densities as a function of the atmospheric layers pointed out by the temperature profile as follows: 1: 0-404 m (black line); 2: 650-1519 m (red line); 3: 1765-10118 m (blue line); 4: 10650-25044 m (pink line); 5: 25289-27191 m (green line). X- and Y-axes are in log-normal scales.





Figure 5: Vertical profile of the simulated fraction of charged particles: a) total; b) in the different size classes. Because of the very low and even zero concertation particles in the largest size-classes, only particles with diameter smaller than 5 µm are considered.

	Positive Ions (#/cm ⁻³)	Negative ions (#/cm ⁻³)
Positive Ions (#/cm ⁻³)	1.00	
Negative ions (#/cm ⁻³)	-0.28	1.00
0.2-0.3 <u>(#/cm⁻³)</u>	0.44*	-0.67*
0.3-0.4 <u>(#/cm⁻³)</u>	0.45*	-0.75*
0.4-0.5 <u>(#/cm⁻³)</u>	0.40*	-0.71*
0.5-0.6 <u>(#/cm⁻³)</u>	0.33*	-0.80*
0.6-0.7 <u>(#/cm⁻³)</u>	0.34*	-0.62*
0.7-0.9 <u>(#/cm⁻³)</u>	0.28	-0.76*
0.9-1.1 <u>(#/cm⁻³)</u>	0.01	-0.62*
1.1-3.0 <u>(#/cm⁻³)</u>	0.31*	-0.22
3.0-5.0 <u>(#/cm⁻³)</u>	0.15	-0.07
5.0-7.5 <u>(#/cm⁻³)</u>	0.12	-0.28*
7.5-10.0 <u>(#/cm⁻³)</u>	0.33*	0.23
10.0-12.5 <u>(#/cm⁻³)</u>	0.17	-0.10
12.5-15. <u>0 (#/cm⁻³)</u>	0.36*	
15.0-17.5 <u>(#/cm⁻³)</u>	0.36*	0.27*
17.5-20.0 <u>(#/cm⁻³)</u>	0.36*	
20.0-22.0 <u>(#/cm⁻³)</u>	0.36*	
22.0-30.0 <u>(#/cm⁻³)</u>	0.32*	-0.18
30.0-40.0 <u>(#/cm⁻³)</u>	0.24	
40.0-50.0 <u>(#/cm⁻³)</u>		0.18
Aerosols > 200 nm_(<u>#/cm⁻³)</u>	0.45*	-0.59*

 Table 1: Spearman's correlation coefficient between ions and number of particles in the different size classes. The asterisk * indicates significant (0.05 significance level, i.e., p<0.05) correlation coefficients; bold indicates strong (R > 0.6 in absolute value)

 5
 correlation, while italics indicates weaker (0.4 < R < 0.6 in absolute values)...</td>