

Interactive  
comment

# ***Interactive comment on “Localized TEC enhancements in the Southern Hemisphere” by Ilya K. Edemskiy***

**Ilya Edemskiy**

ilya@iszf.irk.ru

Received and published: 21 December 2019

1. Authors mentioned (lines 20-22) that: “The Southern Hemisphere contains at least two large anomalous regions: South Atlantic Magnetic Anomaly and Weddell Sea Anomaly. The latter consists in the modulation of TEC’s diurnal oscillations by the solar-modulated seasonal oscillations, which produces a diurnal anomaly in the Discussion paper vicinity of the Weddell Sea during Southern Hemisphere summer (October to March) (Lean et al., 2016).” I recommend author to take the traditional (more clear) definition of WSA phenomenon.

—  
This part of introduction was changed: “It is known that Southern Hemisphere contains



some anomalous regions. South Atlantic Magnetic Anomaly (SAMA) is formed by a configuration of geomagnetic field which has a global minimum of intensity over South Atlantic and South America and makes it easier for energetic particles of inner radiation belt to precipitate, thus increasing ionospheric conductivity over the region (Abdu et al., 2005). South of the SAMA, in the south-eastern Pacific and South Atlantic Antarctic regions, combination of the geomagnetic field features and thermospheric winds produces an inverted diurnal plasma density pattern at equinoxes and in SH summer (October-March): the nighttime maximum is larger than the daytime minimum, and the phenomenon is known as the Weddell Sea Anomaly (WSA) (Horvath, 2006). Jakowski et al. (2015) showed that during periods of low solar activity in Asian longitudinal sector of SH it is possible to observe so called nighttime winter anomaly (NWA), when values of electron concentration are higher in winter than in summer. At the same time Yasyukevich et al. (2018) showed that winter anomaly manifests itself much less intensively in SH than in NH. It is possible to conclude that ionosphere of each hemisphere has some specific features.”

2. (Lines 34-35): “During analysis of ionosphere response to a geomagnetic storm of 15 August 2015, a curious structure was detected in global ionospheric maps (GIMs), which we call localized TEC enhancement or LTE (Edemskiy et al., 2018).” The term localized TEC enhancement was mentioned many years before by Foster and Rideout (2007) and Foster and Coster (2007). Note that Foster et al. studies present localized TEC enhancement in Northern hemisphere many times. I recommend author to read John Foster’s et al. articles in order to understand the morphology and physical explanation of localized TEC enhancement in NH. I believe that these papers should give you new information.

—  
Thank you for your recommendation. The text was changed and reference to Foster et al. papers was added. At the same time it should be noted that their papers mostly describe ionosphere during geomagnetic storms, investigating manifestation of storm

enhanced density (SED), whereas the considered LTEs are observed almost independently on geomagnetic conditions, even during quiet periods ( $K_p=1$ ). The introduction was changed and the following was added: “The most typical irregularities in distribution of electron concentration are produced during geomagnetic storms. Foster and Coster (2007) investigating storm enhanced densities (SED). They showed that during severe and extreme storms it is possible to detect SEDs which in maps of total electron content (TEC) could be observed as localized TEC enhancements (LTE). The authors showed that during a storm recovery phase LTEs could be detected in the night side ionosphere at the middle latitudes of both hemispheres, in magneto-conjugated regions. The authors note that the observed enhancements are approximately corotating in place over the positions in which they were formed earlier in the event. However, the LTE phenomenon studied by Foster and Coster (2007) is different from the LTE phenomenon studied by us. During analysis of ionospheric response to a geomagnetic storm of 15 August 2015 Edemskiy et al. (2018) detected a curious LTE in global ionospheric maps (GIMs). Unlike the LTEs observed by Foster and Coster (2007) this enhancement was detected and observed in sunlit area of the Southern Hemisphere and lasted for several hours. It was not corotating but changing position following the Sun and propagating along the geomagnetic parallels. Using quite a simple detection algorithm Edemskiy et al. (2018) found about 30 similar events in the Southern Hemisphere during 2010-2016 and some of the detected LTEs were observed during relatively quiet periods. The authors showed direct dependence of number of the detected LTEs on solar activity level and suggested that their generation is connected with the orientation of interplanetary magnetic field (IMF), namely with  $B_z$ .”

3. (Lines 45-55). Unfortunately there are many remarks about definition of LTE's. According to first sentence here “The localized TEC enhancement is a positive disturbance of ionosphere.” But according to two detection criteria the LTE is a spatial-temporal structure in the UT map of TEC and is not a disturbance.

[Printer-friendly version](#)

[Discussion paper](#)



Yes, thank you, you are absolutely right. Now it is noted in the text that LTE should not be considered as a disturbance. The criteria were changed as well: "In this paper a TEC enhancement is considered as LTE if it is: - located in middle latitudes of sunlit region. Mainly we investigate LTEs, which are clearly observed in Indian and Southern part of Atlantic Oceans and do not take into account enhancements in Northern Hemisphere. At the same time, LTEs in SH are not accompanied by any LTE in NH and such a focusing on SH LTEs is quite reasonable. - spatially limited by relatively lower TEC values. Normalized difference between squared maximal value in LTE and minimal one at its border ( $\Delta=1 - (l_{edge}/l_{max})^2$ ) should be no less than 20%. Generally that means that there should be a clear trough between an enhancement and the equatorial ionization anomaly (EIA). - confined and have a border of lower TEC values ( $\Delta \geq 20\%$ ) no farther than in  $40^\circ$  in longitude from the location of maximal TEC value. Mainly that means that we do not consider longitudinally stretched enhancements assuming different mechanism of their generation."

4. (Lines 47-48): "1. Spatial limitation and clear borders. An enhancement should not be wider than  $40^\circ$  and  $120^\circ$  in latitude and longitude, respectively. Gradients at an LTE edges should be high enough to make LTE borders possible to distinguish." According to such limitation almost all of winter UT map of TEC should reveal LTE due to 1) short duration (therefore limitation in longitude smaller then  $90^\circ$ ) and significant gradients of TEC diurnal variation during sunlight hours; 2) clear border at sub-auroral latitudes due to pronounce main ionospheric trough structure (for daytime also). So according to these criteria, I don't understand how LTE can be distinguished from the usual TEC maps in winter and equinox seasons. Figure 3 demonstrate many cases of consistence between LTE and typical TEC diurnal variation (that is presented in view of longitude-latitude map for UT epoch). Another problem is statement: "Gradients at an LTE edges should be high enough". Please provide mathematical formulation for "should be high enough". Or this criteria was checked manually for each maps?"

[Printer-friendly version](#)[Discussion paper](#)

Thank you for this remark. The criteria definition was changed to describe an LTE more precisely. Key point which was missed previously is a presence of trough between EIA and the observed enhancement. Demands on the minimal depth of this trough is given mathematically. All the enhancements in fig. 3 fulfill the criteria and are considered as LTE. Several examples of maps showing other types of enhancement or absence of enhancement are available via the link below. Such situations were considered as non-LTE cases and were not considered during the investigation. [https://drive.google.com/drive/folders/1u6GTYRe9bIFb-Kb25gMKkGM5\\_L1vbQAv?usp=sharing](https://drive.google.com/drive/folders/1u6GTYRe9bIFb-Kb25gMKkGM5_L1vbQAv?usp=sharing)

5. (Lines 52-54): "We search LTEs only in the Southern Hemisphere, because Edemskiy et al. (2018) detected LTEs only at SH. A disturbance should follow the Sun having the maximal intensity no latter than 1-2 hours after local noon (in a period 12-14 LT as observed by Edemskiy et al., 2017). I disagree with argument for LTE limitation only in the Southern Hemisphere. Edemskiy et al. (2018) study concern to geomagnetic storm response on particular event on Aug 2017. There are many examples of daytime storm-time localized TEC enhancement (Foster and Rideout, 2007; Zhao et al., 2012) in NH. Why author's algorithm exclude all these situations? I did not found Edemskiy et al., 2017 in the reference list.

—

Misprint "Edemskiy et al., 2017" is corrected: "Edemskiy et al., 2018" The statement was badly formulated. The point was that this particular paper is dedicated only to LTEs detected in Southern Hemisphere, particularly in South Indian and South Atlantic regions. The same LTE was described by Edemskiy et al. (2018) and the idea of the current paper is to find other similar enhancements using GIMs. During the investigation it was found that these structures can be detected not only during magnetic storms but during quiet days as well. The author does not claim absence of LTEs in Northern Hemisphere or in other ranges of longitude. However during SH LTE we do not see any corresponding effect in NH. At the same time the aim was not to describe only storm-

Printer-friendly version

Discussion paper



time LTE, but any such formation which are observed only in Southern Hemisphere. This is reflected in new formulation of the criteria.

6. (Lines 89-92): "An example of a clearly observed LTE was detected at April 5, 2014 (Fig. 1). The disturbance reached the highest intensity in a period 10-12 UT when TEC values in a center of the disturbance exceeded 78 TECU. This value is comparable to equatorial TEC values. The highest values were detected in a latitudinal region 45-70 °S. At the same time, TEC values of the entire region (30-70 °S, 0-90 °E) were enhanced." The reason for SLTE is associated to geomagnetic disturbances during 5 April. Please see AE index on Fig. 1. It is evident that geomagnetic disturbances in AE started at 06 UT on 5 April, 2014 (the same as SLTE). The maximal geomagnetic disturbance occur at 10-12 UT. At the same time the highest intensity of SLTE occur when TEC values in a center of the disturbance exceeded 78 TECU. So SLTE in reality can be SED structure or something else that associated with geomagnetic disturbances.

—

Thank you very much for this remark. It moved the investigation forward. SED structures are mentioned both in introduction and discussion sections and briefly described in the latter. The possible connection between SED and LTE is discussed and some statistical analyses are added as well. At the same time it is shown that clear SLTE are observed during relatively quiet conditions with positive Dst and small AE values; and otherwise: not all storms were accompanied by SLTEs.

7. (Lines 94-95): "As it will be shown later, such a strong SLTE is not typical and in some cases it is not detected at all." Why author to select this case if this case is not typical?

—

This day was chosen since despite the more intense SLTE both the enhancements are

[Printer-friendly version](#)

[Discussion paper](#)



observed quite clearly. This structure fulfills the criteria of LTE and at least partially can be clearly seen in SWARM data. Another map with a LTE confirmed by SWARM is added to the discussion.

8. (Lines 106-107): "In-situ measurement of electron concentration Ne from SWARM satellites allow us to check validity of TEC distribution presented by GIM." In my own opinion Fig. 2 provide clear evidence of SLTE, but not for MLTE. So according to my points 6-8 Figs. 1 and 2 does not give to reader typical examples of LTE. I recommend to add a typical example of MLTE that does not associated with geomagnetic disturbances.

—

Thank you for this remark. Another figure containing MLTE and corresponding SWARM measurements are added to the paper. Some text clarifying this problem was added to the discussion section.

9. Figure 5. IMF intensity is not a good choice of parameter that determine geo-magnetic activity because direction of IMF Bz can be more important for ionospheric disturbances. In my opinion AE or AP index can be more effective in this investigation.

—

Analysis of LTE occurrence rate dependence on geomagnetic indices or IMF components did not reveal any trends that is why these distributions were not shown in figures. The dependence on Bz was the initial hypothesis which was not supported by observations. The fig. 5 is changed now and contains all the basic geomagnetic indices.

10. About discussion part. It is very chaotic. I still did not understand which of the mechanisms, according to the author, is the main one for the formation of LTE. There are a lot of additional questions according to LTE, but I stopped here in order to obtain some clarification about LTE.

[Printer-friendly version](#)

[Discussion paper](#)



The discussion part was almost fully rewritten and now contains more details and suggestions about LTEs.

---

Interactive comment on Ann. Geophys. Discuss., <https://doi.org/10.5194/angeo-2019-124>, 2019.

Interactive  
comment

[Printer-friendly version](#)

[Discussion paper](#)

