

Interactive comment on “Earth’s radiation belts ions: Patterns of the spatial-energy structure and its solar-cyclic variations” *by*
Alexander S. Kovtyukh

Anonymous Referee #1

Received and published: 29 November 2019

article [utf8]inputenc longtable

Comments for the manuscript: “Earth’s radiation belts ions: Patterns of the spatial-energy structure and its solar-cyclic variations” by A. S. Kovtyukh

Thank you very much or the corrections! I think that the manuscript now is much linear
C1

[Printer-friendly version](#)

[Discussion paper](#)



and easy to read. I have just a few more corrections though, mostly concerning the fluency of the text and no more technical questions. The list is reported below.

l p12cm

Line Comment

12 "[...] in the inner regions of the ERB, fluxes [...]"

24-25 "The ERB consist mainly of electrons and protons, but there are also helium nuclei and other [...]"

33 "[...] geomagnetic trap, drift conserving [...] and populate [...]"

34-35 "This layer is called the drift shell. "

37 "For the dipole magnetic field, L is [...]"

42 "[...] along a certain magnetic field line [...]"

43-44 "This dependence is described [...]"

45 "[...] the same magnetic field line, respectively [...]"

48-49 "[...] of radial diffusion of ions towards [...]"

53 "The inner belt ($L < 2.5$) of protons with $E > 10$ MeV is formed by [...]"

55 "For protons with $E < 10$ MeV, this mechanism [...]"

56 "The inner belt of ions with $Z > 4$ is formed [...]"

58-59 "In the intermediate region ($2.5 < L < 3.5$), the mechanism of a ion capture from the Solar Cosmic Rays takes place during strong magnetic storms [...]"

62 "However, for a comprehensive verification [...]"

76-77 "[...] the possibility to create sufficiently complete and reliable empirical models [...]"

79-83 "In the following sections, the spatial-energy structure of the ERB in the $\{E, L\}$ space for protons, helium and CNO group ions are considered (Sect. 2), together with possible physical mechanisms of formation of these structures and their solar-cyclic variations (Sect. 3). Finally, the main conclusions of this work are given (Sect. 4). "

89 "According to this criterion and to the theory of [...]"

91 “[...] represents this very boundary [...] ”

93-94 “A significant number of these discrepancies can be connected to the [...] ”

107-108 “[...] to separate fluxes of ions by their charge. Moreover, for the ions [...] ”

114-115 “[...] the results of every experiment can be compared to the others [...] ”

119 “Figures 1–6 show the spatial-energy distributions [...] ”

120 I suggest removing entirely the quote “hese figures united in pairs: ” and just leave the part describing odd and even Figures

123-124 “The markers are connected by lines of equal intensity [...] ”

132 I suggest removing the quote “In this place, it is need to say a few words about the method of constructing these figures.[...] ”

147 “[...] corresponding set of experimental points (icons); then it was transferred [...] ”

185 “Figure N sums up results from [...] ”

188-189 “ $21^{st} / 22^{nd} / 23^{rd}$ [...] ”

190 See line 185

198 See lines 188-189

205 “From a comparison of Figs. 1 and 2, one can see [...] ”

210 “[...] (2016a,b), which have been constructed from Figs. 1 and 2 confirm [...] ”

214 “[...] $J \propto E^{-\gamma}$, where the index $\gamma =$ [...] ”

219 I would remove the “of the magnetic field” part here, magnetic field lines already describe everything

223 “Segments of iso-lines, that are parallel to the red line, also correspond to [...] ”

227 “at $L = 3-6$, $\gamma = 4.8 \pm 0.5$. [...] ”

228 “between these iso-lines increase with L [...] ”

239 “[...] helium ion fluxes, averaged for quiet periods ($Kp < 2$), are presented [...] ”

240 See line 185

244-245 See lines 188-189

246 See line 185

- 250 See lines 188-189
- 251 "with Figs. 3–4, one can see that [...]"
- 253-254 "[...] $E > 1$ MeV practically do not change, and [...]"
- 255 "Figures 3 and 4 show the same patterns [...]"
- 258 "[...] because there are no experimental data for helium ions in these regions. [...]"
- 266 "For helium spectra [...]"
- 270 "[...] the red line (i.e. in the region of power-law spectra) substantially deviate from [...]"
- 278 "[...] CNO group ions fluxes, averaged for quiet periods ($Kp < 2$), are [...]"
- 280 See line 185
- 283-284 See lines 188-189
- 285 See line 185
- 286-287 "[...] period of activity [...]"
- 289 "[...] nd its configuration differ [...]"
- 291 "[...] Figs. 5–6 one can see that, for ions of CNO group, the [...]"
- 297 "[...] This means that, for ions of the CNO group, the ionization [...]"
- 300 "[...] have not been obtained by the experiments collected in [...]"
- 304 "[...] especially large at the peak of solar activity (Fig. 6): during these times, the slope of iso-lines [...]"
- 306 "At the same time, at $L > 4$ in Fig. 5 and at $L > 3$ in Fig. 6, the iso-lines [...]"
- 316 "[...] at the minimum of solar activity [...]"
- 319 "[...] following the results obtained [...]"
- 324 "[...] and are reduced rapidly with [...]"
- 328 "[...] have not been considered in these works [...]"
- 329 "In quiet periods, only the mechanism of ionization loss is significant [...]"
- 330 "[...] trapped in small L [...]"

Printer-friendly version

Discussion paper



- 331 "[...] the ERB protons are determined, in this mechanism, by the density [...]"
- 336 "[...] the proton supply rates to the inner belt, under the action of the CRAND mechanism, remain [...]"
- 338 "[...] with decreasing solar activity [...]"
- 341 "A proton lifetime on [...]"
- 352 "[...] this was noted in sections [...]"
- 357 "[...] where L^* corresponds to the L shell of protons of the same energy [...]"
- 362 "[...] remains unchanged [...]"
- 363-364 "[...] in fact, these protons form mainly under the action [...]"
- 369 "[...] of radial diffusion of ions during the [...]"
- 377 "[...] values of L, these fluxes begin [...]"
- 384 "[...] highly turbulized region, but [...]"
- 387 "[...] must be generated in the outer [...]"
- 389 "The high-energy part of the ion [...]"
- 390 "[...] has a power-law shape and the exponents [...]"
- 394-395 "[...] along logarithmic axes E and J in a J(E) plane [...]"
- 424 "[...] decreases rapidly with [...]"
- 426 "Then, the lower boundary [...]"
- 429 "Using B_s [...]"
- 434-435 "[...] as a result of their interactions with the current layer [...]"
- 439 "It has been found that in the outer belt [...]"
- 438-439 "[...] in the near equatorial plane [...]"
- 443 "[...] radial diffusion which conserves μ [...]"
- 448 "This kind of dependence of the amplitude [...]"
- 461 "[...] the extensive gaps in $Z \geq 2$ ion data do not allow [...]"

Printer-friendly version

Discussion paper



640-644 For what concerns all the Figures, the captions are very similar, so I suggest a modification in the first one (Figure 1) that should be repeated for all the others: "[...] J, which is given in units of $(\text{cm}^2 \text{ s ster MeV})^{-1}$, is the differential flux of protons [...]" . Also "associated with " and "[...] the power-law tail of the proton spectra, while green line corresponds to [...]"

[Interactive comment](#)

[Printer-friendly version](#)

[Discussion paper](#)

