

**Referee #1** Review on “Quasi 10-day wave modulation of equatorial ionization anomaly during the Southern Hemisphere stratospheric warming of 2002” by Xiaohua Mo

In this work, the author has used the TEC data in the Southern Hemisphere (SH) to demonstrate the effects of quasi 10-day wave on the Northern and Southern TEC crests during the 2002 SH SSW event. The manuscript has been written well and is generally easy to understand. The results in this manuscript do provide a clear evidence of the quasi 10-day modulation of the TEC during the 2002 SSW event. I do have a few concerns and comments, which are mentioned below. However, in general, the manuscript provides some interesting new results and should be accepted after a revision.

**Answer:**

We very appreciate your substantial comments for our study about “Quasi 10-day wave modulation of equatorial ionization anomaly during the Southern Hemisphere stratospheric warming of 2002”. Here are our reply comments. All major revisions are marked in yellow highlights. Thank you very much.

=====

**Specific comments:**

.....

1. Please plot a figure showing the location of the stations used in this work.

**Answer:** The locations of the GPS stations shown in Figures 1 are added in the revised version.

.....

2. Line 97 - “To exclude these long period fluctuations in EIA region associated with solar/magnetosphere forcing, the periods longer than 15 days in the MLAT location and TEC of EIA crest are removed”. How is this process achieved? The author should clarify more regarding the applied method.

**Answer:** To remove the periods longer than 15 days in these parameters, these parameters are subtracted from their respective 15-day moving average.

**Line # [101-104]:** To exclude these long period fluctuations in EIA region associated with solar/magnetosphere forcing, the periods longer than 15 days in the MLAT location and TEC of EIA crest, and EEJ are removed. Specifically, these parameters are subtracted from their respective 15-day moving average.

.....

3. Line 152 - “Moreover, strong planetary wave scale quasi 10-day variation was observed in polar stratospheric temperature during this period”. Please cite the work in which this observation was mentioned.

**Answer:** Relevant references have been cited in the revised version.

**Line # [162-164]:** Moreover, strong planetary wave scale quasi 10-day variation was observed in polar stratospheric temperature during this period (Krüger et al., 2005; Palo et al., 2005), so the quasi 10-day oscillations in EIA region may be related to atmosphere perturbations linking the SSW in the Southern

Hemisphere.

**Technical corrections:**

- Figure 1 caption - Correct to solar flux
- Line 92 - it consists of an eastward-propagating
- Line 105 - too weak to be identified in F10.7
- Line 107 - evolution
- Line 108 - TEC of EIA crest and Kp,
- Line 124, 125 - band-pass
- Line 146 - have
- Line 155 - series of studies have showed

**Answer:** These grammatical and wording mistakes have been corrected in revised one.

**Referee #2** Review on “Quasi 10-day wave modulation of equatorial ionization anomaly during the Southern Hemisphere stratospheric warming of 2002” by Xiaohua Mo

This paper is focused on the possible influence of the quasi 10-day planetary waves (PWs), registered in the high-latitude polar stratospheric temperature before and around the Southern Hemisphere (SH) sudden stratospheric warming (SSW) in 2002, on the oscillations of the equatorial ionization anomaly (EIA) crests and their Total Electron Content (TEC). The locations of the EIA crests are calculated from the observations of the two GPS stations in China which are situated near the northern and southern EIA crests and the TEC data are derived from the International GNSS Service global ionospheric TEC maps in Asia. The SH SSW is described by the temperature and zonal mean zonal wind taken from the NCEP while the geomagnetic and solar variability are characterized by the Kp-index and solar radio flux F10.7 respectively. The period from July 21 to October 18, 2002 is considered (day numbers 200-300) and the quasi-10-day variability associated with the SSW is found in both the location of the EIA crests and the TEC between days 220-290. The author suggested that the observed ~10-day oscillation of the EIA region is generated through modulating the equatorial fountain effect.

The topic of the paper is certainly appropriate for the journal. In general, the paper is written clearly; actually it follows the pattern of the previous paper of the author, Mo et al. (2014) cited here. This study is certainly useful for the scientific community working on the vertical coupling of the atmosphere-ionosphere system, however I think that it has serious deficiencies. Due to this I will suggest the publication of this paper but after serious revision and addressing the comments mentioned below.

**Answer:**

We very appreciate your substantial comments for our study about “Quasi 10-day wave modulation of equatorial ionization anomaly during the Southern Hemisphere stratospheric warming of 2002”. Here

are our reply comments. All major revisions are marked in yellow highlights. Thank you very much.

---

---

**Major comments:**

.....  
(1) I have serious concern about the significance of the observed ~10-d oscillations particularly in the location of the EIA crests because the amplitude of these oscillations is only around 1.5° (Fig. 5a). Additionally, in Lomb-Scargle periodograms these oscillations are significant only above 90% confidence level (Fig. 3) that is not enough. It has been mentioned above that this study is similar to the previous one Mo et al. (2014). However, while there the quasi 16-day oscillations of the EIA crests were evident even in the raw data here the quasi-10-day ones are not. Usually only waves with significance at least above 95% confidence level are considered in studying the atmospheric and ionospheric perturbations. The author does not mention anything about the error in calculating the coordinates of the EIA crests. Without knowing the error in calculating the MLAT of the EIA crests it is difficult to accept the 10-day variability of the EIA region as significant one.

**Answer:** The perturbations amplitude in EIA region during SH SSW are smaller than those during NH SSW (Olson et al., 2013), so the Quasi 10-day wave in EIA region is not quite obvious. To make the results more credible, the 90% confidence level has been changed to 95% confidence level in Figures 4. It can be seen the quasi 10-day periodic component of most EIA parameters can reach 95% confidence level. Moreover, the significance levels of quasi 10-day oscillations revealed by the Morlet wavelet spectral analysis are higher than 95%. The error of the collected EIA crest latitude according to IPP trajectory is determined by the selected ionospheric shell height. Due to ionospheric variation, the shell height varies in different time and condition. Now, the constant shell height is used in almost all TEC derivation method, and the value of this height affects the resolution of the IPP location and the derived TEC values. So it is difficult to estimate the error of the EIA latitude. In the original manuscript, we only roughly estimate the error of the latitude from the sample rate of the raw GPS data according to the IPP velocity in the ionospheric shell.

**Line #[104-108]:** The residuals are subjected to Lomb-Scargle (L-S) spectral analysis (Lomb,1976; Scargle, 1982), and the results are shown in Figures 4a, 4b, 4c, 4d, and 4e. The horizontal dashed lines represent the 95% confidence level. It is evident that the MLAT location and TEC of EIA crest, and EEJ all exhibit significant quasi 10-day periodic component, which exceed or approach 95% confidence level, suggesting that the whole dynamical process in EIA region is modulated by quasi 10-day wave.

**Line # [69-74]:** Since the ionospheric vertical TEC usually reach the maximum at EIA crest, the location of EIA crest can be obtained by vertical TEC values at each ionospheric penetration point (IPP), which is the intersection of the line of sight and the ionospheric shell (assumed to be 400 km) (Mo et al., 2014). The relative accuracy of the TEC is 0.02 total electron content unit (1TECU= $10^{16}$  el m<sup>-2</sup>) (Hofmann-Wellenhof et al., 1992). The sample rate of these GPS stations were 30s, so the resolution of the location of EIA crest is less than 25 km (Mo et al., 2017).

(2) In order to propose a mechanism for generating the 10-day variability of the EIA region the authors used indirect approach based on some general references on dynamics as well as references connected with the SH SSW in 2002. The important citations as: Eswarajah et al. (2018) or Olson et al. (2013) which however present ground-based measurements at high latitudes or at Peruvian longitude sector cannot be considered as serious evidences because the author investigates different region, low latitudes over China. I cannot understand why the author does not use a meteor wind data from a Chinese radar at low latitudes and to check if there are quasi-10-day wave or modulated tides which are able to affect the fountain effect. Further, to see if the electric currents are modulated the author may consider the perturbations in the geomagnetic fields revealed from magnetometer measurements. Only then a solid evidence can be presented in support of the suggested mechanism.

**Answer:** The Meteor and MF radar was ever set up and was tested after 2003 in China, so the direct observations of MLT neutral wind are not available in China low latitude during 2002 SH SSW. The equatorial electrojet (EEJ) estimated by geomagnetic field cannot be obtained in China sector, because the geomagnetic field data in Bac Lieu in Vietnam are missing during the period from Aug 28, 2002 to Sep 26, 2002 (day number 240-269). In order to compensate this weakness, the EEJ estimated by geomagnetic field in Indian sector is added in revised manuscript. The EEJ in Indian sector also exhibit significant quasi 10-day periodic component.

**Line # [83-86]:** To demonstrate the dynamical process in EIA region, the EEJ is also used in this study, which can be estimated by the difference between the horizontal component of geomagnetic field at TIR (8.7°N, 77.8°E, MLAT~0.03°N) and VSK (17.68°N, 83.32°E, MLAT~8.56°N) (Rastogi et al., 1990). The results are shown in Figures 2e.

.....

(3) Important studies on atmospheric dynamics and the ionospheric response to the SSW events are not cited.

Concrete comments:

P. 2, lines 37-38: Please, add the following references: Chau et al. (GRL, 36, L05101, 2009, doi:10.1029/2008GL036785) giving evidence for the vertical plasma drift changes during the SSW and Pancheva and Mukhtarov (JASTP, 73, 1697–1702, 2011, doi:10.1016/j.jastp.2011.03.006) presenting which main characteristics of the EIA and how they are changed during the major SSW.

P. 2, line 38: Please, add Jin et al. (JGR, 117, A10323, doi:10.1029/2012JA017650) where for the first time a comparison between the results from a whole atmosphere ionosphere coupled model (GAIA) with the COSMIC and TIMED/SABER observations during the major SSW in January 2009 was presented.

**Answer:** These references have been added in the introduction part of revised manuscript.

**Line # [35-38]:** Although the main processes of SSW occur in the middle atmosphere, its effects on the ionosphere have been observed in significant changes of equatorial electrojet (EEJ), vertical plasma drift, and equatorial ionization anomaly (EIA) (Vineeth et al., 2007; Chau et al., 2009; Goncharenko et al., 2010; Pancheva and Mukhtarov, 2011; Jin et al., 2012).

.....

P. 2, line 41: “Since planetary waves in the Southern Hemisphere (SH) generally have smaller amplitudes than in the Northern Hemisphere (NH).....” generally this is true only for the SPWs; the climatology of some other well known PWs, as for example, the quasi-2-day W3 wave or the quasi-6-day W1 wave are both stronger in the SH.

**Answer:** Thank you for your remind. “planetary waves” has been changed to “stationary planetary waves”.

.....  
P. 3, line 79:.....  $\sim\pm 15^\circ$  N MLAT.... please, delete N

**Answer:** Revised.

.....  
P. 4, line 117: “In additional, we .....”; please, delete al

**Answer:** Revised.

.....  
P. 5, lines 126-127: “Note that quasi 10-day oscillations of northern and southern EIA crests are in-phase, which...”; sorry, both oscillations are not in phase; the oscillation of the NH crest indicates a delay of a day with respect to the SH one. Please, calculate the cross-correlation function between both times, particularly between days 220-290 when they have large amplitudes, and will see that the largest cross-correlation will be found at different from zero time lag.

**Answer:** It is a very good comment. According to this suggestion, the cross-correlation function is used to reveal the phase relationship between northern and southern EIA crests. The results show that the quasi 10-day wave of northern EIA crest delay 1 day behind southern EIA crest.

**Line # [136-141]:** To further verify this, Figure 7 shows the cross-correlation of quasi 10-day waves in MLAT location (a) and TEC (b) between northern and southern EIA crests. The cross-correlation coefficients of MLAT location and TEC reach 0.8 and 0.93, respectively. Moreover, the maximum cross-correlation coefficients for MLAT location is at 1 day, indicating that the wave of northern EIA crest delay 1 day behind southern EIA crest. This phase difference between northern and southern EIA crests may be due to differences in longitude between two GPS stations.

1     **Quasi 10-day wave modulation of equatorial ionization anomaly during the Southern**  
2                                   **Hemisphere stratospheric warming of 2002**

3  
4     Xiaohua Mo<sup>1</sup>, Donghe Zhang<sup>2\*</sup>

5  
6     <sup>1</sup>College of Science, Key Laboratory for Ionospheric Observation and Simulation, Guangxi University for  
7     Nationalities, Nanning, China

8     <sup>2</sup>Department of Geophysics, Peking University, Beijing, 100871, China

9  
10    Correspondence: Donghe Zhang ([zhangdh@pku.edu.cn](mailto:zhangdh@pku.edu.cn))

11  
12  
13    **Abstract**

14        The present paper studies the perturbations in equatorial ionization anomaly (EIA) region during the  
15    Southern Hemisphere (SH) sudden stratospheric warming (SSW) of 2002, using the location of EIA crests  
16    derived from Global Positioning System (GPS) station observations, the Total Electron Content (TEC)  
17    obtained by International GNSS Service (IGS) global ionospheric TEC map (GIMs), and the equatorial  
18    electrojet (EEJ) estimated by geomagnetic field in Asian sector. A strong quasi 10-day periodic oscillation  
19    is clearly identified in northern and southern EIA region. This quasi 10-day oscillation is also seen in the  
20    polar stratospheric temperature and EEJ, suggesting the enhanced quasi-10-day planetary wave associated  
21    with SSW produced oscillation in EIA region through modulating the equatorial fountain effect. Our  
22    results reveal some newer features of ionospheric variation that have not been reported during Northern  
23    Hemisphere (SH) SSWs.

31 **1. Introduction**

32 Sudden stratospheric warming (SSW) is large-scale meteorological process in the polar stratosphere  
33 which is characterized by rapid rise in temperatures and deceleration/reversal in the zonal mean flows  
34 (Scherhag, 1952). The primary driver of SSW is thought to be a rapid growth of quasi-stationary planetary  
35 wave interacting with zonal mean flow (Matsuno, 1971). Although the main processes of SSW occur in the  
36 middle atmosphere, its effects on the ionosphere have been observed in significant changes of equatorial  
37 electrojet (EEJ), vertical plasma drift, and equatorial ionization anomaly (EIA) (Vineeth et al., 2007; Chau  
38 et al., 2009; Goncharenko et al., 2010; Pancheva and Mukhtarov, 2011; Jin et al., 2012). These ionospheric  
39 variations mainly display similar semidiurnal pattern and 13- to 16-day wave signatures which have been  
40 associated with planetary wave, solar and lunar tide wave (Pedatella and Forbes, 2009; Goncharenko et al.,  
41 2010; Fejer et al., 2010; Park et al., 2012). Since stationary planetary waves in the Southern Hemisphere  
42 (SH) generally have smaller amplitudes than in the Northern Hemisphere (NH) where orographic and  
43 thermal forcing is stronger (Andrews et al., 1987), major SSWs often occur in NH. Therefore, most studies  
44 about SSW effects on the ionosphere are during NH SSW period.

45 In August to September 2002, three minor SSWs and a major SSW appeared in SH (Varotsos 2002;  
46 Baldwin et al., 2003). There is sufficient evidence that a series of unusual atmospheric states occurred in  
47 this period, i.e., planetary wave scale quasi 10-day variation (Krüger et al., 2005; Palo et al., 2005),  
48 short-term semidiurnal tide variability with zonal wave number  $s=1$  (Chang et al., 2009) and the winds  
49 oscillation with ~14-days period (Andrew et al., 2004), are all linked to the extremely large planetary wave  
50 events. Although the atmospheric activity in connection with 2002 SH SSW has been well revealed in  
51 observations and numerical modeling, relatively little is known about the ionosphere effects of 2002 SH  
52 SSW. Recently, Olson et al. (2013) studied the equatorial electrodynamic perturbations in Peruvian sector  
53 during 2002 SH SSW and found enhanced quasi 2-day fluctuations and large amplitude multi-day  
54 perturbations in EEJ and vertical drifts. The researches of ionospheric behavior during SH SSW periods  
55 are useful for verifying the existing explanation about the origin of ionospheric perturbations during NH  
56 SSW periods and revealing some newer features of ionospheric variation, so further investigation of 2002  
57 SH SSW effect on ionosphere with more ionospheric parameters is still warranted.

58 In the present study, we present the first observational evidence of quasi 10-day oscillation in EIA  
59 region during 2002 SH SSW which has not been reported during NH SSWs, based on the location of EIA  
60 crests derived from Global Positioning System (GPS) station observations, the Total Electron Content

61 (TEC) obtained by International GNSS Service (IGS) global ionospheric TEC map (GIMs), and the EEJ  
62 estimated by geomagnetic field in Asian sector.

63

## 64 2. Data and Methods

65 The location of EIA crests derived from GPS observations are used to analyze the variation in EIA  
66 region during 2002 SH SSW from July 20, 2002 to October 27, 2002. The GPS stations are GUAN  
67 (23.19°N, 113.34°E, MLAT~12.52°N) and BAKO (6.49°S, 106.84°E, MLAT~17.18°S) which are near  
68 northern and southern EIA crest, respectively. The locations of the GPS stations are shown in Figures 1.  
69 Since the ionospheric vertical TEC usually reach the maximum at EIA crest, the location of EIA crest can  
70 be obtained by vertical TEC values at each ionospheric penetration point (IPP), which is the intersection of  
71 the line of sight and the ionospheric shell (assumed to be 400 km) (Mo et al., 2014). The relative accuracy  
72 of the TEC is 0.02 total electron content unit (1TECU= $10^{16}$  eI m<sup>-2</sup>) (Hofmann-Wellenhof et al., 1992). The  
73 sample rate of these GPS stations were 30s, so the resolution of the location of EIA crest is less than 25 km  
74 (Mo et al., 2017). Figures 2a and 2b show the daily average geomagnetic latitude (MLAT) of northern and  
75 southern EIA crests during 2002 SH SSW.

76 The TEC from GIMs are also used to analyze the variation in EIA region. The GIMs provides maps of  
77 TEC obtained from a global network of GPS receivers, which have temporal resolution of 2 hours and  
78 spatial resolution of 5° in longitude and 2.5° in latitude (Mannucci et al., 1998). The EIA crest usually  
79 reaches its maximum development near 14:00 LT (Huang et al., 1989; Yeh et al., 2001), so the daily  
80 average TEC obtained by GIMs at 12~14 LT,  $\pm 5^\circ \sim \pm 15^\circ$  MLAT, 100°~150°E every day in Asian sector  
81 are used to describe the variation in northern and southern EIA region, the results are shown in Figures 2c  
82 and 2d.

83 To demonstrate the dynamical process in EIA region, the EEJ is also used in this study, which can be  
84 estimated by the difference between the horizontal component of geomagnetic field at TIR (8.7°N, 77.8°E,  
85 MLAT~0.03°N) and VSK (17.68°N, 83.32°E, MLAT~8.56°N) (Rastogi et al., 1990). The results are  
86 shown in Figures 2e. In addition, the polar stratospheric temperature (90°S, 10hPa) and zonal mean zonal  
87 winds (60°S, 10hPa) obtained from National Centers for Environment Prediction (NCEP) are used to  
88 examine the extent of the SSW, the results are shown in Figures 2f and 2g. The background of  
89 geomagnetic activity index (Kp) and solar flux index (F10.7) from the websites <http://spidr.ngdc.noaa.gov/>  
90 are depicted in Figures 2h and 2i.



### 91 3. Results and Analysis

92 It can be seen from Figures 2f and 2g that there were three obvious minor SH SSW events around day  
93 number 230-260 and a major SH SSW event around day number 263-288 (Olson et al., 2013). Figure 3  
94 shows the contour map of polar stratospheric temperature (80°S, 10hPa) obtained from NCEP from July 20,  
95 2002 to October 27, 2002. An eastward phase progression of quasi 10-day wave is clearly observed around  
96 day number 210-270. With SABER temperature data, Palo et al. (2005) also observed similar disturbance  
97 and suggested it consists of an eastward-propagating quasi 10-day wave with zonal wave numbers  $s=1$   
98 superimposed upon a large stationary planetary wave with  $s=1$ .

99 Now we examine the impact of this quasi 10-day wave on EIA region. It should be noted the  
100 solar/magnetospheric forcing on ionosphere are strong due to 2002 SH SSW event occurring during solar  
101 maximum year. To exclude these long period fluctuations in EIA region associated with  
102 solar/magnetosphere forcing, the periods longer than 15 days in the MLAT location and TEC of EIA crest,  
103 and EEJ are removed. Specifically, these parameters are subtracted from their respective 15-day moving  
104 average. The residuals are subjected to Lomb-Scargle (L-S) spectral analysis (Lomb,1976; Scargle, 1982),  
105 and the results are shown in Figures 4a, 4b, 4c, 4d, and 4e. The horizontal dashed lines represent the 95%  
106 confidence level. It is evident that the MLAT location and TEC of EIA crest, and EEJ all exhibit significant  
107 quasi 10-day periodic component, which exceed or approach 95% confidence level, suggesting that the  
108 whole dynamical process in EIA region is modulated by quasi 10-day wave. Figures 4f and 4g show the  
109 L-S spectral analysis of Kp and F10.7. It can be seen that spectral component of Kp also has quasi 10-day  
110 periodic component which will be related to solar wind high-speed streams (Lei et al., 2008). However,  
111 this quasi 10-day periodic component is too weak to be identified in F10.7, indicating that variation in the  
112 solar flux cannot account for this quasi 10-day oscillation in EIA region.

113 To investigate the time evolution of quasi 10-day periodic variation, the Morlet wavelet spectral  
114 analysis is applied to MLAT location and TEC of EIA crest, EEJ and Kp which exhibit quasi 10-day  
115 oscillation. The periods longer than 15 days in the MLAT location and TEC of EIA crest , and EEJ are  
116 removed before the wavelet spectra is generated, and the results are illustrated in Figures 5a, 5b, 5c, 5d,  
117 and 5e. The black solid contours in each panel indicate a significance level higher than 95%. The white line  
118 in each panel represents the cone of influence of the wavelet analysis. The color bar number is the power  
119 strength for each parameter. Obviously, the most predominant periodic component in the MLAT location  
120 and TEC of EIA crest, and EEJ are quasi 10-day period, which mainly appeared around day number

121 210-290, indicating quasi 10-day oscillations in EIA region go through three minor SSWs and a major  
122 SSW period. The time evolution of the power in MLAT location and TEC of northern EIA crest match well  
123 those of southern EIA crest, respectively. In addition, we note both the MLAT location and the TEC of EIA  
124 crest show the quasi 2-day oscillations during major SSW period (around day number 260-270), which are  
125 also found on equatorial ionospheric electric fields and currents at the same period (Olson et al., 2013).  
126 Figure 5f shows the wavelet spectral analysis of Kp index. It can be seen that quasi 10-day periodic  
127 component is nearly absent in Kp around day number 230-290, suggesting that magnetic activity should  
128 not be the driving force for this quasi 10-day oscillation in EIA region.

129 In order to demonstrate the phase relationship of the quasi 10-day oscillations between northern and  
130 southern EIA crests, the band-pass filter is performed on the MLAT location and TEC of EIA crest. The  
131 absolute values of the MLAT location of EIA crest are used. The band-pass filter is centered at the period  
132 of 10-day, with half-power points at 8-day and 12-day, and the results are shown in Figure 6. The quasi  
133 10-day wave amplitudes of northern and southern EIA crests are roughly equivalent, which exceed 1.7  
134 degree for MLAT location and 7 TECU for TEC, respectively. Although the quasi 10-day wave of northern  
135 EIA crest match well those of southern EIA crest, the wave of northern EIA crest seemed to delay behind  
136 southern EIA crest, especially for MLAT location. To further verify this, Figure 7 shows the  
137 cross-correlation of quasi 10-day waves in MLAT location (a) and TEC (b) between northern and southern  
138 EIA crests. The cross-correlation coefficients of MLAT location and TEC reach 0.8 and 0.93, respectively.  
139 Moreover, the maximum cross-correlation coefficients for MLAT location is at 1 day, indicating that the  
140 wave of northern EIA crest delay 1 day behind southern EIA crest. This phase difference between northern  
141 and southern EIA crests may be due to differences in longitude between two GPS stations.

142

#### 143 4. Discussions

144 In recent years a series of reports have focused on ionospheric perturbations during SSW event. The  
145 most predominant features in ionosphere associated with SSW event are semidiurnal pattern and 13- to  
146 16-day wave variations, which are attributed to nonlinear interaction of planetary wave, solar and lunar  
147 tide wave (Pedatella and Forbes, 2009;Goncharenko et al., 2010; Fejer et al., 2010;Park et al., 2012). As  
148 major SSW often occurs in NH, most studies about SSW effects on the ionosphere are during NH SSW  
149 period. In August to September 2002, the first major SSW was observed in SH. The NH and SH SSW  
150 occurred in Arctic and Antarctic winter, respectively, so the occurring time and location of SH SSW are

151 opposite to those of NH SSW. The researches of ionospheric behavior during SH SSW periods are useful  
152 for testing the general rule of ionospheric perturbations during NH SSW periods. For example, Olson et al.  
153 (2013) demonstrated that multi-day ionospheric perturbations responding to 2002 SH SSW resemble those  
154 observed during NH SSWs and these ionospheric perturbations were associated with enhanced lunar tidal  
155 effects.

156 In the current study we present observations of quasi 10-day oscillation in EIA region during the  
157 2002 SH SSW that **have** not been reported during NH SSWs. This quasi 10-day oscillation is absent and  
158 weak in Kp and F10.7 index, indicating that the magnetic activity and solar flux cannot account for this  
159 quasi 10-day oscillation in EIA region. Meanwhile, an unusual atmospheric state occurred in this period  
160 that the ozone hole over the Antarctic has a smaller size and splits into two separate holes (Varotsos 2002;  
161 Baldwin et al., 2003). This phenomenon is thought to be due to high temperatures the Antarctic, which was  
162 contributed to by upward propagation of a planetary wave (Venkat Ratnam et al., 2004). Moreover, strong  
163 planetary wave scale quasi 10-day variation was observed in polar stratospheric temperature during this  
164 period (Krüger et al., 2005; Palo et al., 2005), so the quasi 10-day oscillations in EIA region may be related  
165 to atmosphere perturbations linking the SSW in the Southern Hemisphere.

166 **A series of studies have showed** how the quasi 10-day planetary wave in stratosphere can penetrate  
167 into the ionosphere E region. Krüger et al. (2005) revealed the eastward-traveling waves with periods near  
168 10 days and their interaction with quasi-stationary planetary waves forced in the troposphere during 2002  
169 SH SSW event, supporting the observational and numerical evidence that the eastward traveling wave  
170 interacts with the stationary wave to produce a quasi-periodic amplitude modulation of the stationary  
171 waves (Hirota et al., 1990; Ushimaru and Tanaka, 1992). Palo et al. (2005) found an eastward-propagating  
172 quasi 10-day wave with zonal wave numbers  $s=1$  and  $s=2$ , and a quasi-stationary planetary waves with  $s=1$   
173 extend from the lower stratosphere to the 100-120 km height region with little amplitude attenuation.  
174 While the quasi-stationary planetary wave is confined to the high latitude atmosphere and cannot directly  
175 propagate to equatorial ionosphere, the tides were introduced into planetary wave modulation mechanism.  
176 Eswaraiyah et al. (2018) reported that zonal diurnal and semidiurnal tide amplitudes in Antarctica  
177 mesosphere and lower thermosphere were enhanced around day number 230-290 during 2002 SH SSW,  
178 which coincides with the enhanced period of quasi 10-day oscillations in EIA region shown in Figure 5.  
179 Moreover, Chang et al. (2009) showed that the short-term variability of the  $s=1$  semidiurnal tide is strongly  
180 dependent upon the PW1 events (quasi-10-day wave) prior to the major warming during 2002 SH SSW,

181 supporting the suggestion that the quasi-stationary planetary wave can influence migrating and  
182 nonmigrating solar tides globally (Liu et al., 2010; Pedatella and Forbes, 2010). So the interactions  
183 between quasi-10-day planetary wave and tide will modify the ionosphere E-region winds, which can  
184 produce E-region electric fields via the E-region dynamo process. In this study, the evidence of quasi  
185 10-day periodic variation of EIA crests and EEJ strongly supports the suggestion that quasi-10-day  
186 planetary wave produced oscillation in EIA region through modulating E-region electric fields.  
187 Specifically, the E-region electric fields map to F-region along the magnetic field lines and generate an  
188 eastward electric field (Goncharenko, 2010). At the magnetic equator, the eastward electric field with quasi  
189 10-day periodic variation change electron density distribution in the low-latitude region via  $\vec{E} \times \vec{B}$  drift,  
190 and finally leads to quasi 10-day planetary waves characteristic variations in EIA region. The effects of  
191 planetary wave on the equatorial fountain effect have been revealed by the evidence that the vertical and  
192 latitudinal structures of the 6-day oscillation in the F region ionosphere peak on both sides of the equator in  
193 the EIA region (Gu et al., 2014). The synchronous 10-day oscillation between northern and southern EIA  
194 crest in our results are consistent with these observations.

195 In our prior studies, a 14- to 15-day wave during several NH SSW events is ascribed to lunar tide  
196 (Mo et al., 2018). So the source of quasi 10-day oscillations in EIA region during 2002 SH SSW is  
197 different from 14- to 15-day waves during NH SSW. Moreover, no obvious 14- to 15-day oscillation is  
198 found in EIA region during 2002 SH SSW, which may be that the equatorial lunar semidiurnal effects  
199 during September-October are weaker than that during January-February (Stening et al., 2011; Pedatella,  
200 2014). Olson et al. (2013) also reported that the perturbations amplitude of EEJ and vertical drifts  
201 modulated by lunar semidiurnal tides during SH SSW are smaller than those during NH SSW.

202

## 203 5. Conclusions

204 Using the location and TEC of EIA crests derived from GPS station observations and GIMs, we found  
205 a quasi 10-day periodic variability in northern and southern EIA region in Asian sector during the SH SSW  
206 of 2002. In the same time period, this quasi 10-day oscillation is also seen in the polar stratospheric  
207 temperature and EEJ, which is absent and weak in Kp and F10.7 index, respectively. Previous studies have  
208 shown that a strong quasi 10-day planetary wave with zonal wave numbers  $s=1$  extend from the lower  
209 stratosphere to mesosphere and lower thermosphere (Palo et al., 2005), so the quasi 10-day variation in  
210 EIA region should be ascribed to enhanced 10-day planetary wave in lower atmosphere associated with

211 SSW.

212 **Acknowledgements:** The GPS data were from the Crustal Movement Observation Network of China (via  
213 e-mail to yglyang@cma.gov. cn) and IGS (available at <http://sopac.ucsd.edu>). The geomagnetic data at  
214 TIR and VSK were from WDC for Geomagnetism, Kyoto (available  
215 at <http://wdc.kugi.kyoto-u.ac.jp/hyplt/index.html>). The GIMs were downloaded from the site  
216 <ftp://cddis.gsfc.nasa.gov>. This research was jointly supported by the National Natural Science Foundation  
217 of China (41864006, 41674157, and 41464006), Guangxi Natural Science Foundation  
218 (2016GXNSFAA380132), and Chinese Meridian Project. We gratefully acknowledge National Center for  
219 Environmental Prediction (NCEP) for providing public access to stratospheric data (available at  
220 <https://www.esrl.noaa.gov/psd/data/reanalysis/>).

## 221 References

- 222 Andrews, D. G., Holton, J. R., and Leovy, C. B.: Middle Atmosphere Dynamics, Academic, San Diego,  
223 Calif, 1987.
- 224 Andrew, J. D., Vincent, R. A., Murphy, D. J., Tsutsumi, M., Riggin, D. M., and Jarvis, M. J.: The  
225 large-scale dynamics of the mesosphere-lower thermosphere during the Southern Hemisphere  
226 stratospheric warming of 2002, *Geophys. Res. Lett*, 31, L14102, doi:10.1029/2004GL020282, 2004
- 227 Baldwin, M., Hirooka, T., O'Neill, A., and Yoden, S.: major stratospheric warming in the Southern  
228 Hemisphere in 2002: Dynamical aspects of the ozone hole split, *SPARC Newsl.*, 20, 24-26, 2003.
- 229 Chang, L. C., Palo, S. E., and Liu, H. -L.: Short-term variation of the s=1 nonmigrating semidiurnal tide  
230 during the 2002 stratospheric sudden warming, *J. Geophys. Res.*, 114, D03109,  
231 doi:10.1029/2008JD010886, 2009.
- 232 Chau, J. L., Fejer, B. G., and Goncharenko, L. P.: Quiet variability of equatorial  $E \times B$  drift during a  
233 sudden stratospheric warming event. *Geophys. Res. Lett*, 36,  
234 L05101, <https://doi.org/10.1029/2008GL036785>, 2009.
- 235 Eswaraiah, S., Kim, Y. H., Lee, J., Ratnam, M. V., and Rao, S. V. B.: Effect of Southern Hemisphere  
236 sudden stratospheric warmings on Antarctica mesospheric tides: First observational study. *J. Geophys.*  
237 *Res. Space Physics*, 123, 2127–2140, <https://doi.org/10.1002/2017JA024839>, 2018.
- 238 Fejer, B. G., Olson, M. E., Chau, J. L., Stolle, C., Lühr, H., Goncharenko, L. P., Yumoto, K., and  
239 Nagatsuma, T.: Lunar-dependent equatorial ionospheric electrodynamic effects during sudden  
240 stratospheric warmings, *J. Geophys. Res.*, 115, A00G03, doi:10.1029/2010JA015273, 2010.

241 Goncharenko L. P., Chau, J. L., Liu, H. -L., and Coster, A. J.: Unexpected connections between the  
242 stratosphere and ionosphere, *Geophys. Res. Lett.*, 37, L10101, doi:10.1029/2010GL043125, 2010.

243 Gu, S.-Y., Liu, H.-L., Li, T., Dou, X., Wu, Q., and Russell III, J. M.: Observation of the neutral-ion  
244 coupling through 6 day planetary wave, *J. Geophys. Res. Space Physics*, 119, 10,376–10,383,  
245 doi:10.1002/2014JA020530, 2014.

246 Hirota, I., Kuroi, K., and Shiotani, M.: Midwinter warmings in the Southern Hemisphere stratosphere in  
247 1988, *Q. J. R. Meteorol., Soc.*, 116, 929-941, 1990.

248 Hofmann - Wellenhof, B., Lichtenegger, H., and Collins, J.: *GPS—Theory and Practice*, Springer, New  
249 York, 1992.

250 Huang, Y. N., Cheng, K., and Chen, S. W.: On the equatorial anomaly of the ionospheric total electron  
251 content near the northern anomaly crest region, *J. Geophys. Res.*, 94(A10), 13,515–13,525, 1989.

252 Jin, H., Miyoshi, Y., Pancheva, D., Mukhtarov, P., Fujiwara, H., and Shinagawa, H.: Response of the  
253 migrating tides to the stratospheric sudden warming in 2009 and their effects on the ionosphere studied  
254 by a whole atmosphere-ionosphere model GAIA with COSMIC and TIMED/SABER observations, *J.*  
255 *Geophys. Res.*, 117, A10323, doi:10.1029/2012JA017650, 2012.

256 Krüger, K., Naujokat, B., and Labitzke, K.: The unusual midwinter warming in the Southern Hemisphere  
257 stratosphere 2002: A comparison to Northern Hemisphere phenomena, *J. Atmos. Sci.*, 62, 603-613,  
258 2005.

259 Lei, J., Thayer, J. P., Forbes, J. M., Wu, Q., She, C., Wan, W., and Wang, W.: Ionosphere response to solar  
260 wind high-speed streams, *Geophys. Res. Lett.*, 35, L19105, doi:10.1029/2008GL035208, 2008.

261 Liu, H. -L, Wang, W., Richmond, A. D., and Roble, R. G.: Ionospheric variability due to planetary waves  
262 and tides for solar minimum conditions, *J. Geophys. Res.*, 115, A00G01, doi:10.1029/2009JA015188,  
263 2010.

264 Lomb, N. R.: Least-squares frequency analysis of unequally spaced data, *Astrophys.Space Sci.*, 39,  
265 447-462, 1976.

266 Mannucci, A. J., Wilson, B. D., Yuan, D. N., Ho, C. M., Lindqwister, U. J., and Runge, T. F.: A global  
267 mapping technique for GPS derived ionospheric total electron content measurements, *Radio Sci.*, 33,  
268 565-582, doi:10.1029/97RS02707, 1998.

269 Matsuno, T.: A dynamical model of the stratospheric sudden warming, *J. Atmos. Sci.*, 28, 1479-1494, 1971.

270 Mo, X. H., Zhang, D. H., Goncharenko, L. R., Hao, Y. Q., and Xiao, Z.: Quasi-16-day periodic meridional

271 movement of the equatorial ionization anomaly, *Ann. Geophys.*, 32, 121-131, 2014.

272 Mo, X. H., Zhang, D. H., Goncharenko, L. R., Zhang, S. R., Hao, Y. Q., Xiao, Z., Pei, J. Z., Yoshikawa, A.,  
273 Chau, H. D.: Meridional movement of northern and southern equatorial ionization anomaly crests in the  
274 East-Asian sector during 2002–2003 SSW, *Science China Earth Sciences*, 60(4), 776–785, [https://](https://doi.org/10.1007/s11430-016-0096-y)  
275 [doi.org/10.1007/s11430-016-0096-y](https://doi.org/10.1007/s11430-016-0096-y), 2017.

276 Mo, X. H., and Zhang, D. H.: Lunar tidal modulation of periodic meridional movement of equatorial  
277 ionization anomaly crest during sudden stratospheric warming, *J. Geophys. Res. Space Physics*, 123,  
278 1488-1499. <https://doi.org/10.1002/2017JA024718>, 2018.

279 Olson, M. E., Fejer, B. G., Stolle, C., Lühr, H., and Chau, J. L.: Equatorial ionospheric electrodynamic  
280 perturbations during Southern Hemisphere stratospheric warming events, *J. Geophys. Res. Space*  
281 *Physics*, 118, 1190-1195, doi:10.1002/jgra.50142, 2013.

282 Palo, S. E., Forbes, J. M., Zhang, X., Russell III, J. M., Mertens, C. J., Mlynczak, M. G., Burns, G. B., Espy,  
283 P. J., and Kawahara, T. D.: Planetary wave coupling from the stratosphere to the thermosphere during  
284 the 2002 Southern Hemisphere pre-stratwam period, *Geophys. Res. Lett.*, 32, L23809,  
285 doi:10.1029/2005GL024298, 2005.

286 Pancheva, D, and Mukhtarov, P.: Stratospheric warmings: The atmosphere-ionosphere coupling paradigm,  
287 *J. Atmos. Sol. Terr. Phys.*, 73, 1697-1702, doi:10.1016/j.jastp.2011.03.006, 2011.

288 Park, J., Lühr, H., Kunze, M., Fejer, B. G., and Min, K. W.: Effect of sudden stratospheric warming on  
289 lunar tidal modulation of the equatorial electrojet, *J. Geophys. Res.*, 117, A03306,  
290 doi:10.1029/2011JA017351, 2012.

291 Pedatella, N. M., and Forbes, J. M.: Modulation of the equatorial F-region by the quasi-16 day planetary  
292 wave, *Geophys. Res. Lett.*, 34, L09105, doi:10.1029/2009GL037809, 2009.

293 Pedatella, N. M., and Forbes, J. M.: Evidence for stratosphere sudden warming-ionosphere coupling due to  
294 vertically propagating tides, *Geophys. Res. Lett.*, 37, L11104, doi:10.1029/2010GL043560, 2010.

295 Pedatella, N. M.: Observations and simulations of the ionospheric lunar tide: Seasonal variability, *J.*  
296 *Geophys. Res. Space Physics*, 119, 5800-5806, doi:10.1002/2014JA020189, 2014.

297 Rastogi, R. G., and Klobuchar, J. A. : Ionospheric electron content within the equatorial F2 layer anomaly  
298 belt, *J. Geophys. Res.*, 95(A11), 19,045–19,052, <https://doi.org/10.1029/JA095iA11p19045>, 1990.

299 Scargle, J. D.: Studies in astronomical time series analysis. II. Statistical aspects of spectral analysis of  
300 unevenly spaced data, *Astrophys. J.*, 263, 835-853, 1982.

301 Scherhag, R.: Die explosionsartigen Stratosphärenwärmungen des Spätwinters 1952, Ber. Dtsch  
302 Wetterdienstes USZone, 6, 51-63, 1952.

303 Stening, R. J.: Lunar tide in the equatorial electrojet in relation to stratospheric warmings, J. Geophys. Res.,  
304 116, A12315, doi:10.1029/2011JA017047, 2011.

305 Ushimaru, S., and Tanaka, H.: A numerical study of the interaction between stationary Rossby waves and  
306 eastward-traveling waves in the Southern Hemisphere stratosphere, J. Atmos., Sci., 49, 1354-1373,  
307 1992.

308 Varotsos, C.: The Southern Hemisphere ozone hole split in 2002, Environ. Sci. Pollut. Res., 9, 375-376,  
309 2002.

310 Venkat Ratnam, M., Tsuda, T., Jacobi, C., and Aoyama, Y.: Enhancement of gravity wave activity observed  
311 during a major Southern Hemisphere stratospheric warming by CHAMP/GPS measurements, Geophys.  
312 Res. Lett., 31, L16101, doi:10.1029/2004GL019789, 2004.

313 Vineeth, C., Pant, T. K., Devasia, C. V., and Sridharan, R.: Atmosphere-ionosphere coupling observed over  
314 the dip equatorial MLTI region through the quasi 16-day wave. Geophys. Res. Lett., 34,  
315 L12102, <https://doi.org/10.1029/2007GL030010>, 2007.

316 Yeh, K. C., Franke, S. J., Andreeva, E. S., and Kunitsyn, V. E.: An investigation of motions of the  
317 equatorial anomaly crest, Geophys. Res. Lett., 28(24), 4517-4520, doi:10.1029/2001GL013897, 2001.


318

319

320


321






322  
323

**Figure 1.** Location of the GPS stations in Asian sector.



324  
 325 **Figure 2.** The magnetic latitude (MLAT ) location of (a) northern and (b) southern equatorial ionization  
 326 anomaly (EIA) crest; The TEC of (c) northern and (d) southern EIA crest; the (e) equatorial electrojet  
 327 (EEJ), (f) polar stratospheric temperature (at 90°S, 10hPa) and (g) zonal wind (at 60°S, 10hPa) from  
 328 National Centers for Environment Prediction; the (h) Geomagnetic activity index, Kp and (i) solar flux  
 329 index F10.7 during the period from July 20, 2002 to October 27, 2002.




330

331 **Figure 3.** The contour map of polar stratospheric temperature (80°S, 10hPa) obtained from NCEP during

332

the same period as in Figure 2.




333

334

335

336

**Figure 4.** Lomb-Scargle periodograms of the MLAT location of (a) northern and (b) southern EIA crest, the TEC of (c) northern and (d) southern EIA crest, (e) EEJ, (f) Kp index and (g) F10.7 during the same period as in Figure 2.




337

338 **Figure 5.** The wavelet power spectra of the MLAT location of (a) northern and (b) southern EIA crest, the

339 TEC of (c) northern and (d) southern EIA crest, (e) EEJ and (f) Kp index during the same period as in

340 Figure 2. The white line in each panel represents the cone of influence of the wavelet analysis.



341


342

**Figure 6.** The band-pass filter results of the (a) MLAT location of (solid line) northern and (dash-dotted line) southern EIA crest, the (b) TEC of (solid line) northern and (dash-dotted line) southern EIA crest

343

344

during the same period as in Figure 2.



345

346

347

**Figure 7.** The cross-correlation of quasi 10-day waves in MLAT location (a) and TEC (b) between northern and southern EIA crest.