

Review on “Quasi 10-day wave modulation of equatorial ionization anomaly during the Southern Hemisphere stratospheric warming of 2002” by Xiaohua Mo and Donghe Zhang

Using the location and TEC of EIA crests derived from GPS station observations and GIMs, the authors report on a quasi 10-day periodic variability in northern and southern EIA region in Asian sector during the 2002 SH SSW. Around the same time period, this quasi 10-day oscillation is also seen in the polar stratospheric temperature and EEJ, which is absent and weak in Kp and F10.7 index, respectively. Given previous work showing a strong quasi 10-day planetary wave with zonal wave numbers $s=1$ extend from the lower stratosphere to mesosphere and lower thermosphere the authors infer that the quasi 10-day variation in the EIA region should be ascribed to enhanced 10-day planetary wave in lower atmosphere associated with SSW.

General comment:

This manuscript contains some interesting results and I find it improved in this revised version. In particular I find the results of the 10-day modulation of the TEC during the 2002 SH SSW event compelling. A major source of concern is lack of sufficient evidence on potential connections between the 10-day periodic variation of EIA crests and SSW/EEJ. Further statistical analysis that demonstrates this connection is required before publication can be recommended.

Answer:

We very appreciate your substantial comments for our study about “Quasi 10-day wave modulation of equatorial ionization anomaly during the Southern Hemisphere stratospheric warming of 2002”. The connections between the 10-day periodic variation of EIA crests and SSW/EEJ have been further analyzed and discussed in the introduction and discussions part of revised manuscript. All major revisions are marked in yellow highlights. Thank you very much.

Line # [39-67]: Since stationary planetary waves in the Southern Hemisphere (SH) generally have smaller amplitudes than in the Northern Hemisphere (NH) where orographic and thermal forcing is stronger (Andrews et al., 1987), major SSWs often occur in NH. Therefore, most studies about SSW effects on the ionosphere are during NH SSW period. In recent years, a great deal of research has been focused on the variation of the low latitude ionosphere during SSW period in the northern hemisphere, and the quasi-16-day periodic disturbance and the lunar tide characteristics have been found in some ionospheric parameters, for example, EEJ, vertical plasma drifts, ionospheric electron density (Vineeth et al., 2007; Chau et al., 2009; Pedatella and Forbes, 2009; Goncharenko et al., 2010). Some researchers considered that this kind of quasi-16-day periodic variations is related to the enhanced planetary wave during the SSW period (Chau et al., 2009; Pedatella and Forbes, 2009; Liu et al., 2010). But others believe that this kind of quasi-16-day period is related to the semi-diurnal lunar tides (Fejer et al., 2010; Park et al., 2012). The direct evidence is the typical SSW feature appears in some ionospheric parameters as morning enhancement and afternoon decrease, in a semi-diurnal pattern that progresses to later local times within several days. However, some studies believe that the local time variation characteristics are not necessarily caused by the semidiurnal lunar tides, Pedatella et al. (2012) demonstrate that the phase of the semidiurnal solar tide changes in a manner that makes it similar to the phase of the lunar semidiurnal tide. Besides,

although many studies on the variation of the low latitude ionosphere during the SSW period, the physical connection between the SSW in the polar region and the featured variations in low latitude ionosphere is still not clear. Some studies even consider that the SSW and the featured variation of low latitude ionosphere is co-source, which is the effect of enhanced planetary waves in different regions (Stening et al., 2011).

In comparison, the atmospheric parameters in the mesosphere and lower thermosphere (MLT) are very limited. The atmospheric variation in MLT is usually indirectly reflected through the EEJ obtained from the geomagnetic field data in equatorial and low latitude region. The EEJ is driven by zonal electric field which also produces EIA via upward $\vec{E} \times \vec{B}$ drift (fountain effect). The zonal electric field modulated by tidal winds in the lower thermosphere through the E-region dynamo process is easy to be influenced by various atmospheric waves, so these ionospheric variations often display similar semidiurnal pattern and 13- to 16-day wave signatures which have been associated with enhanced planetary wave, solar and lunar tide wave during SSW period (Pedatella and Forbes, 2009; Goncharenko et al., 2010; Fejer et al., 2010; Park et al., 2012).

Line # [186-190]: The wave interactions between eastward-propagating waves with periods near 10 days, quasi-stationary planetary waves, and the zonal mean atmospheric state were eventually driven towards total break-down of the polar vortex and a major warming of the stratosphere (Krüger et al., 2005; Palo et al., 2005). So the quasi 10-day oscillations in EIA region should be ascribed to atmosphere perturbations linking the SSW in the Southern Hemisphere.

Line # [211-227]: In this study, the EEJ driven by equatorial zonal electric field also exhibits quasi 10-day oscillation, indicating that the upward-propagating planetary waves interacted with the tide produced oscillation in EIA region through modulating E-region electric fields. Specifically, the E-region electric fields map to lower ionospheric F-region along the magnetic field lines and generate an eastward electric field (Goncharenko et al., 2010). At the magnetic equator, the eastward electric field with quasi 10-day periodic variation change electron density distribution in the low-latitude region via $E \times B$ drift, and finally leads to quasi 10-day planetary waves characteristic variations in EIA region. Previous studies have revealed a strong correlation between ionospheric perturbations and the occurrence of NH SSW. During NH SSW period, quasi 16-day oscillations and semidiurnal pattern are observed in equatorial mesopause temperature, the MLT meridional and zonal wind, EEJ, electron density and TEC (Vineeth et al., 2007; Pedatella and Forbes, 2009; Park et al., 2012 ; Jonah et al., 2014). Some researchers attribute these ionospheric perturbations to the strong dynamical coupling between the lower atmosphere and ionosphere through the intensification of planetary wave activity (Chau et al., 2009), lunar (Fejer et al., 2010) and solar (Pedatella et al., 2012) tide. In this study, the consistent quasi 10-day oscillations appear in EEJ, the location and TEC of northern and southern EIA crest, indicating that coupling mechanism between the lower atmosphere and ionosphere during SH SSW period is consistent with that during NH SSW period.

=====
Other comments:
.....

1. The abstract is not sufficiently developed.

Answer: The abstract has been developed in the revised version.

Line # [18-27]: The present paper studies the perturbations in equatorial ionization anomaly (EIA) region during the Southern Hemisphere (SH) sudden stratospheric warming (SSW) of 2002, using the location of EIA crests derived from Global Positioning System (GPS) station observations, the Total Electron Content (TEC) obtained by International GNSS Service (IGS) global ionospheric TEC map (GIMs), and the equatorial electrojet (EEJ) estimated by geomagnetic field in Asian sector. The results indicate the existence of an obvious quasi 10-day periodic oscillation in the location and TEC of northern and southern EIA crest. An eastward phase progression of quasi 10-day wave producing the SH SSW of 2002 is also identified in polar stratospheric temperature. Previous studies have shown that a strong quasi 10-day planetary wave with zonal wave numbers $s=1$ extend from the lower stratosphere to mesosphere and lower thermosphere during the SH SSW of 2002 (Palo et al., 2005). Moreover, the EEJ driven by equatorial zonal electric field exhibits quasi 10-day oscillation, suggesting the enhanced quasi-10-day planetary wave associated with SSW penetrates into the ionosphere E region and produces oscillation in EIA region through modulating the E-region electric fields. Our results reveal some newer features of ionospheric variation that have not been reported during Northern Hemisphere (NH) SSWs.

.....

2. Line 23 ‘Norther Hemisphere (SH)’, is this a typo or the authors mean both NH and SH?

Answer: This is a typo, and the ‘Northern Hemisphere (SH)’ has been changed to ‘Northern Hemisphere (NH)’.

.....

3. Line 54: ‘The researches’, change to ‘Research’.

Answer: the ‘researches’ has been changed to ‘research’.

.....

4. Unit missing in the legend of Figure 5.

Answer: The color bar number in Figure 5 is the wavelet power with no units.

.....

5. Line 166: ‘A series of studies’, which ones? Include appropriate references.

Answer: the corresponding references have been cited in the revised version.

Line # [192-194]: A series of studies have showed how the quasi 10-day planetary wave in stratosphere can penetrate into the ionosphere E region (Krüger et al., 2005; Palo et al., 2005; Chang et al., 2009).

.....

6. Line 208-221: While the observed 10-day oscillation may be ascribable to the SSW the manuscript presents insufficient evidence to justify this claim.

Answer: The connections between the 10-day periodic variation of EIA crests and SSW have been further analyzed and discussed in the introduction and discussions parts of revised manuscript. In the conclusions part, we summarize these results as follows:

Line # [238-249]: Using the location and TEC of EIA crests derived from GPS station observations and GIMs, we found a quasi 10-day periodic variability in northern and southern EIA region in Asian sector during the SH SSW of 2002. In the same time period, this quasi 10-day oscillation is also seen in the polar stratospheric temperature, which is absent and weak in Kp and F10.7 index, respectively. The SH SSW of 2002 itself is generated by quasi 10-day planetary wave. Previous studies have shown that a strong quasi 10-day planetary wave with zonal wave numbers $s=1$ extend from the lower stratosphere to mesosphere and lower thermosphere during the SH SSW of 2002 (Palo et al., 2005). Moreover, the EEJ driven by equatorial zonal electric field exhibits quasi 10-day oscillation, indicating that the upward-propagating planetary waves interacted with the tide will modify E-region electric fields, thereby altering the plasma structures through upward $E \times B$ drift, which results in the periodical variations in these ionospheric parameters in F region. These results support the suggestion that the quasi 10-day variation in EIA region should be ascribed to enhanced 10-day planetary wave in lower atmosphere associated with SSW.

.....

1 Quasi 10-day wave modulation of equatorial ionization anomaly during the Southern 2 Hemisphere stratospheric warming of 2002

3
4 Xiaohua Mo¹, Donghe Zhang^{2*}
5

6 ¹College of Science, Key Laboratory for Ionospheric Observation and Simulation, Guangxi University for
7 Nationalities, Nanning, China

8 ²Department of Geophysics, Peking University, Beijing, 100871, China
9

10 Correspondence: Donghe Zhang (zhangdh@pku.edu.cn)
11
12

13 Abstract

14 The present paper studies the perturbations in equatorial ionization anomaly (EIA) region during the
15 Southern Hemisphere (SH) sudden stratospheric warming (SSW) of 2002, using the location of EIA crests
16 derived from Global Positioning System (GPS) station observations, the Total Electron Content (TEC)
17 obtained by International GNSS Service (IGS) global ionospheric TEC map (GIMs), and the equatorial
18 electrojet (EEJ) estimated by geomagnetic field in Asian sector. The results indicate the existence of an
19 obvious quasi 10-day periodic oscillation in the location and TEC of northern and southern EIA crest. An
20 eastward phase progression of quasi 10-day wave producing the SH SSW of 2002 is also identified in polar
21 stratospheric temperature. Previous studies have shown that a strong quasi 10-day planetary wave with
22 zonal wave numbers $s=1$ extend from the lower stratosphere to mesosphere and lower thermosphere during
23 the SH SSW of 2002 (Palo et al., 2005). Moreover, the EEJ driven by equatorial zonal electric field
24 exhibits quasi 10-day oscillation, suggesting the enhanced quasi-10-day planetary wave associated with
25 SSW penetrates into the ionosphere E region and produces oscillation in EIA region through modulating
26 the E-region electric fields. Our results reveal some newer features of ionospheric variation that have not
27 been reported during Northern Hemisphere (NH) SSWs.
28
29
30

31 1. Introduction

32 Sudden stratospheric warming (SSW) is large-scale meteorological process in the polar stratosphere
33 which is characterized by rapid rise in temperatures and deceleration/reversal in the zonal mean flows
34 (Scherhag, 1952). The primary driver of SSW is thought to be a rapid growth of quasi-stationary planetary
35 wave interacting with zonal mean flow (Matsuno, 1971). Although the main processes of SSW occur in the
36 polar middle atmosphere, its effects on the ionosphere have been observed in significant changes of
37 equatorial electrojet (EEJ), vertical plasma drift, and equatorial ionization anomaly (EIA) (Vineeth et al.,
38 2007; Chau et al., 2009; Goncharenko et al., 2010; Pancheva and Mukhtarov, 2011; Jin et al., 2012).

39 Since stationary planetary waves in the Southern Hemisphere (SH) generally have smaller amplitudes
40 than in the Northern Hemisphere (NH) where orographic and thermal forcing is stronger (Andrews et al.,
41 1987), major SSWs often occur in NH. Therefore, most studies about SSW effects on the ionosphere are
42 during NH SSW period. In recent years, a great deal of research has been focused on the variation of the
43 low latitude ionosphere during SSW period in the northern hemisphere, and the quasi-16-day periodic
44 disturbance and the lunar tide characteristics have been found in some ionospheric parameters, for example,
45 EEJ, vertical plasma drifts, ionospheric electron density (Vineeth et al., 2007; Chau et al., 2009; Pedatella
46 and Forbes, 2009; Goncharenko et al., 2010). Some researchers considered that this kind of quasi-16-day
47 periodic variations is related to the enhanced planetary wave during the SSW period (Chau et al., 2009;
48 Pedatella and Forbes, 2009; Liu et al., 2010). But others believe that this kind of quasi-16-day period is
49 related to the semi-diurnal lunar tides (Fejer et al., 2010; Park et al., 2012). The direct evidence is the
50 typical SSW feature appears in some ionospheric parameters as morning enhancement and afternoon
51 decrease, in a semi-diurnal pattern that progresses to later local times within several days. However, some
52 studies believe that the local time variation characteristics are not necessarily caused by the semidiurnal
53 lunar tides, Pedatella et al. (2012) demonstrate that the phase of the semidiurnal solar tide changes in a
54 manner that makes it similar to the phase of the lunar semidiurnal tide. Besides, although many studies on
55 the variation of the low latitude ionosphere during the SSW period, the physical connection between the
56 SSW in the polar region and the featured variations in low latitude ionosphere is still not clear. Some
57 studies even consider that the SSW and the featured variation of low latitude ionosphere is co-source,
58 which is the effect of enhanced planetary waves in different regions (Stening et al., 2011).

59 In comparison, the atmospheric parameters in the mesosphere and lower thermosphere (MLT) are very
60 limited. The atmospheric variation in MLT is usually indirectly reflected through the EEJ obtained from the

61 geomagnetic field data in equatorial and low latitude region. The EEJ is driven by zonal electric field
62 which also produces EIA via upward $E \times B$ drift (fountain effect). The zonal electric field modulated by tidal
63 winds in the lower thermosphere through the E-region dynamo process is easy to be influenced by various
64 atmospheric waves, so these ionospheric variations often display similar semidiurnal pattern and 13- to
65 16-day wave signatures which have been associated with enhanced planetary wave, solar and lunar tide
66 wave during SSW period (Pedatella and Forbes, 2009; Goncharenko et al., 2010; Fejer et al., 2010; Park et
67 al., 2012).

68 In August to September 2002, three minor SSWs and a major SSW appeared in SH (Varotsos 2002;
69 Baldwin et al., 2003). There is sufficient evidence that a series of unusual atmospheric states occurred in
70 this period, i.e., planetary wave scale quasi 10-day variation (Krüger et al., 2005; Palo et al., 2005),
71 short-term semidiurnal tide variability with zonal wave number $s=1$ (Chang et al., 2009) and the winds
72 oscillation with ~ 14 -days period (Andrew et al., 2004), are all linked to the extremely large planetary wave
73 events. Although the atmospheric activity in connection with 2002 SH SSW has been well revealed in
74 observations and numerical modeling, relatively little is known about the ionosphere effects of 2002 SH
75 SSW. Recently, Olson et al. (2013) studied the equatorial electrodynamic perturbations in Peruvian sector
76 during 2002 SH SSW and found enhanced quasi 2-day fluctuations and large amplitude multi-day
77 perturbations in EEJ and vertical drifts. The research of ionospheric behavior during SH SSW periods are
78 useful for verifying the existing explanation about the origin of ionospheric perturbations during NH SSW
79 periods and revealing some newer features of ionospheric variation, so further investigation of 2002 SH
80 SSW effect on ionosphere with more ionospheric parameters is still warranted.

81 In the present study, we present the first observational evidence of quasi 10-day oscillation in EIA
82 region during 2002 SH SSW which has not been reported during NH SSWs, based on the location of EIA
83 crests derived from Global Positioning System (GPS) station observations, the Total Electron Content
84 (TEC) obtained by International GNSS Service (IGS) global ionospheric TEC map (GIMs), and the EEJ
85 estimated by geomagnetic field in Asian sector.

86

87 **2. Data and Methods**

88 The location of EIA crests derived from GPS observations are used to analyze the variation in EIA
89 region during 2002 SH SSW from July 20, 2002 to October 27, 2002. The GPS stations are GUAN
90 (23.19°N, 113.34°E, MLAT \sim 12.52°N) and BAKO (6.49°S, 106.84°E, MLAT \sim 17.18°S) which are near

91 northern and southern EIA crest, respectively. The locations of the GPS stations are shown in Figures 1.
92 Since the ionospheric vertical TEC usually reach the maximum at EIA crest, the location of EIA crest can
93 be obtained by vertical TEC values at each ionospheric penetration point (IPP), which is the intersection of
94 the line of sight and the ionospheric shell (assumed to be 400 km) (Mo et al., 2014). The relative accuracy
95 of the TEC is 0.02 total electron content unit ($1\text{TECU}=10^{16}\text{ el m}^{-2}$) (Hofmann-Wellenhof et al., 1992). The
96 sample rate of these GPS stations were 30s, so the resolution of the location of EIA crest is less than 25 km
97 (Mo et al., 2017). Figures 2a and 2b show the daily average geomagnetic latitude (MLAT) of northern and
98 southern EIA crests during 2002 SH SSW.

99 The TEC from GIMs are also used to analyze the variation in EIA region. The GIMs provides maps of
100 TEC obtained from a global network of GPS receivers, which have temporal resolution of 2 hours and
101 spatial resolution of 5° in longitude and 2.5° in latitude (Mannucci et al., 1998). The EIA crest usually
102 reaches its maximum development near 14:00 LT (Huang et al., 1989; Yeh et al., 2001), so the daily
103 average TEC obtained by GIMs at 12~14 LT, $\pm 5^\circ\sim\pm 15^\circ$ MLAT, $100^\circ\sim 150^\circ\text{E}$ every day in Asian sector are
104 used to describe the variation in northern and southern EIA region, the results are shown in Figures 2c and
105 2d.

106 To demonstrate the dynamical process in EIA region, the EEJ is also used in this study, which can be
107 estimated by the difference between the horizontal component of geomagnetic field at TIR (8.7°N , 77.8°E ,
108 $\text{MLAT}\sim 0.03^\circ\text{N}$) and VSK (17.68°N , 83.32°E , $\text{MLAT}\sim 8.56^\circ\text{N}$) (Rastogi et al., 1990). The results are shown
109 in Figures 2e. In addition, the polar stratospheric temperature (90°S , 10hPa) and zonal mean zonal winds
110 (60°S , 10hPa) obtained from National Centers for Environment Prediction (NCEP) are used to examine the
111 extent of the SSW, the results are shown in Figures 2f and 2g. The background of geomagnetic activity
112 index (Kp) and solar flux index (F10.7) from the websites <http://spidr.ngdc.noaa.gov/> are depicted in
113 Figures 2h and 2i.

114 **3. Results and Analysis**

115 It can be seen from Figures 2f and 2g that there were three obvious minor SH SSW events around day
116 number 230-260 and a major SH SSW event around day number 263-288 (Olson et al., 2013). Figure 3
117 shows the contour map of polar stratospheric temperature (80°S , 10hPa) obtained from NCEP from July 20,
118 2002 to October 27, 2002. An eastward phase progression of quasi 10-day wave is clearly observed around
119 day number 210-270. With SABER temperature data, Palo et al. (2005) also observed similar disturbance
120 and suggested it consists of an eastward-propagating quasi 10-day wave with zonal wave numbers $s=1$

121 superimposed upon a large stationary planetary wave with $s=1$.

122 Now we examine the impact of this quasi 10-day wave on EIA region. 2002 was at solar maximum
123 phase, the ionosphere maybe exhibit some variations caused by periodic solar irradiance and high speed
124 solar streams related to solar rotation during 2002 SH SSW event, for example, 27-day periodic variation.

125 To exclude these long period fluctuations in EIA region, the periods longer than 15 days in the MLAT
126 location and TEC of EIA crest, and EEJ are removed. Specifically, these parameters are subtracted from
127 their respective 15-day moving average. The residuals are subjected to Lomb-Scargle (L-S) spectral
128 analysis (Lomb,1976; Scargle, 1982), and the results are shown in Figures 4a, 4b, 4c, 4d, and 4e. The
129 horizontal dashed lines represent the 95% confidence level. It is evident that the MLAT location and TEC
130 of EIA crest, and EEJ all exhibit significant quasi 10-day periodic component, which exceed or approach
131 95% confidence level, suggesting that the whole dynamical process in EIA region is modulated by quasi
132 10-day wave. Figures 4f and 4g show the L-S spectral analysis of Kp and F10.7. It can be seen that both Kp
133 and F10.7 do not exhibit quasi 10-day periodic component, indicating that variation in the solar irradiance
134 and geomagnetic activity cannot account for this quasi 10-day oscillation in EIA region.

135 To investigate the time evolution of quasi 10-day periodic variation, the Morlet wavelet spectral
136 analysis is applied to MLAT location and TEC of EIA crest, EEJ and Kp. The periods longer than 15 days
137 in the MLAT location and TEC of EIA crest , and EEJ are removed before the wavelet spectra is generated,
138 and the results are illustrated in Figures 5a, 5b, 5c, 5d, and 5e. The black solid contours in each panel
139 indicate a significance level higher than 95%. The white line in each panel represents the cone of influence
140 of the wavelet analysis. The color bar number is the power strength for each parameter. Obviously, the most
141 predominant periodic component in the MLAT location and TEC of EIA crest, and EEJ are quasi 10-day
142 period, which mainly appeared around day number 210-290, indicating quasi 10-day oscillations in EIA
143 region go through three minor SSWs and a major SSW period. The time evolution of the power in MLAT
144 location and TEC of northern EIA crest match well those of southern EIA crest, respectively. In addition,
145 we note both the MLAT location and the TEC of EIA crest show the quasi 2-day oscillations during major
146 SSW period (around day number 260-270), which are also found on equatorial ionospheric electric fields
147 and currents at the same period (Olson et al., 2013). Figure 5f shows the wavelet spectral analysis of Kp
148 index. It can be seen that quasi 10-day periodic component is nearly absent in Kp around day number
149 230-290, suggesting that magnetic activity should not be the driving force for this quasi 10-day oscillation
150 in EIA region.

151 In order to demonstrate the phase relationship of the quasi 10-day oscillations between northern and
152 southern EIA crests, the band-pass filter is performed on the MLAT location and TEC of EIA crest. The
153 absolute values of the MLAT location of EIA crest are used. The band-pass filter is centered at the period of
154 10-day, with half-power points at 8-day and 12-day, and the results are shown in Figure 6. The quasi 10-day
155 wave amplitudes of northern and southern EIA crests are roughly equivalent, which exceed 1.7 degree for
156 MLAT location and 7 TECU for TEC, respectively. Although the quasi 10-day wave of northern EIA crest
157 match well those of southern EIA crest, the wave of northern EIA crest seemed to delay behind southern
158 EIA crest, especially for MLAT location. To further verify this, Figure 7 shows the cross-correlation of
159 quasi 10-day waves in MLAT location (a) and TEC (b) between northern and southern EIA crests. The
160 cross-correlation coefficients of MLAT location and TEC reach 0.8 and 0.93, respectively. Moreover, the
161 maximum cross-correlation coefficients for MLAT location is at 1 day, indicating that the wave of northern
162 EIA crest delay 1 day behind southern EIA crest. This phase difference between northern and southern EIA
163 crests may be due to differences in longitude between two GPS stations.

164

165 **4. Discussions**

166 In recent years a series of reports have focused on ionospheric perturbations during SSW event. The
167 most predominant features in low latitude ionosphere associated with SSW event are semidiurnal pattern
168 and 13- to 16-day periodic variations, which are attributed to nonlinear interaction of planetary wave, solar
169 and lunar tide wave (Pedatella and Forbes, 2009;Goncharenko et al., 2010; Fejer et al., 2010;Park et al.,
170 2012). As major SSW often occurs in NH, most studies about SSW effects on the ionosphere are during
171 NH SSW period. In August to September 2002, the first major SSW was observed in SH. The NH and SH
172 SSW occurred in Arctic and Antarctic winter, respectively, so the occurring time and location of SH SSW
173 are opposite to those of NH SSW. The researches of ionospheric behavior during SH SSW periods are
174 useful for testing the general rule of ionospheric perturbations during NH SSW periods. For example,
175 Olson et al. (2013) demonstrated that multi-day ionospheric perturbations responding to 2002 SH SSW
176 resemble those observed during NH SSWs and these ionospheric perturbations were associated with
177 enhanced lunar tidal effects.

178 In this study we present observations of quasi 10-day oscillation in EIA region during the 2002 SH
179 SSW that has not been reported during NH SSWs. This quasi 10-day periodic component is absent or very
180 weak in Kp and F10.7 index, indicating that the magnetic activity and solar irradiance cannot account for

181 this quasi 10-day oscillation in EIA region. Meanwhile, an unusual atmospheric state occurred in this
182 period that the ozone hole over the Antarctic has a smaller size and splits into two separate holes (Varotsos
183 2002; Baldwin et al., 2003). This phenomenon is thought to be due to high temperatures in the Antarctic
184 stratosphere, which was contributed to by upward propagation of planetary waves (Venkat Ratnam et al.,
185 2004). Moreover, strong planetary wave scale quasi 10-day variation was observed in polar stratospheric
186 temperature during this period. The wave interactions between eastward-propagating waves with periods
187 near 10 days, quasi-stationary planetary waves, and the zonal mean atmospheric state were eventually
188 driven towards total break-down of the polar vortex and a major warming of the stratosphere (Krüger et al.,
189 2005; Palo et al., 2005). So the quasi 10-day oscillations in EIA region should be ascribed to atmosphere
190 perturbations linking the SSW in the Southern Hemisphere.

191 The coupling process of 10-day oscillation between the lower atmosphere and ionosphere can be
192 demonstrated by existing observations and simulations. A series of studies have showed how the quasi
193 10-day planetary wave in stratosphere can penetrate into the ionosphere E region (Krüger et al., 2005; Palo
194 et al., 2005; Chang et al., 2009). Krüger et al. (2005) revealed the eastward-traveling waves with periods
195 near 10 days and their interaction with quasi-stationary planetary waves forced in the troposphere during
196 2002 SH SSW event, supporting the observational and numerical evidence that the eastward traveling wave
197 interacts with the stationary wave to produce a quasi-periodic amplitude modulation of the stationary waves
198 (Hirota et al., 1990; Ushimaru and Tanaka, 1992). Palo et al. (2005) found an eastward-propagating quasi
199 10-day wave with zonal wave numbers $s=1$ and $s=2$, and a quasi-stationary planetary waves with $s=1$
200 extend from the lower stratosphere to the 100-120 km height region with little amplitude attenuation. While
201 the quasi-stationary planetary wave is confined to the high latitude atmosphere and cannot directly
202 propagate to equatorial ionosphere, the tides were introduced into planetary wave modulation mechanism.
203 Eswaraiah et al. (2018) reported that zonal diurnal and semidiurnal tide amplitudes in Antarctica
204 mesosphere and lower thermosphere were enhanced around day number 230-290 during 2002 SH SSW,
205 which coincides with the enhanced period of quasi 10-day oscillations in EIA region shown in Figure 5.
206 Moreover, Chang et al. (2009) showed that the short-term variability of the $s=1$ semidiurnal tide is strongly
207 dependent upon the PW1 events (quasi-10-day wave) prior to the major warming during 2002 SH SSW,
208 supporting the suggestion that the quasi-stationary planetary wave can influence migrating and
209 nonmigrating solar tides globally (Liu et al., 2010; Pedatella and Forbes, 2010). So the interactions between
210 quasi-10-day planetary wave and tide will modify the ionosphere E-region winds, which can produce

211 E-region electric fields via the E-region dynamo process. In this study, the EEJ driven by equatorial zonal
212 electric field also exhibits quasi 10-day oscillation, indicating that the upward-propagating planetary waves
213 interacted with the tide produced oscillation in EIA region through modulating E-region electric fields.
214 Specifically, the E-region electric fields map to lower ionospheric F-region along the magnetic field lines
215 and generate an eastward electric field (Goncharenko et al., 2010). At the magnetic equator, the eastward
216 electric field with quasi 10-day periodic variation change electron density distribution in the low-latitude
217 region via $E \times B$ drift, and finally leads to quasi 10-day planetary waves characteristic variations in EIA
218 region. Previous studies have revealed a strong correlation between ionospheric perturbations and the
219 occurrence of NH SSW. During NH SSW period, quasi 16-day oscillations and semidiurnal pattern are
220 observed in equatorial mesopause temperature, the MLT meridional and zonal wind, EEJ, electron density
221 and TEC (Vineeth et al., 2007; Pedatella and Forbes, 2009; Park et al., 2012 ; Jonah et al., 2014). Some
222 researchers attribute these ionospheric perturbations to the strong dynamical coupling between the lower
223 atmosphere and ionosphere through the intensification of planetary wave activity (Chau et al., 2009), lunar
224 (Fejer et al., 2010) and solar (Pedatella et al., 2012) tide. In this study, the consistent quasi 10-day
225 oscillations appear in EEJ, the location and TEC of northern and southern EIA crest, indicating that
226 coupling mechanism between the lower atmosphere and ionosphere during SH SSW period is consistent
227 with that during NH SSW period.

228 In our prior studies, a 14- to 15-day wave during several NH SSW events is ascribed to lunar tide
229 (Mo et al., 2018). So the source of quasi 10-day oscillations in EIA region during 2002 SH SSW is different
230 from 14- to 15-day waves during NH SSW. For this 10-day periodic event, it seems that the effect of the
231 planetary wave is more obvious. Moreover, no obvious 14- to 15-day oscillation is found in EIA region
232 during 2002 SH SSW, which may be that the equatorial lunar semidiurnal effects during
233 September-October are weaker than that during January-February (Stening et al., 2011; Pedatella, 2014).
234 Olson et al. (2013) also reported that the perturbations amplitude of EEJ and vertical drifts modulated by
235 lunar semidiurnal tides during SH SSW are smaller than those during NH SSW.

236

237 5. Conclusions

238 Using the location and TEC of EIA crests derived from GPS station observations and GIMs, we found
239 a quasi 10-day periodic variability in northern and southern EIA region in Asian sector during the SH SSW
240 of 2002. In the same time period, this quasi 10-day oscillation is also seen in the polar stratospheric

241 temperature, which is absent and weak in Kp and F10.7 index, respectively. The SH SSW of 2002 itself is
242 generated by quasi 10-day planetary wave. Previous studies have shown that a strong quasi 10-day
243 planetary wave with zonal wave numbers $s=1$ extend from the lower stratosphere to mesosphere and lower
244 thermosphere during the SH SSW of 2002 (Palo et al., 2005). Moreover, the EEJ driven by equatorial zonal
245 electric field exhibits quasi 10-day oscillation, indicating that the upward-propagating planetary waves
246 interacted with the tide will modify E-region electric fields, thereby altering the plasma structures through
247 upward $E \times B$ drift, which results in the periodical variations in these ionospheric parameters in F region.
248 These results support the suggestion that the quasi 10-day variation in EIA region should be ascribed to
249 enhanced 10-day planetary wave in lower atmosphere associated with SSW.

250 **Acknowledgements:** The GPS data were from the Crustal Movement Observation Network of China (via
251 e-mail to yglyang@cma.gov.cn) and IGS (available at <http://sopac.ucsd.edu>). The geomagnetic data at
252 TIR and VSK were from WDC for Geomagnetism, Kyoto (available
253 at <http://wdc.kugi.kyoto-u.ac.jp/hyplt/index.html>). The GIMs were downloaded from the site
254 <ftp://cddis.gsfc.nasa.gov>. This research was jointly supported by the National Natural Science Foundation
255 of China (41864006, 41674157, and 41464006), Guangxi Natural Science Foundation
256 (2016GXNSFAA380132), and Chinese Meridian Project. We gratefully acknowledge National Center for
257 Environmental Prediction (NCEP) for providing public access to stratospheric data (available at
258 <https://www.esrl.noaa.gov/psd/data/reanalysis/>).

259 **References**

- 260 Andrews, D. G., Holton, J. R., and Leovy, C. B.: Middle Atmosphere Dynamics, Academic, San Diego,
261 Calif, 1987.
- 262 Andrew, J. D., Vincent, R. A., Murphy, D. J., Tsutsumi, M., Riggin, D. M., and Jarvis, M. J.: The
263 large-scale dynamics of the mesosphere-lower thermosphere during the Southern Hemisphere
264 stratospheric warming of 2002, *Geophys. Res. Lett.*, 31, L14102, doi:10.1029/2004GL020282, 2004
- 265 Baldwin, M., Hirooka, T., O'Neill, A., and Yoden, S.: major stratospheric warming in the Southern
266 Hemisphere in 2002: Dynamical aspects of the ozone hole split, *SPARC Newsl.*, 20, 24-26, 2003.
- 267 Chang, L. C., Palo, S. E., and Liu, H. -L.: Short-term variation of the $s=1$ nonmigrating semidiurnal tide
268 during the 2002 stratospheric sudden warming, *J. Geophys. Res.*, 114, D03109,
269 doi:10.1029/2008JD010886, 2009.
- 270 Chau, J. L., Fejer, B. G., and Goncharenko, L. P.: Quiet variability of equatorial $E \times B$ drift during a

271 sudden stratospheric warming event. *Geophys. Res. Lett.*, 36,
272 L05101, <https://doi.org/10.1029/2008GL036785>, 2009.

273 Eswaraiah, S., Kim, Y. H., Lee, J., Ratnam, M. V., and Rao, S. V. B.: Effect of Southern Hemisphere
274 sudden stratospheric warmings on Antarctica mesospheric tides: First observational study. *J. Geophys.*
275 *Res. Space Physics*, 123, 2127–2140, <https://doi.org/10.1002/2017JA024839>, 2018.

276 Fejer, B. G., Olson, M. E., Chau, J. L., Stolle, C., Lühr, H., Goncharenko, L. P., Yumoto, K., and
277 Nagatsuma, T.: Lunar-dependent equatorial ionospheric electrodynamic effects during sudden
278 stratospheric warmings, *J. Geophys. Res.*, 115, A00G03, doi:10.1029/2010JA015273, 2010.

279 Goncharenko L. P., Chau, J. L., Liu, H. -L., and Coster, A. J.: Unexpected connections between the
280 stratosphere and ionosphere, *Geophys. Res. Lett.*, 37, L10101, doi:10.1029/2010GL043125, 2010.

281 Hirota, I., Kuroi, K., and Shiotani, M.: Midwinter warmings in the Southern Hemisphere stratosphere in
282 1988, *Q. J. R. Meteorol. Soc.*, 116, 929-941, 1990.

283 Hofmann - Wellenhof, B., Lichtenegger, H., and Collins, J.: *GPS—Theory and Practice*, Springer, New
284 York, 1992.

285 Huang, Y. N., Cheng, K., and Chen, S. W.: On the equatorial anomaly of the ionospheric total electron
286 content near the northern anomaly crest region, *J. Geophys. Res.*, 94(A10), 13,515–13,525, 1989.

287 Jin, H., Miyoshi, Y., Pancheva, D., Mukhtarov, P., Fujiwara, H., and Shinagawa, H.: Response of the
288 migrating tides to the stratospheric sudden warming in 2009 and their effects on the ionosphere studied
289 by a whole atmosphere-ionosphere model GAIA with COSMIC and TIMED/SABER observations, *J.*
290 *Geophys. Res.*, 117, A10323, doi:10.1029/2012JA017650, 2012.

291 **Jonah, O. F., de Paula E. R., Kherani, E. A., Dutra, S. L. G., and Paes, R. R.: Atmospheric and ionospheric**
292 **response to sudden stratospheric warming of January 2013, *J. Geophys. Res. Space Physics*, 119,**
293 **4973–4980, doi.org/10.1002/2013JA019491, 2014.**

294 Krüger, K., Naujokat, B., and Labitzke, K.: The unusual midwinter warming in the Southern Hemisphere
295 stratosphere 2002: A comparison to Northern Hemisphere phenomena, *J. Atmos. Sci.*, 62, 603-613,
296 2005.

297 Liu, H. -L, Wang, W., Richmond, A. D., and Roble, R. G.: Ionospheric variability due to planetary waves
298 and tides for solar minimum conditions, *J. Geophys. Res.*, 115, A00G01, doi:10.1029/2009JA015188,
299 2010.

300 Lomb, N. R.: Least-squares frequency analysis of unequally spaced data, *Astrophys.Space Sci.*, 39,

301 447-462, 1976.

302 Mannucci, A. J., Wilson, B. D., Yuan, D. N., Ho, C. M., Lindqwister, U. J., and Runge, T. F.: A global
303 mapping technique for GPS derived ionospheric total electron content measurements, *Radio Sci.*, 33,
304 565-582, doi:10.1029/97RS02707, 1998.

305 Matsuno, T.: A dynamical model of the stratospheric sudden warming, *J. Atmos. Sci.*, 28, 1479-1494, 1971.

306 Mo, X. H., Zhang, D. H., Goncharenko, L. R., Hao, Y. Q., and Xiao, Z.: Quasi-16-day periodic meridional
307 movement of the equatorial ionization anomaly, *Ann. Geophys.*, 32, 121-131, 2014.

308 Mo, X. H., Zhang, D. H., Goncharenko, L. R., Zhang, S. R., Hao, Y. Q., Xiao, Z., Pei, J. Z., Yoshikawa, A.,
309 Chau, H. D.: Meridional movement of northern and southern equatorial ionization anomaly crests in the
310 East-Asian sector during 2002–2003 SSW, *Science China Earth Sciences*, 60(4), 776–785, [https://](https://doi.org/10.1007/s11430-016-0096-y)
311 doi.org/10.1007/s11430-016-0096-y, 2017.

312 Mo, X. H., and Zhang, D. H.: Lunar tidal modulation of periodic meridional movement of equatorial
313 ionization anomaly crest during sudden stratospheric warming, *J. Geophys. Res. Space Physics*, 123,
314 1488-1499. <https://doi.org/10.1002/2017JA024718>, 2018.

315 Olson, M. E., Fejer, B. G., Stolle, C., Lühr, H., and Chau, J. L.: Equatorial ionospheric electrodynamic
316 perturbations during Southern Hemisphere stratospheric warming events, *J. Geophys. Res. Space*
317 *Physics*, 118, 1190-1195, doi:10.1002/jgra.50142, 2013.

318 Palo, S. E., Forbes, J. M., Zhang, X., Russell III, J. M., Mertens, C. J., Mlynczak, M. G., Burns, G. B., Espy,
319 P. J., and Kawahara, T. D.: Planetary wave coupling from the stratosphere to the thermosphere during the
320 2002 Southern Hemisphere pre-stratwarn period, *Geophys. Res. Lett.*, 32, L23809,
321 doi:10.1029/2005GL024298, 2005.

322 Pancheva, D., and Mukhtarov, P.: Stratospheric warmings: The atmosphere-ionosphere coupling paradigm, *J.*
323 *Atmos. Sol. Terr. Phys.*, 73, 1697-1702, doi:10.1016/j.jastp.2011.03.006, 2011.

324 Park, J., Lühr, H., Kunze, M., Fejer, B. G., and Min, K. W.: Effect of sudden stratospheric warming on
325 lunar tidal modulation of the equatorial electrojet, *J. Geophys. Res.*, 117, A03306,
326 doi:10.1029/2011JA017351, 2012.

327 Pedatella, N. M., and Forbes, J. M.: Modulation of the equatorial F-region by the quasi-16 day planetary
328 wave, *Geophys. Res. Lett.*, 34, L09105, doi:10.1029/2009GL037809, 2009.

329 Pedatella, N. M., and Forbes, J. M.: Evidence for stratosphere sudden warming-ionosphere coupling due to
330 vertically propagating tides, *Geophys. Res. Lett.*, 37, L11104, doi:10.1029/2010GL043560, 2010.

331 Pedatella, N. M., Liu, H. L., and Richmond, A. D.: Atmospheric semidiurnal lunar tide climatology
332 simulated by the Whole Atmosphere Community Climate Model, *J. Geophys. Res.*, 117, A06327,
333 doi:10.1029/2012JA017792, 2012.

334 Pedatella, N. M.: Observations and simulations of the ionospheric lunar tide: Seasonal variability, *J.*
335 *Geophys. Res. Space Physics*, 119, 5800-5806, doi:10.1002/2014JA020189, 2014.

336 Rastogi, R. G., and Klobuchar, J. A. : Ionospheric electron content within the equatorial F2 layer anomaly
337 belt, *J. Geophys. Res.*, 95(A11), 19,045–19,052, <https://doi.org/10.1029/JA095iA11p19045>, 1990.

338 Scargle, J. D.: Studies in astronomical time series analysis. II. Statistical aspects of spectral analysis of
339 unevenly spaced data, *Astrophys. J.*, 263, 835-853, 1982.

340 Scherhag, R.: Die explosionsartigen Stratosphärenerwärmungen des Spätwinters 1952, *Ber. Dtsch*
341 *Wetterdienstes USZone*, 6, 51-63, 1952.

342 Stening, R. J.: Lunar tide in the equatorial electrojet in relation to stratospheric warmings, *J. Geophys. Res.*,
343 116, A12315, doi:10.1029/2011JA017047, 2011.

344 Ushimaru, S., and Tanaka, H.: A numerical study of the interaction between stationary Rossby waves and
345 eastward-traveling waves in the Southern Hemisphere stratosphere, *J. Atmos., Sci.*, 49, 1354-1373,
346 1992.


347 Varotsos, C.: The Southern Hemisphere ozone hole split in 2002, *Environ. Sci. Pollut. Res.*, 9, 375-376,
348 2002.

349 Venkat Ratnam, M., Tsuda, T., Jacobi, C., and Aoyama, Y.: Enhancement of gravity wave activity observed
350 during a major Southern Hemisphere stratospheric warming by CHAMP/GPS measurements, *Geophys.*
351 *Res. Lett.*, 31, L16101, doi:10.1029/2004GL019789, 2004.

352 Vineeth, C., Pant, T. K., Devasia, C. V., and Sridharan, R.: Atmosphere-ionosphere coupling observed over
353 the dip equatorial MLTI region through the quasi 16-day wave. *Geophys. Res. Lett.*, 34,
354 L12102, <https://doi.org/10.1029/2007GL030010>, 2007.


355 Yeh, K. C., Franke, S. J., Andreeva, E. S., and Kunitsyn, V. E.: An investigation of motions of the
356 equatorial anomaly crest, *Geophys. Res. Lett.*, 28(24), 4517-4520, doi:10.1029/2001GL013897, 2001.

357
358
359
360




361
362

Figure 1. Location of the GPS stations in Asian sector.



363
 364 **Figure 2.** The magnetic latitude (MLAT) location of (a) northern and (b) southern equatorial ionization
 365 anomaly (EIA) crest; The TEC of (c) northern and (d) southern EIA crest; the (e) equatorial electrojet (EEJ),
 366 (f) polar stratospheric temperature (at 90°S, 10hPa) and (g) zonal wind (at 60°S, 10hPa) from National
 367 Centers for Environment Prediction; the (h) Geomagnetic activity index, Kp and (i) solar flux index F10.7
 368 during the period from July 20, 2002 to October 27, 2002.




369

370 **Figure 3.** The contour map of polar stratospheric temperature (80°S, 10hPa) obtained from NCEP during


371

the same period as in Figure 2.



372

373 **Figure 4.** Lomb-Scargle periodograms of the MLAT location of (a) northern and (b) southern EIA crest, the
 374 TEC of (c) northern and (d) southern EIA crest, (e) EEJ, (f) Kp index and (g) F10.7 during the same period
 375 as in Figure 2.




376

377 **Figure 5.** The wavelet power spectra of the MLAT location of (a) northern and (b) southern EIA crest, the

378 TEC of (c) northern and (d) southern EIA crest, (e) EEJ and (f) Kp index during the same period as in

379 Figure 2. The white line in each panel represents the cone of influence of the wavelet analysis.




380

381 **Figure 6.** The band-pass filter results of the (a) MLAT location of (solid line) northern and (dash-dotted

382 line) southern EIA crest, the (b) TEC of (solid line) northern and (dash-dotted line) southern EIA crest

383

during the same period as in Figure 2.



384

385 **Figure 7.** The cross-correlation of quasi 10-day waves in MLAT location (a) and TEC (b) between northern

386

and southern EIA crest.

387