

Interactive comment on “Mirror mode physics: Amplitude limit” by Rudolf A. Treumann and Wolfgang Baumjohann

Rudolf A. Treumann and Wolfgang Baumjohann

treumannr@gmail.com

Received and published: 7 September 2019

We thank the reviewer for the favourable comments.

We would like to direct him to the General Response we have posted in the Discussion where most questions he has have been answered in the interest of all 4 Reviewers. Here some brief specific remarks on the questions raised by the report:

1. It is absolutely true: the pairs are not phase bunched. They are just all in gyration only which is sufficient for the surface current. Phase bunching cannot be achieved and is not necessary. It suffices that at zero parallel speed the surviving pairs are on a common shell. This we mean by coherence. We have said this now explicitly in the paper and weakened the expression of quasi-coherent.

C1

2. The mirror mode provides the ideal conditions for pair formation. We would believe that this is a rare case because of the bounce motion. However, it might be that in magnetic holes which also probably permit for bouncing and as well other places, say reconnection geometries or also the auroral magnetosphere (i.e. in all cases of magnetic trapping or quasi-trapping like in reconnection where the particles bounce between plasmoids) the possibility exists for this kind of pairing as it only requires reflection at mirror points with conservation of the magnetic moment, presence of waves to resonate (here ion sound, otherwise parallel propagating waves mostly of electrostatic nature). The formal expression for the potential allows for all kinds of waves but the resonance condition sets another requirement which is not easy to satisfy.

3. Yes, of course the London length could be used. However, at the present state of the theory which in these last subsections is rather speculative as it just suggests a possibility, I would dare to apply it. The factor α is purely heuristic and very uncertain yet. Before doing it one would have to solve the stability problem of the pairs which is a difficult task which we did not attack at this moment. At this time the value of α is rather large, and one would expect that it is much less because some pairs will dissolve, being replaced by new pairs such that the number will strongly fluctuate. The London length itself is very uncertain, just a rough parameter even in superconductivity theory which is certainly more precise than plasma theory at the high temperatures where for instance pressure balance is not better than roughly 10

4. Eq 7, thanks this is trivial. Slightly changed.

5. Thanks for the language corrections.

Interactive comment on Ann. Geophys. Discuss., https://doi.org/10.5194/angeo-2019-86, 2019.

C2