

1 **Ionospheric Pc1 waves during a storm recovery phase**

2 **observed by CSES**

3 Xiaochen Gou¹, Lei Li^{1*}, Yiteng Zhang^{1*}, Bin Zhou¹, Yongyong Feng¹, Bingjun Cheng¹,
4 Tero Raita³, Ji Liu¹, ZhiMa Zeren², Xuhui Shen²

5
6 ¹State Key Laboratory of Space Weather, National Space Science Center, Chinese
7 Academy of Sciences, Beijing, China;

8 ²Institute of Crustal Dynamic, China Earthquake Administration, Beijing, China;

9 ³Sodankylä Geophysical Observatory, University of Oulu, Sodankylä, Finland

11 **Abstract**

12 During the storm recovery phase on August 27, 2018, the China Seismo-
13 Electromagnetic Satellite (CSES) detected Pc1 wave activities both in the Northern and
14 Southern hemispheres in the high latitude post-midnight ionosphere with a central
15 frequency about 2 Hz. Meanwhile, the typical Pc1 waves were simultaneously
16 observed by the Sodankylä Geophysical Observatory (SGO) stations on the ground for
17 several hours. In this paper, we study the propagation characteristics and possible
18 source regions of those waves. Firstly, we find that the Pc1 waves observed by the
19 satellites exhibited mixed polarization and the wave normal is almost parallel with the
20 background magnetic field. The field-aligned Poynting fluxes point downward in both
21 hemispheres, implying the satellites are close to the wave injection regions in the
22 ionosphere at about $L=3$. Furthermore, we also find that the estimated position of the
23 plasmopause calculated by models is almost at $L=3$. Therefore, we suggest the possible
24 sources of waves are near the plasmopause, which is consistent with previous studies
25 that the outward expansion of the plasmasphere into the ring current during the
26 recovery phase of geomagnetic storms may generate electromagnetic ion cyclotron
27 (EMIC) waves and then these EMIC waves propagate along the background magnetic
28 field northward and southward to the ionosphere at about $L=3$. Additionally, the
29 ground station data show that Pc1 wave power attenuates with increasing distance
30 from $L=3$, supporting the idea that CSES observes the wave activities near the injection
31 region. The observations are unique in that the Pc1 waves are observed in the
32 ionosphere in nearly conjugate regions, where transvers Alfvén waves propagate down
33 into the ionosphere.

35 **1 Introduction**

36 Electromagnetic ion cyclotron (EMIC) waves are in the typical frequency range of 0.1–

37 5Hz which corresponds to Pc1 pulsations on the ground. Generally, in the
38 magnetosphere, EMIC wave can be excited by cyclotron instability of hot ions (1-100
39 keV) with temperature anisotropy ($T_{\perp} > T_{\parallel}$) near the Earth's magnetic equator,
40 particularly, in the region with large plasma density and weak magnetic field, such as
41 the plasmopause, ring current and plasma sheet [Cornwall et al., 1965; Erlandson et
42 al., 1993; Horne and Thorne, 1993; Anderson et al., 1996; Lin et al., 2014]. Previous
43 studies indicate that hot ion temperature anisotropy ($T_{\perp} > T_{\parallel}$) near the Earth's magnetic
44 equator can be caused by several possible mechanisms, such as plasmopause
45 expanding into ring current region during storm recovery phase [Cornwall et al., 1970;
46 Russell & Thorne, 1970], mid-energy ions penetrating into the ring current region from
47 the plasma sheet [Bossen et al., 1976], the solar wind dynamic pressure enhancement
48 or the magnetosphere compression [Olson & Lee, 1983; Anderson & Hamilton, 1993;
49 McCollough et al., 2010; Usanova et al. 2012]. Statistical results show that EMIC waves
50 are associated with increased magnetic activity and have a peak occurrence during the
51 storm recovery phase [Wentworth, 1964; Erlandson & Ukhorskiy, 2001; Bortnik et al.,
52 2008].

53

54 Generally, EMIC waves are excited at or near the Earth's magnetic equator, and then
55 propagate along the background magnetic field toward the high latitude region, can
56 penetrate into the upper ionosphere under certain conditions. The left-hand polarized
57 (LHP) Alfvén waves incident from the magnetosphere can couple to the right-hand
58 polarized (RHP) compressional, isotropic waves in the ionosphere by the anisotropic
59 ionospheric Hall currents [Fraser et al., 1975a, 1975b; Fujita and Tamao 1988]. Since
60 the wavelength of EMIC waves with frequency about 1Hz is comparable with the scale
61 size of the ionospheric minimum in the Alfvén speed, they can be trapped and ducted
62 in this region of low Alfvén speed [Lysak et al., 1999]. Thus, the EMIC waves can be
63 observed both at the low earth orbit (LEO) and on the ground as Pc1 geomagnetic
64 pulsations with different characteristics.

65

66 At ionospheric altitudes, satellite observations of Pc1 waves are usually provided by
67 the onboard magnetometers. Magsat observed Pc1 waves at an ionospheric altitude
68 of 350-550km, with both LH and RH polarizations in a latitudinally narrow (<100 km)
69 region [Iyemori and Hayashi, 1989]. In recent years, with the development of LEO
70 satellites, various statistical studies of EMIC waves have been carried out to reveal the
71 global propagation characteristics, spatial distribution, and geomagnetic dependence
72 of Pc1 waves. Since the field-aligned currents mask the Pc1 pulsations in the high
73 latitude zone, excluding data at auroral latitudes, according to the statistical analysis
74 of CHAMP satellite data during one solar cycle, Park et al. [2013] found that Pc1 waves

75 are mostly linearly polarized, having a peak occurrence at sub-auroral latitudes, and
76 weakly dependent on magnetic activity and the solar wind velocity. Similarly, the
77 Swarm data show a peak occurrence rate of Pc1 waves at middle latitude including
78 sub-auroral region. Moreover, these waves are linear polarization dominated,
79 propagating oblique to the background magnetic field, and preferably occur during the
80 late recovery phase of magnetic storms [Kim et al. 2018a].

81

82 In this paper, we report a Pc1 wave event observed by the China Seismo-
83 Electromagnetic Satellite (CSES), as well as the Swarm satellite. Based on both electric
84 and magnetic field measurements, we study the propagation characteristics and
85 possible source regions of those Pc1 waves occurring at high latitude in the Northern
86 and Southern hemisphere ionosphere during the recovery phase of the geomagnetic
87 storm on 25-28 August 2018.

88

89 **2 Data sources**

90 The China Seismo-Electromagnetic Satellite (CSES) was launched on February 2, 2018,
91 into a sun-synchronous circular orbit at an altitude of 507 km with an inclination angle
92 of 97.4°. The local time of the descending node is 14:00. We use the magnetic field
93 data from the High Precision Magnetometer (HPM) and the electric field data from the
94 Electric Field Detector (EFD) onboard CSES. HPM includes two three-components
95 fluxgate sensors to collect vector magnetic field data with a sampling rate of 60Hz, and
96 the noise of the sensors are less than 0.02nT / $\sqrt{\text{Hz}}$ @1 Hz [Zhou et al., 2018; 2019].
97 EFD consists of four spherical sensors, which can realize three-components electric
98 field detection at a broad frequency range from DC to 3.5MHz, in which the ULF band
99 provides 125Hz sampled waveform signal [Huang et al., 2018]. Swarm was launched
100 on November 22, 2013, which has three satellites (Alpha, Bravo, and Charlie) at
101 altitudes of 450 – 550 km with an inclination angle of 88° [Friis-Christensen et al.,
102 2006]. For this study, we used magnetic field data from Swarm A with a high sampling
103 rate of 50 Hz and a noise level of 0.01 nT/ $\sqrt{\text{Hz}}$ @1Hz [Merayo, 2014]. We also use the
104 Finnish pulsation magnetometers data from Sodankylä Geophysical Observatory (SGO)
105 including Sodankylä (SOD), Oulu (OUL), and Nurmijärvi (NUR) with a sampling rate of
106 40 Hz. In addition, the solar wind data of OMNI is from CDA Web. The Dst index is from
107 WDC Web and the plasmopause simulation data is from NASA CCMC Web.

108

109 **3 Observations**

110 Figure 1 shows the variation of solar wind parameters and the geomagnetic index
111 during the Pc 1 wave event in this study. The Dst index, interplanetary magnetic field,

112 solar wind speed and solar wind dynamic pressure from Aug. 25 to 29, 2018 are shown
113 from top to bottom. It can be seen that during the magnetic storm, the Dst index
114 decreased to -170 nT at 8:00 26 August. The Pc1 waves were observed by CSES and
115 Swarm between UTC 22:50 – 23:30 (marked by the black box in Figure 1) with
116 northward interplanetary magnetic field (IMF) and minor increased Dst index in the
117 magnetic storm recovery phase on Aug. 27, 2018.

118

119 **3.1 Spatial-temporal characteristics of Pc1 waves**

120

121 On Aug. 27, 2018, CSES and Swarm -A satellites passed through the ionospheric Pc1
122 wave regions for three times, in the Northern and Southern hemispheres, marked by
123 squares (CSES) and triangles (Swarm) in Figure 2. Firstly, at around UTC 23:00 (local
124 time about 02:06 to 02:34), Swarm -A and CSES satellites successively observed Pc1
125 waves in the Southern hemisphere at geomagnetic latitude about $56^{\circ} \text{S} \sim 53^{\circ} \text{S}$ with L
126 shell value about $3.0 \sim 3.4$. The distance between the two satellites is about 300km.
127 Swarm -A observed the Pc1 waves at about UTC 22:50 (QD-LAT= 56°S , $L=3.4$) about 10
128 minutes before CSES, with a maximum amplitude about 12 nT and a central frequency
129 about 2 Hz, lasting for 1 minute, as shown in Figure 3. Then, CSES observed the Pc1
130 wave at UTC 23:02 (QD-LAT= 54°S , $L=3.1$) by the HPM Magnetometer (shown in Figure
131 4), with a maximum amplitude about 1.5 nT and a central frequency about 2 Hz, lasting
132 a minute and a half. Thereafter, at about UTC 23:30 (local time about 01:27 to 01:22),
133 the CSES flew away to the Northern hemisphere, passing through the Pc1 wave region
134 again at geomagnetic latitudes about 54°N , L values about 3.1. As shown in Figure 5,
135 the maximum amplitude is about 10 nT and the central frequency is about 2 Hz, with
136 a duration about 1 minute. Around this time, since the Swarm satellite was about 6000
137 km northeast of the CSES satellite, no Pc1 waves were observed by Swarm.

138

139 At the same time, the typical Pc1 waves were also observed by the SGO stations on
140 the ground for several hours. As shown in Figure 6, from UTC 21:35 to 24:00, SGO
141 stations recorded continuous pulsations with a central frequency of about 2-3 Hz. In
142 Figure 6, from top to bottom are the observations from SGO stations: Sodankylä (SOD;
143 $L = 5.3$, 64.3°N , 105.6°E , QD), Oulu (OUL; $L = 4.5$, 61.9°N , 104.1°E , QD), and Nurmijärvi
144 (NUR; $L = 3.4$, 57.1°N , 101.2°E , QD) from $\sim 21:00$ to 24:00 UT. The wave power of Pc1
145 pulsations increases monotonically with the decrease of L shell values of SGO stations,
146 with the maximum power at NUR station, which is close to the region where CSES
147 observed Pc1 in the Northern hemisphere. Because of the ducting effect of Pc1 waves
148 in the ionospheric waveguide, Pc1 waves are likely to be seen at a long distance away
149 from the source region [e.g., Fujita and Taomao, 1988; Kim et al., 2010]. Since the

150 boundary of the waveguide is not a perfect conductor, some absorption may happen
151 when waves propagate in the waveguide, resulting in attenuation of the wave power.
152 So, comparing the wave power observed by different ground stations, it is possible to
153 infer the probable location of the wave source. Therefore, in our case, we suggest that
154 the injection source region of the Pc1 waves in the Northern hemisphere should be
155 near (QD-LAT=54 - 56°N, $L \sim 3.3$), where CSES and NUR observed the pulsations, and
156 after incidence on the ionosphere, the waves were ducted toward northeast, observed
157 by the ground stations located at higher latitudes.

158

159 **3.2 Propagation characteristics of Pc1 waves**

160 Wave polarization is another property that provides information on the wave source
161 and spatial characteristics of wave propagation. According to theoretical studies, the
162 incident LHP Alfvén waves in the ionosphere can gradually change to RHP as the waves
163 propagate in the ionosphere away from the injection region [e.g., Fujita and Taoumao
164 1986]. Close to the injection region, the polarization pattern is usually complex,
165 because the waves near the injection source are combined with incident waves and
166 ducting waves [Hayashi et al., 1981; Kim et al., 2010].

167

168 We further analyzed the propagation characteristics of Pc1 waves observed by CSES
169 and Swarm satellites in the Northern and Southern hemispheres during the magnetic
170 storm recovery phase. Firstly, we converted the magnetic field into field-aligned
171 coordinates (FAC) and then applied polarization analysis according to the method of
172 Means et al. [1972]. From top to bottom, Figs. 7a-e shows Swarm magnetic field
173 components in FAC (including perpendicular components B_r and B_a marked in blue
174 and green and the parallel component B_z marked in red), magnetic wave power
175 spectrum in perpendicular direction and parallel direction, wave normal angle (0°
176 indicates parallel propagation and 90° indicates perpendicular propagation to the
177 background magnetic field), ellipticity (positive indicates RHP and negative indicates
178 LHP). For CSES, electric components in FAC, electric wave power spectrum in
179 perpendicular direction and parallel direction, and field-aligned Poynting flux are also
180 included in Figures 8 and 9.

181

182 It can be seen from the Swarm and CSES data in the Southern (Figure 7,8) and Northern
183 hemispheres (Figure 9), that wave normal angles (Figs. 7d, 8g and 9g) predominate
184 below $\sim 20^\circ$, indicating that Pc1 waves propagated almost parallel to the background
185 magnetic field. Our result is somewhat different from the nightside observations in the
186 ionosphere by Pisa et al. (2015) and Kim et al. (2018), which show the wave normal
187 angles are scattered or have different tendency between two hemispheres. For CSES,

188 based on the HPM and EFD data, we also calculate the field-aligned Poynting flux of
189 Pc1 waves (shown by Figs. 8i and 9i), which is positive in the Northern hemisphere,
190 negative in the Southern hemisphere, indicating that Pc1 waves observed by CSES
191 propagate along the background magnetic field downward into the ionosphere in the
192 both hemispheres.

193

194 Additionally, we find that the waves have dominant perpendicular power, and the
195 parallel power (compressional power) is almost zero (shown at Figs. 7b-c, 8e-f and 9e-
196 f), which means the waves are transverse. The transverse wave is one of the
197 characteristics of the incident wave near the wave injection region [Engebretson et
198 al.,2008; Kim et al., 2010]. The transverse wave also explains why the
199 downward(upward) component in the local North-East-Down(up) coordinates has the
200 minimum wave power, as observed by satellites and ground stations (Figure 2-3, Figure
201 6). Near the injection region with a geomagnetic latitude of $\sim 55^\circ$, the dip angle of the
202 geomagnetic field is about 73° . For a transverse wave, the power projected to the
203 downward direction should be small. We further find the wave normal, electric field
204 vector, background magnetic field are almost lie in the same plane (not shown here)
205 with a deviation less than $\pm 8^\circ$, which confirms that the incident transverse wave is
206 Alfvénic.

207

208 From Figs. 7e, 8h and 9h, the ellipticity of Pc1 waves shows mixed polarization for the
209 waves detected by CSES and Swarm in both hemispheres. To check whether our
210 calculation results truly represent these wave properties, we also use Minimum and
211 Maximum Variance Analysis (MVA) [Sonnerup & Scheible, 1998] to get the MVA
212 hodograph and the wave normal direction (not shown here), which are also consistent
213 with current results. Therefore, it seems that all the Pc1 waves observed by CSES and
214 Swarm have mixed ellipticities and propagate along the background magnetic field.

215

216 **4 Discussion**

217 In 1970, Cornwall et al. proposed that during storm recovery phase, the plasmopause
218 expanding into the ring current region can excite EMIC wave. Through simulation,
219 Horne and Thorne et al. [1993] found that the growth rate of EMIC wave inside the
220 plasmopause is obviously lower than that outside the plasmopause, and its peak is
221 near the plasmopause.

222

223 To identify the source of the Pc1 waves observed by CSES and Swarm, we use the
224 dynamic plasmasphere model from the Community Coordinated Modeling Center
225 (CCMC) [Pierrard et al., 2008] to obtain the variation of the position of the

226 plasmopause during this magnetic storm on August 26, 2018 (as shown in Figure 10).
 227 The dots correspond to the position of the plasmopause and the red star represents
 228 the conjugate location of Pc1 waves observed by CSES in the Southern hemisphere.
 229 From 11 to 21 MLT there is a plume rotating with the plasmasphere in the eastward
 230 direction. Such plumes are mostly formed during geomagnetic storm recovery phase
 231 [Pierrard and Cabrera, 2005]. Meanwhile, the simulation cycle of dynamic
 232 plasmasphere model always start at 02 MLT because the plasmas are unstable at post-
 233 midnight since the convection electric field has the largest value. Additionally, the
 234 simulation does not stop after one full cycle at 02 MLT but continues farther up to 05
 235 MLT showing two plasmopause branches between 02 MLT and 05 MLT and the gaps
 236 are caused by the loss of some of the plasma elements at large Kp jumps [Verbanac et
 237 al., 2018; Bandic et al., 2019]. Results show that the plasmopause moves outward at
 238 about UTC 23:00 on August 27, and the L value reaches about 3 near local time 02:00.
 239 Moreover, based on the formula in Carpenter and Anderson [1992] (shown as
 240 equation 1), the position of the plasmopause is estimated at about L=2.98. Therefore,
 241 we suggest that the possible sources of Pc1 waves are nearly located at the
 242 plasmopause, and this is consistent with previous studies, that the outward expansion
 243 of the plasmasphere into the ring current during the recovery phase of geomagnetic
 244 storms may generate EMIC waves, which propagate along the background magnetic
 245 field to the ionosphere, and be observed by multi-ground stations [Wentworth, 1964;
 246 Cornwall et al., 1970; Russell & Thorne, 1970].

$$247 \quad \hat{L}_{pp} = 5.6 - 0.46 \times \max_{-24, -4} K_p \quad (1)$$

248 According to the wave analysis performed using CSES and Swarm data, together with
 249 ground station observations, we suggest that the satellites are close to the wave
 250 injection regions in the Southern and northern hemisphere, during the recovery phase
 251 of the storm. The incident waves propagate almost along the background magnetic
 252 field, as transvers Alfvén waves, which has long been predicted by theoretical studies,
 253 although direct observations are rare. However, the ellipticity of the waves shows a
 254 complex pattern, which is different from the polarizations of EMIC waves (LHP) in the
 255 magnetosphere found by previous works [Fraser et al., 1975a, b; Erlandson et al.,
 256 1990]. Theoretical studies predict that EMIC waves triggered near the Earth's magnetic
 257 equator propagate toward the ionosphere, changing wave characteristics such as
 258 ellipticity and wave normal angle when they pass through multicomponent plasma
 259 [Denton, 2018; Johnson & Cheng, 1999; Kim & Johnson, 2016]. The mixed polarization
 260 pattern observed in our case might either result from incident waves with complex
 261 polarization pattern, or be attributed to the interference between the incident wave
 262 and ducting waves in the ionospheric waveguide.

263

264 Joint magnetic field and electric field observations onboard CSES provide
265 unambiguous evidence that Pc1 waves propagate downward into the ionosphere in
266 the nearly conjugate ionospheric regions. Although the observations at north and
267 south are temporally separated by about 30 mins, it seems reasonable to infer that
268 the EMIC waves propagate northward and southward from the magnetic equatorial
269 region simultaneously, and wave reflection from the ionosphere is insignificant. Our
270 result is in accord with the CRRES satellite measurements reported by Loto'aniu et al.
271 (2005), which observed that outside a region of about $\pm 11^\circ$ MLAT around the equator,
272 the Poynting vectors of the EMIC waves are directed away from the equator along the
273 magnetic field lines.

274

275 Pc1 waves sometimes have repetitive wave packet structures, which have been
276 explained by a bouncing wave packet model [e.g., Jacobs and Watanabe, 1964].
277 According to this model, a wave packet triggered in the equatorial region travels along
278 the magnetic field line, and is reflected between conjugate hemispheres. The Poynting
279 vector is an important parameter for establishing the propagation direction of wave
280 packet energy. CSES observations of Poynting vector in the ionospheric do not seem
281 to support this model.

282

283 **5 Conclusion**

284 In this paper, using the simultaneous observations from CSES and Swarm satellites and
285 the ground geomagnetic stations data, we investigated the typical Pc1 waves in the
286 Northern and Southern ionospheric hemispheres. Our principal results are as follows.

287 1. During the storm recovery phase on Aug. 27, 2018, the typical Pc1 waves were
288 recorded by the SGO stations on the ground for several hours. Meanwhile, the Pc1
289 waves were detected by the China Seismo-Electromagnetic Satellite (CSES) and Swarm
290 both in Northern and Southern hemispheres in the high latitude post-midnight
291 ionosphere region with a central frequency about 2 Hz.

292

293 2. In the field-aligned coordinate system, the power spectrum, ellipticity and normal
294 wave angle, Poynting vector are analyzed. Results show that the satellites observed
295 transverse Alfvén waves with mixed polarizations, propagating almost parallel to the
296 background magnetic field downward, which imply the satellites were close to the
297 wave injection region in the ionosphere at about $L=3$. Attenuation of Pc1 wave power
298 at ground stations with increasing distance from $L=3$ also supports the idea that CSES
299 observes the wave activity near the injection region.

300

301 3. Furthermore, it is also found that the position of the plasmapause calculated by the

302 CCMC model and the equation of Carpenter and Anderson is almost at $L=3$. Therefore,
303 we suggest the possible sources of waves are near the plasmopause, which is
304 consistent with previous studies that the outward expansion of the plasmasphere into
305 the ring current during the recovery phase of geomagnetic storms may generate
306 electromagnetic ion cyclotron (EMIC) waves. Downward pointing Poynting fluxes
307 measured by CSES at nearly conjugate hemispheres suggest EMIC waves propagate
308 northward and southward simultaneously to the ionosphere at about $L=3$.

309

310 **Acknowledgments**

311 The work is supported by NSFC grant 41904147, National Key Research and
312 Development Programs of Ministry of Science and Technology of the People's Republic
313 of China (MOST) (2016YBF0501503, 2018YFC1503501). This research made use of the
314 data from CSES mission, a project funded by China National Space Administration
315 (CNSA) and China Earthquake Administration (CEA). Additionally, thanks to CSES teams
316 for providing CSES HPM and EFD data from website link <http://www.leos.ac.cn/>, ESA
317 Swarm teams for providing Swarm FGM data from the <ftp://Swarm-diss.eo.esa.int/>,
318 and the PI of pulsation magnetometers from Sodankylä Geophysical Observatory(SGO)
319 for providing the geomagnetic pulsation data(<https://www.sgo.fi/Data/Pulsation/>),
320 and NASA CDA Web (<https://cdaweb.sci.gsfc.nasa.gov/index.html/>) for providing the
321 OMNI solar wind and magnetic field data, and NASA CCMC Web
322 (<https://ccmc.gsfc.nasa.gov/>) for providing the plasmopause simulation data.

323

324 **References**

325 Anderson, B. J., Takahashi, K., Erlandson, R. E., & Zanetti, L. J.: Pc1 pulsations observed
326 by AMPTE/CCE in the Earth's outer magnetosphere, *Geophysical Research Letters*,
327 17(11), 1853–1856, <https://doi.org/10.1029/gl017i011p01853>, 1990.

328 Anderson, B. J., & Hamilton, D. C.: Electromagnetic ion cyclotron waves stimulated by
329 modest magnetospheric compressions, *Journal of Geophysical Research*, 98(A7),
330 11369, <https://doi.org/10.1029/93ja00605>, 1993.

331 Anderson, B. J., Denton, R. E., Ho, G., Hamilton, D. C., Fuselier, S. A., and Strangeway,
332 R. J.: Observational test of local proton cyclotron instability in the Earth's
333 magnetosphere, *Journal of Geophysical Research: Space Physics*, 101(A10), 21527–
334 21543, <https://doi.org/10.1029/96ja01251>, 1996.

335 Bandić, M., Verbanac, G., & Pierrard, V.: Relationship between global plasmopause
336 characteristics and plasmopause structures in the frame of interchange instability
337 mechanism. *Journal of Geophysical Research: Space Physics*.
338 <https://doi:10.1029/2019ja026768>, 2019.

339 Bortnik, J., Cutler, J. W., Dunson, C., Bleier, T. E., & McPherron, R. L.: Characteristics of
340 low-latitude Pc1 pulsations during geomagnetic storms, *Journal of Geophysical*
341 *Research: Space Physics*, 113(A4), <https://doi.org/10.1029/2007ja012867>, 2008.

342 Bossen, M., McPherron, R. L., & Russell, C. T.: A statistical study of Pc 1 magnetic
343 pulsations at synchronous orbit, *Journal of Geophysical Research*, 81(34), 6083–6091,
344 <https://doi.org/10.1029/ja081i034p06083>, 1976.

345 Carpenter, D. L., & Anderson, R. R.: An ISEE/whistler model of equatorial electron
346 density in the magnetosphere, *Journal of Geophysical Research*, 97(A2), 1097,
347 <https://doi.org/10.1029/91ja01548>, 1992.

348 Cornwall, J. M.: Cyclotron instabilities and electromagnetic emission in the ultralow
349 frequency and very low frequency ranges, *Journal of Geophysical Research*, 70(1), 61–
350 69, <https://doi.org/10.1029/jz070i001p00061>, 1965.

351 Cornwall, J. M., Coroniti, F. V. and Thorne, R. M.: Turbulent loss of ring current protons,
352 *J. Geophys. Res.*, 75(25), 4699–4709, <https://doi.org/10.1029/JA075i025p04699>, 1970.

353 Denton, R. E.: Electromagnetic Ion Cyclotron Wave fields in a Realistic Dipole Field,
354 *Journal of Geophysical Research: Space Physics*, 123(2), 1208–1223,
355 <https://doi.org/10.1002/2017ja024886>, 2018.

356 Engebretson, M. J., Posch, J. L., Westerman, A. M., Otto, N. J., Slavin, J. A., Le, G.,
357 Strangeway, R. J. and Lessard, M. R.: Temporal and spatial characteristics of Pc1 waves
358 observed by ST5, *Journal of Geophysical Research: Space Physics*, 113(A7),
359 <https://doi.org/10.1029/2008ja013145>, 2008.

360 Erlandson, R. E., Aggson, T. L., Hoge, W. R., & Slavin, J. A.: Simultaneous observations
361 of subauroral electron temperature enhancements and electromagnetic ion cyclotron
362 waves, *Geophysical Research Letters*, 20(16), 1723–1726,
363 <https://doi.org/10.1029/93gl01975>, 1993.

364 Fraser, B. J.: Polarization of Pc 1 pulsations at high and middle latitudes, *Journal of*
365 *Geophysical Research*, 80(19), 2797–2807, [doi:10.1029/ja080i019p02797](https://doi.org/10.1029/ja080i019p02797), 1975a.

366 Fraser, B. J.: Ionospheric duct propagation and Pc 1 pulsation sources, *Journal of*
367 *Geophysical Research*, 80(19), 2790–2796, <https://doi.org/10.1029/ja080i019p02790>,
368 1975b.

369 Friis-Christensen, E., Lühr, H. and Hulot, G.: Swarm: A constellation to study the Earth's
370 magnetic field. *Earth Planet Sp* 58, 351–358, <https://doi.org/10.1186/BF03351933>,
371 2006.

372 Fujita, S., and Tamao, T.: Duct propagation of hydromagnetic waves in the upper
373 ionosphere, 1, Electromagnetic field disturbances in high latitudes associated with
374 localized incidence of a shear Alfvén wave. *Journal of Geophysical Research*, 93(A12),
375 14665. <https://doi.org/10.1029/ja093ia12p14665>, 1988.

376 Hayashi, K., Kokubun, S., Oguti, T., Tsuruda, K., Machida, S., Kitamura, T., Saka, O.,

377 Watanabe, T.: The extent of Pc 1 source region in high latitudes, *Canadian Journal of*
378 *Physics*, 59(8), 1097–1105, <https://doi.org/10.1139/p81-145>, 1981.

379 Horne, R. B., & Thorne, R. M.: On the preferred source location for the convective
380 amplification of ion cyclotron waves, *Journal of Geophysical Research*, 98(A6), 9233,
381 <https://doi.org/10.1029/92ja02972>, 1993.

382 Huang, J., Lei, J., Li, S., Zeren, Z., Li, C., Zhu, X., Yu, W.: The Electric Field Detector (EFD)
383 onboard the ZH-1 satellite and first observational results, *Earth and Planetary Physics*,
384 2(6), 469–478, <https://doi.org/10.26464/epp2018045>, 2018.

385 Iyemori, T., and Hayashi, K.: PC 1 micropulsations observed by Magsat in the
386 ionospheric F region, *Journal of Geophysical Research*, 94(A1), 93,
387 <https://doi.org/10.1029/ja094ia01p00093>, 1989.

388 Jacobs, J.A., Watanabe, T.: Micropulsation whistlers, *Journal of Atmospheric and*
389 *Terrestrial Physics* 26, 825–829, [https://doi.org/10.1016/0021-9169\(64\)90180-1](https://doi.org/10.1016/0021-9169(64)90180-1), 1964.

390 Johnson, J. R., & Cheng, C. Z.: Can Ion Cyclotron Waves Propagate to the Ground?
391 *Geophysical Research Letters*, 26(6), 671–674, <https://doi.org/10.1029/1999gl900074>,
392 1999.

393 Kim, H., Lessard, M. R., Engebretson, M. J., Lühr, H.: Ducting characteristics of Pc 1
394 waves at high latitudes on the ground and in space, *Journal of Geophysical Research:*
395 *Space Physics*, 115(A9), <https://doi.org/10.1029/2010ja015323>, 2010.

396 Kim, E.-H., and Johnson, J. R.: Full-wave modeling of EMIC waves near the He⁺
397 gyrofrequency, *Geophysical Research Letters*, 43(1), 13–21,
398 <https://doi.org/10.1002/2015gl066978>, 2016.

399 Kim, H., Hwang, J., Park, J., Bortnik, J., Lee, J.: Global characteristics of electromagnetic
400 ion cyclotron waves deduced from Swarm satellites, *Journal of Geophysical Research:*
401 *Space Physics*, 123, 1325–1336, <https://doi.org/10.1002/2017JA024888>, 2018.

402 Lin, R. L., Zhang, J. C., Allen, R. C., Kistler, L. M., Mouikis, C. G., Gong, J. C., Liu, L. Q.,
403 Klecker, B., Sauvaud, J. A., Dunlop, M. W.: Testing linear theory of EMIC waves in the
404 inner magnetosphere: Cluster observations, *Journal of Geophysical Research: Space*
405 *Physics*, 119(2), 1004–1027, <http://doi.org/10.1002/2013ja019541>, 2014.

406 Loto'aniu, T. M.: Propagation of electromagnetic ion cyclotron wave energy in the
407 magnetosphere, *Journal of Geophysical Research*, 110(A7).
408 <https://doi.org/10.1029/2004ja010816>, 2005.

409 Lysak, R. L: Propagation of Alfvén waves through the ionosphere: Dependence on
410 ionospheric parameters, *J. Geophys. Res.*, 104, 10,017,
411 <https://doi.org/10.1029/1999JA900024>, 1999.

412 McCollough, J. P., Elkington, S. R., Usanova, M. E., Mann, I. R., Baker, D. N., Kale, Z. C.:
413 Physical mechanisms of compressional EMIC wave growth, *Journal of Geophysical*
414 *Research: Space Physics*, 115(A10), <https://doi.org/10.1029/2010ja015393>, 2010.

415 Means, J. D.: Use of the three-dimensional covariance matrix in analyzing the
416 polarization properties of plane waves, *Journal of Geophysical Research*, 77(28), 5551–
417 5559, <https://doi.org/10.1029/ja077i028p05551>, 1972.

418 Merayo, J. M. G.: The Swarm Vector Field Magnetometer (VFM): instrument
419 commissioning & performance assessment, 3rd Swarm Science Meeting, Copenhagen,
420 Denmark, 19-20 June 2014.

421 Olson, J. V., & Lee, L. C.: Pc1 wave generation by sudden impulses, *Planetary and Space
422 Science*, 31(3), 295–302, [https://doi.org/10.1016/0032-0633\(83\)90079-x](https://doi.org/10.1016/0032-0633(83)90079-x), 1983.

423 Park, J., Lühr, H., and Rauberg, J.: Global characteristics of Pc1 magnetic pulsations
424 during solar cycle 23 deduced from CHAMP data, *Annales de Geophysique*, 31(9),
425 1507–1520, <https://doi.org/10.5194/angeo-31-1507-2013>, 2013.

426 Pierrard, V., and Stegen: A three-dimensional dynamic kinetic model of the
427 plasmasphere, *J. Geophys. Res.*, 113, A10209, <https://doi.org/10.1029/2008JA013060>,
428 2008.

429 Píša, D., Parrot, M., Santolík, O. and Menietti, J. D.: EMIC waves observed by the low-
430 altitude satellite DEMETER during the November 2004 magnetic storm, *J. Geophys.
431 Res. Space Physics*, 120, 5455–5464, <https://doi.org/10.1002/2014JA020233>, 2015.

432 Russell, C. T., and Thorne, R. M.: on the structure of the inner magnetosphere, *Cosmic
433 Electrodynamics*, Vol. 1, p. 67 – 89, 1970.

434 Sonnerup, B. U. O., & Scheible, M.: Minimum and maximum variance analysis, Chap.
435 1. In G. Paschmann & P. W. Daly (Eds.), *Analysis methods for multi-spacecraft data*, No.
436 SR-001 in *ISSI Scientific Reports* (pp. 185–220). Noordwijk, Netherlands: ESA
437 Publications Division, 1998.

438 Usanova, M. E., Mann, I. R., Bortnik, J., Shao, L., Angelopoulos, V.: THEMIS observations
439 of electromagnetic ion cyclotron wave occurrence: Dependence on AE, SYMH, and
440 solar wind dynamic pressure, *Journal of Geophysical Research: Space Physics*,
441 117(A10), <https://doi.org/10.1029/2012ja018049>, 2012.

442 Wentworth, R. C.: Enhancement of hydromagnetic emissions after geomagnetic
443 storms, *J. Geophys. Res.*, 69(11), 2291–2298,
444 <https://doi.org/10.1029/JZ069i011p02291>, 1964.


445 Verbanac, G., Bandić, M., Pierrard, V., & Cho, J.: MLT Plasmapause Characteristics:
446 Comparison Between THEMIS Observations and Numerical Simulations. *Journal of
447 Geophysical Research: Space Physics*. <https://doi:10.1002/2017ja024573>, 2018.

448 Zhou, B., Yang, Y. Y., Zhang, Y. T., Gou, X. C., Cheng, B. J., Wang, J. D., and Li, L.: Magnetic
449 field data processing methods of the China Seismo-Electromagnetic Satellite, *Earth
450 Planet. Phys.*, 2(6), 455–461, <https://doi.org/10.26464/epp2018043>, 2018.

451 Zhou, B., Cheng, B. J., Gou, X.C., Li, L., Zhang, Y. T., Wang, J. D., Magnes, W., Lammegger,
452 R., Pollinger, A., Ellmeier, M., Xiao, Q., Zhu, X. H., Yua, S. G., Yang, Y. Y. and Shen, X. H.:

453 First in-orbit results of the vector magnetic field measurement of the High Precision
 454 Magnetometer onboard the China Seismo-Electromagnetic Satellite, Earth Planets
 455 Space 71, 119, <https://doi.org/10.1186/s40623-019-1098-3>, 2019.


456



457

458


459 Figure 1. The solar wind conditions and geomagnetic index from Aug. 25 to 29, 2018.
 460 From top to bottom: Dst index, interplanetary magnetic field, solar wind speed and
 461 solar wind dynamic pressure, respectively. The occurrence of Pc1 waves is marked by
 462 the black box.




463

464 Figure 2. The locations of Pc1 waves observed by CSES (squares) and Swarm (triangles)
 465 satellites. The pentagram, rhombus and circle represent three the SGO stations:


466 Nurmijärvi (NUR; $L = 3.4$, 57.1°N , 101.2°E , QD), Oulu (OUL; $L = 4.5$, 61.9°N , 104.1°E ,
 467 QD), and Sodankylä (SOD; $L = 5.3$, 64.3°N , 105.6°E , QD), respectively. The black dotted
 468 and solid lines denote the trajectories of CSES and Swarm-A satellites, respectively and
 469 the red arrows represents three Pc1 wave observations.



470 2018-08-27 UTC 22:49:20 22:50:00 22:50:40 22:51:20 22:52:00
 471 Figure 3. The power spectral densities (PSDs) of the magnetic fields in the local North-
 472 East-Down coordinates during the Pc1 wave period (UTC 22:50-22:51) observed by
 473 Swarm -A.



474 2018-08-27 UTC 23:00:40 23:01:20 23:02:00 23:02:40 23:03:20
 475 Figure 4. The power spectral densities (PSDs) of the magnetic fields in the local North-
 476 East-Down coordinates during the Pc1 wave period (UTC 23:01-23:02) observed by
 477 CSES.




478

479


480

481

Figure 5. The power spectral densities (PSDs) of the magnetic fields in the local North-East-Down coordinates during the Pc1 wave period (UTC 23:30-23:31) observed by CSES.




482
 483 Figure 6. The power spectral densities (PSDs) of the magnetic fields in the local North-
 484 East-Up coordinates during the Pc1 wave period (UTC 21:35-24:00) observed by SGO
 485 ground stations at different L shell values.



486


487 Figure 7. The wave propagation and polarization features of the Pc1 waves observed
 488 by Swarm. From top to bottom, (a) magnetic field components (including
 489 perpendicular components Ba and Br marked in blue and green, parallel component
 490 Bz marked in red), (b) wave power spectrum in perpendicular and (c) parallel directions,
 491 (d) wave normal angle and (e) ellipticity computed by wave vector analysis of Means
 492 [1972]. (positive indicates right-handed polarization and negative indicates left-
 493 handed polarization).



494


495 Figure 8. The wave propagation and polarization features of the Pc1 waves observed
 496 by CSES in the Southern hemisphere. From top to bottom, (a) electric field components
 497 (including perpendicular components E_a and E_r marked in blue and green, parallel
 498 component E_z marked in red), (b) electric wave power spectrum in perpendicular and

499 (c) parallel directions; (d) magnetic field components (including perpendicular
 500 components Ba and Br marked in blue and green, parallel component Bz marked in
 501 rad), (e) wave power spectrum in perpendicular and (f) parallel directions, (g) magnetic
 502 wave normal angle and (h) ellipticity, (i) the field-aligned Poynting fluxes.



503

504 Figure 9. The wave propagation and polarization features of the Pc1 waves observed
505 by CSES in the Northern hemisphere, same format as Figure 8.
506



507
508 Figure 10. The Kp index (upper) and the simulated plasmopause location (lower) in the
509 geomagnetic equatorial plane marked by blue dots at UTC 23:00 on August 27, 2018
510 from CCMC Web. The red star represents the conjugate location of Pc1 waves observed
511 by CSES in the Southern hemisphere.