

# Estimating the maximum of smoothed highest 3-hourly *aa* index in 3 days by the preceding minimum for the solar cycle

Zhanle Du<sup>1,2</sup>

<sup>1</sup>Key Laboratory of Solar Activity, National Astronomical Observatories, Chinese Academy of Sciences, Beijing 100101, People's Republic of China

<sup>2</sup>University of Chinese Academy of Science, Beijing, People's Republic of China

Correspondence to: Z. L. Du (zldu@nao.cas.cn)

## Abstract

Predicting the maximum intensity of geomagnetic activity for an upcoming solar cycle is important in space weather service and for planning future space missions. This study analyzed the highest/lowest 3-hourly  $aa$  index ( $aa_H/aa_L$ ) in 3-day-interval, smoothed by 363 days to analyze their variation with the 11-year solar cycle. It is found that the maximum of  $aa_H$  ( $aa_{Hmax}$ ) is well correlated with the preceding minimum of either  $aa_H$  ( $aa_{Hmin}$ ,  $r = 0.85$ ) or  $aa_L$  ( $aa_{Lmin}$ ,  $r = 0.89$ ) for the solar cycle. Based on these relationships, the intensity of  $aa_{Hmax}$  for solar cycle 25 is estimated to be  $aa_{Hmax}(25) = 83.7 \pm 6.9$  (nT), about 29% stronger than that of solar cycle 24. This value is equivalent to the  $Ap$  index of  $Ap_{max}(25) = 47.4 \pm 4.4$  (nT) if employing the high correlation between  $Ap$  and  $aa$  ( $r = 0.93$ ). The maximum of  $aa_L$  ( $aa_{Lmax}$ ) is also well correlated with the preceding  $aa_{Hmin}$  ( $r = 0.80$ ). The maximum amplitude of the sunspot cycle ( $R_m$ ) is much better correlated with the high geomagnetic activity ( $aa_{Hmax}$ ,  $r = 0.79$ ) than with the low one ( $aa_{Lmax}$ ,  $r = 0.37$ ). The rise time from  $aa_{Hmin}$  to  $aa_{Hmax}$  is weakly anti-correlated to the following  $aa_{Hmax}$  ( $r = -0.42$ ). Similar correlations are also found for the 13-month smoothed monthly mean  $aa$  index. These results are expected to be useful in understanding the geomagnetic activity intensity of solar cycle 25.

## 1 Introduction

Studying and predicting geomagnetic activity are important in both geophysics and space weather. Severe geomagnetic activity may cause intense geomagnetic storms (Gonzalez et al., 1989, 1994; Chen et al., 2019), leading to disruptions in communication and deviations of orbital motions of satellites (Yoshida and Yamagishi, 2010; Petrovay, 2020). Now, the old solar cycle 24 being over, satellite and spacecraft-related departments want to know the maximum intensity of both solar and geomagnetic activity in the new solar cycle 25 for planning future space missions.

Among various indices to quantitatively describe geomagnetic activity, the  $aa$  index

(Mayaud, 1972), derived from the 3-hourly K index at two near-antipodal midlatitude stations in England and Australia, is the longest time series (since 1868) and has been widely used to analyze the long-term trend of global geomagnetic activity (Russell and Mulligan, 1995; Marat et al., 2017; Du, 2011b; El-Borie et al., 2019) and its correlation with both climate change (Cliver et al., 1998; Dobrica et al., 2009; Gavrilyeva et al., 2017) and solar activity (Echer et al., 2004; Prestes et al., 2006; Lukianova et al., 2009; Du, 2011a,c; Du and Wang, 2012; Singh and et al., 2019). The minimum  $aa$  index ( $aa_{\min}$ ), at or near the minimum of the solar cycle, has been widely used to predict the maximum amplitude of the sunspot cycle ( $R_m$ ), the so-called Ohl's precursor method (Brown and Williams, 1969; Ohl and Ohl, 1979; Du et al., 2009). But it is seldom used to directly predict the maximum  $aa$  index ( $aa_{\max}$ ) of an ensuing solar cycle.

The planetary geomagnetic index  $A_p$  (available since 1932, Bartels, 1963), derived from the average of the measurements at 13 observatories around the globe, is a daily measure of the response of geomagnetic field to variations in the interplanetary magnetic field (IMF) and the solar wind (Li, 1997; McPherron, 1999; Tsurutani et al., 2006). It is the main global magnetic index forecasted by government agencies (McPherron, 1999). Most works on forecasting the geomagnetic activity have been over short intervals, on the order of hours or days (McPherron, 1999; Abunina et al., 2013). In the earlier years, Kane (1988) pointed out that it is impossible to forecast the long-term geomagnetic activity through analyzing the daily, monthly and annual  $A_p$  and  $aa$  indices. Gordon (2015) demonstrated that the long-term geomagnetic activity can only be predicted to within a limited threshold of accuracy due to the irregular trends and cycles in the annual data and nonlinear variability in the monthly series, through analyzing the  $aa$  index.

In this study, we analyze the relationship between the maximum  $aa$  index and its preceding minimum for the 11-year solar cycle. The data and parameters used in this study are described in Sect. 2. We find out the highest/lowest 3-hourly  $aa$  index ( $aa_H/aa_L$ ) in each 3 days' interval, smoothed by 363 days (121 points) to analyze their variation with the solar cycle. In Sect. 3, it is found that the maximum of  $aa_H$  ( $aa_{H\max}$ )

is well correlated with the preceding minimum of either  $aa_H$  ( $aa_{Hmin}$ ) or  $aa_L$  ( $aa_{Lmin}$ ) for the solar cycle (Sect. 3.1), which can be used to estimate the intensity of  $aa_{Hmax}$  for solar cycle 25. The maximum of  $aa_L$  ( $aa_{Lmax}$ ) is also found to be well correlated with the preceding  $aa_{Hmin}$  (Sect. 3.2). The rise time from  $aa_{Hmin}$  to  $aa_{Hmax}$  is only weakly anti-correlated to the following  $aa_{Hmax}$  (Sect. 3.3). Using the relationship between  $A_p$  and  $aa$ , the maximum intensity of the  $A_p$  index for solar cycle 25 is estimated (Sect. 3.4). Similar correlations are analyzed using the 13-month smoothed (with half weight at the two ends) monthly mean  $aa$  index (Sect. 4). Some conclusions are discussed and summarized in Sect. 5.


## 2 Data

We use the 3-hourly  $aa$  index (in unit of nT) since 1 January 1868 (updated to 10 October 2020) from the International Service of Geomagnetic Indices (ISGI, [isgi.unistra.fr/](http://isgi.unistra.fr/)). We find out the highest/lowest  $aa$  index ( $aa_H/aa_L$ ) from 24 values of the 3-hourly  $aa$  index in each 3 days' interval, as shown in Fig. 1. The values of  $aa_H$  are in the range [5, 715] nT and those of  $aa_L$  are in the range [2, 45] nT.

It is seen in the figure that both  $aa_H$  and  $aa_L$  vary dramatically: the average absolute differences of the adjacent values are 36.7 nT and 2.5 nT, respectively. It is hard to see the variation in  $aa_H$  or  $aa_L$  with the solar cycle. Especially, most (63%) values of  $aa_L$  are the minimum (2 nT). In order to analyze the long-term trend of  $aa$  with the solar cycle conventionally represented by the 13-month smoothed monthly mean sunspot number, both  $aa_H$  and  $aa_L$  are smoothed by  $w = 121$  points (363 days) using the following running smoothing technique,


$$\bar{x}(i) = \frac{1}{w} \sum_{j=-(w-1)/2}^{(w-1)/2} x(i+j), \quad i = (w-1)/2, \dots, N_0 - (w-1)/2, \quad (1)$$

for  $x = aa_H$  and  $aa_L$  ( $N_0 = 18600$ ). There are  $N = N_0 - (w-1) = 18480$  data points left after removing the  $(w-1)$  incomplete-smooth points at the two ends of the series, as



**Fig. 1.** (a) The highest ( $aa_H$ ) and (b) lowest ( $aa_L$ ) 3-hourly  $aa$  index in each 3 days.

shown in Fig. 2 (solid). The smoothed monthly mean ( $w = 13$  months with half weight



**Fig. 2.** (a) The highest ( $aa_H$ ) and (b) lowest ( $aa_L$ ) 3-hourly  $aa$  index in each 3 days (black solid), smoothed by 121 points (363 days). The numbers in the figure represent the 11-year solar cycles. The upper dashed and lower dash-dotted lines indicate the maxima and minima, respectively, for the solar cycle. The red dotted line shows the 13-month smoothed monthly mean sunspot number ( $R_t$ ) for comparison.

at the two ends) International sunspot number series ( $R_i$ , Clette et al., 2016),

$$R_i(i) = \frac{1}{12} \left( \frac{R(i-6)}{2} + \sum_{j=-5}^5 R(i+j) + \frac{R(i+6)}{2} \right), \quad (2)$$

of the second version ([www.sidc.be/silso/datafiles](http://www.sidc.be/silso/datafiles)) is used for comparison (dotted), where  $R$  represents the (non-smoothed) monthly value (Petrovay, 2020). The smooth-  
width is selected to be  $w = 121$  points (363 days) as it is close to that (1 year = 12  
months) used in the smoothing of  $R_i$ .

In the upper panel of the figure, the dashed (dash-dotted) line indicates the maximum  
(minimum) of  $aa_H$ ,  $aa_{Hmax}$  ( $aa_{Hmin}$ ). In the lower panel, the dashed (dash-dotted) line  
indicates the maximum (minimum) of  $aa_L$ ,  $aa_{Lmax}$  ( $aa_{Lmin}$ ). The value of  $aa_{Hmax}$  ( $aa_{Lmax}$ )  
is the maximum of  $aa_H$  ( $aa_L$ ) during the time period between two adjacent solar cycle  
minima ( $R_{min}$ ) determined by the smoothed monthly mean  $R_i$ . The value of  $aa_{Hmin}$   
( $aa_{Lmin}$ ) is the minimum of  $aa_H$  ( $aa_L$ ) during the time period between two adjacent  
 $aa_{Hmax}$  ( $aa_{Lmax}$ ). These parameters are displayed in Table 1, in which  $T_r$  is the rise time  
from  $aa_{Hmin}$  to  $aa_{Hmax}$  and  $R_m$  the maximum of  $R_i$  for the 11-year solar cycle. The last  
row denotes the averages of the parameters.

### 3 Result

The correlation coefficients between the parameters in Table 1 are listed in Table 2  
for comparison. It is seen in Table 2 that  $R_m$  is well correlated with  $aa_{Hmin}$  ( $r = 0.84$ ),  
 $aa_{Hmax}$  ( $r = 0.79$ ),  $aa_{Lmin}$  ( $r = 0.81$ ), and positively correlated with  $aa_{Lmax}$  ( $r = 0.37$ ). This  
implies that the stronger the solar activity ( $R_i$ ), the higher the geomagnetic activity ( $aa$ ).  
But the maximum amplitude of sunspot cycle ( $R_m$ ) is much better correlated with the  
high geomagnetic activity ( $aa_{Hmax}$ ,  $r = 0.79$ ) than with the low one ( $aa_{Lmax}$ ,  $r = 0.37$ ),  
implying that the low geomagnetic activity depends less on the solar activity than the  
high one does. The correlation between  $R_m$  and  $aa_{Hmin}$  ( $aa_{Lmin}$ ),  $r = 0.84$  (0.81), is

**Table 1.** The minimum ( $aa_{Hmin}$ ) and maximum ( $aa_{Hmax}$ ) of 363-day smoothed highest 3-hourly  $aa$  index ( $aa_H$ ) in each 3 days, the rise time from  $aa_{Hmin}$  to  $aa_{Hmax}$  ( $T_r$ ), the minimum ( $aa_{Lmin}$ ) and maximum ( $aa_{Lmax}$ ) of 363-day smoothed lowest 3-hourly  $aa$  index ( $aa_L$ ) in each 3 days, and the maximum ( $R_m$ ) of 13-month smoothed monthly mean  $R_1$  for solar cycles 11-25.

$n$ (unit)	$aa_{Hmin}$ (nT)	$aa_{Hmax}$ (nT)	$T_r$ (year)	$aa_{Lmin}$ (nT)	$aa_{Lmax}$ (nT)	$R_m$
11		88.24			4.29	234.0
12	21.41	79.60	3.81	2.00	4.59	124.4
13	31.88	86.92	2.08	2.58	4.47	146.5
14	19.41	59.54	8.47	2.00	4.37	107.1
15	25.31	73.28	5.72	2.19	5.80	175.7
16	30.80	75.90	5.56	2.07	7.56	130.2
17	41.60	84.01	4.01	2.52	6.36	198.6
18	51.13	88.31	6.57	3.03	7.18	218.7
19	49.60	100.99	5.89	3.08	7.66	285.0
20	43.08	79.31	7.54	2.55	8.78	156.6
21	49.35	97.87	2.63	3.52	7.96	232.9
22	48.93	103.89	4.78	3.57	9.73	212.5
23	47.84	99.75	5.92	2.97	10.22	180.3
24	25.02	64.81	5.73	2.17	5.27	116.4
25	29.57			2.94		
Av.	36.78	84.46	5.29	2.66	6.73	179.9

related to the Ohl's precursor method (Ohl and Ohl, 1979) for predicting  $R_m$ . Some other correlations are analyzed to estimate  $aa_{Hmax}$  (Sect. 3.1),  $aa_{Lmax}$  (Sect. 3.2), and  $T_r$  (Sect. 3.3).



**Table 2.** Correlation coefficients between the parameters in Table 1.

$r$	$aa_{Hmin}$	$aa_{Hmax}$	$T_r$	$aa_{Lmin}$	$aa_{Lmax}$	$R_m$
$aa_{Hmin}$	1.00	0.85	-0.10	0.84	0.80	0.84
$aa_{Hmax}$	0.85	1.00	-0.42	0.89	0.63	0.79
$T_r$	-0.10	-0.42	1.00	-0.28	0.13	-0.18
$aa_{Lmin}$	0.84	0.89	-0.28	1.00	0.70	0.81
$aa_{Lmax}$	0.80	0.63	0.13	0.70	1.00	0.37

### 3.1 Relationship between $aa_{Hmax}$ and its preceding $aa_{Hmin}/aa_{Lmin}$


One can see in Table 2 that  $aa_{Hmax}$  is well correlated with its preceding  $aa_{Hmin}$  ( $r = 0.85$ ) and  $aa_{Lmin}$  ( $r = 0.89$ ), as shown in Fig. 3 for the scatter plots of  $aa_{Hmax}$  against  $aa_{Hmin}$  (a) and  $aa_{Lmin}$  (b). The solid lines represent the linear fits of  $aa_{Hmax}$  to  $aa_{Hmin}$  and  $aa_{Lmin}$  with the least-squares-fit regression equations given by

$$\begin{aligned} aa_{Hmax} &= 47.1 \pm 7.1 + (0.99 \pm 0.18)aa_{Hmin}, \sigma = 7.3, \\ aa_{Hmax} &= 25.3 \pm 9.5 + (22.3 \pm 3.5)aa_{Lmin}, \sigma = 6.5, \end{aligned} \quad (3)$$

where  $\pm$  indicates the  $1-\sigma$  deviation of the fitting coefficient and  $\sigma$  is the standard deviation of the regression.

Based on the above relationships, the minimum  $aa$  index (around the solar minimum) can be used as an indicator to estimate the following maximum. One can estimate  $aa_{Hmax}$  for solar cycle 25 by substituting the values of  $aa_{Hmin}$  (29.57 nT) and  $aa_{Lmin}$  (2.94 nT) into the above equations,

$$\begin{aligned} aa_{Hmax1}(25) &= 76.4 \pm 7.3(\text{nT}), \text{ from } aa_{Hmin}, \\ aa_{Hmax2}(25) &= 91.0 \pm 6.5(\text{nT}), \text{ from } aa_{Lmin}, \end{aligned} \quad (4)$$



**Fig. 3.** Scatter plots of  $aa_{Hmax}$  against  $aa_{Hmin}$  (a) and  $aa_{Lmin}$  (b).

(labelled by asterisks). Since these values are derived by the fitting equations (Eq. (3)) with similar correlation coefficients (0.85 and 0.89), we take their average,

$$\begin{aligned}
 aa_{Hmax}(25) &= \frac{1}{2}[aa_{Hmax1}(25) + aa_{Hmax2}(25)] \\
 &= 83.7 \pm 6.9(\text{nT}),
 \end{aligned}
 \tag{5}$$

as an estimate of  $aa_{Hmax}(25)$ . It implies that the 363-day smoothed highest 3-hourly  $aa$  index in 3-day-interval during the maximum period of solar cycle 25 is estimated to be close to the average (84.46 nT) over the past cycles (Table 1), but higher than that (64.81 nT) of solar cycle 24 by about 29.2%.

It should be pointed out that the above estimate may be an upper limit of  $aa_{Hmax}(25)$  as the values of  $aa_{Hmin}(25)$  and  $aa_{Lmin}(25)$  may not be finally determined (refer to Discussion in Sect. 5). Although we are not quite sure if the current  $aa_H$  ( $aa_L$ ), 30.69 (3.13) in April 2020, would decrease to a smaller value than that, 29.57 (2.94), used in the current work, there would not be significant variations in  $aa_{Hmin}$ ,  $aa_{Lmin}$ , and the above estimate, because solar cycle 25 (in term of the smoothed monthly mean  $R_l$ ) has already started (December 2019) a few months.

### 3.2 Relationship between $aa_{Lmax}$ and the preceding $aa_{Hmin}$


One can also see in Table 2 that  $aa_{Lmax}$  is well correlated with the preceding  $aa_{Hmin}$  ( $r = 0.80$ ) or  $aa_{Lmin}$  ( $r = 0.70$ ). Fig. 4(a) shows the scatter plot of  $aa_{Lmax}$  against  $aa_{Hmin}$ . The linear fitting equation of  $aa_{Lmax}$  to  $aa_{Hmin}$  (solid) is

$$aa_{Lmax} = 2.0 \pm 1.2 + (0.131 \pm 0.030)aa_{Hmin}, \sigma = 1.2. \quad (6)$$

Substituting  $aa_{Hmin}(25) = 29.57$  (nT) into this equation, one can estimate the 363-day smoothed *lowest* 3-hourly  $aa$  index in 3-day-interval during the maximum period of solar cycle 25,  $aa_{Lmax} = 5.9 \pm 1.2$  (nT). This value is slightly lower than the average (6.73 nT) over the past cycles, but higher than that (5.27 nT) of solar cycle 24 by 12.0%.

### 3.3 Relationship between the rise time and the following maximum

Now, we analyze if the rise time of the  $aa$  geomagnetic index for the solar cycle is correlated with the following maximum so that it can be used to estimate the rise time, as the case often used in the solar (sunspot) cycle (Waldmeier, 1939).



**Fig. 4.** Scatter plots of  $aa_{Lmax}$  against  $aa_{Hmin}$  (a) and  $T_r$  against  $aa_{Hmax}$  (b).

Figure 4(b) shows the scatter plot of the rise time ( $T_r$ ) from  $aa_{Hmin}$  to  $aa_{Hmax}$  for the solar cycle against its following maximum ( $aa_{Hmax}$ ). The solid line indicates the linear

fit of  $T_r$  to  $aa_{H_{\max}}$  by

$$T_r = 9.9 \pm 3.0 - (0.055 \pm 0.036)aa_{H_{\max}}, \sigma = 1.6 \text{ (years)}. \quad (7)$$

The anti-correlation coefficient between  $T_r$  and  $aa_{H_{\max}}$ ,  $r = -0.42$  (at a confidence level of about 84%), is so weak that it can hardly be used to estimate the rise time ( $T_r$ ). If the rise time is computed from the minimum ( $R_{\min}$ ) of the sunspot cycle to  $aa_{H_{\max}}$ , the correlation is even weaker,  $r = -0.14$ .

### 3.4 Relationship between $A_p$ and $aa$

Another important index often used to evaluate geomagnetic activity is the  $A_p$  index ([www.gfz-potsdam.de/en/kp-index](http://www.gfz-potsdam.de/en/kp-index)). Similar correlations as those for the  $aa$  index can also be obtained. But the  $A_p$  index is available only from 1932 to April 2018. In this section, we employ the relationship between  $A_p$  and  $aa$  to estimate the maximum intensity of the  $A_p$  index for solar cycle 25 from the previous result. Figure 5(a) shows the scatter plot of the 363-day smoothed 3-hourly  $A_p$  against  $aa$  (dots). The solid line represents the linear fit of  $A_p$  to  $aa$  by


$$A_p = 0.12 \pm 0.01 + (0.5647 \pm 0.0005)aa, \sigma = 2.1. \quad (8)$$

It is obvious that  $A_p$  is highly correlated with  $aa$ ,  $r = 0.93$ , as they are based on the same observations.

According to this equation, the estimated  $aa_{H_{\max}}(25) = 83.7 \pm 6.9$  (nT) in Sect. 3.1 is equivalent to  $A_{p_{\max}}(25) = 47.4 \pm 3.9 \pm 2.1 = 47.4 \pm 4.4$  (nT), in which  $\pm 3.9$  is derived from the uncertainty ( $\pm 6.9$ ) of  $aa_{H_{\max}}(25)$ ,  $\pm 2.1$  is the standard deviation of the regression, and  $\sqrt{3.9^2 + 2.1^2} = 4.4$ .

## 4 Result for the 13-month smoothed monthly mean $aa$ index

At last in this section, we simply analyze the previous correlations using the 13-month smoothed monthly mean  $aa$  index, as shown in Fig. 5(b). The upper dashed and




**Fig. 5.** (a) Scatter plot of the 363-day smoothed 3-hourly  $A_p$  against  $aa$  (dots) and the linear fit (solid). (b) The 13-month smoothed monthly mean  $aa$  (solid) and  $R_1$  (dotted). The numbers in the figure indicate the solar cycles. The upper dashed and lower dash-dotted lines indicate the maximum ( $aa_{max}$ ) and minimum ( $aa_{min}$ ) of  $aa$ , respectively, for the solar cycle.

**Table 3.** Parameters of the 13-month smoothed monthly mean  $aa$  for the solar cycle.

$n$	$aa_{\min}(\text{nT})$	$aa_{\max}(\text{nT})$	$T_r(\text{month})$
11		21.10	
12	6.07	20.25	44
13	10.77	23.66	23
14	5.64	17.12	93
15	8.26	22.60	63
16	9.57	25.39	68
17	12.06	24.66	64
18	15.26	28.56	79
19	15.34	31.42	62
20	12.56	29.07	109
21	15.33	32.51	31
22	16.18	32.20	51
23	14.69	32.90	72
24	7.85	19.33	67
25	12.78		
Av.	11.60	25.77	63.5

lower dash-dotted lines indicate the maximum ( $aa_{\max}$ ) and minimum ( $aa_{\min}$ ) of the  $aa$  index, respectively, for the solar cycle. The value of  $aa_{\max}$  is the maximum of  $aa$  during the time period between two adjacent solar cycle minima ( $R_{\min}$ ) determined by the smoothed monthly mean  $R_l$  (dotted). The value of  $aa_{\min}$  is the minimum of  $aa$  during the time period between two adjacent  $aa_{\max}$ . The parameters are listed in Table 3, in which  $T_r$  is the rise time from  $aa_{\min}$  to  $aa_{\max}$ .

Figure 6(a) shows the scatter plot of  $aa_{\max}$  against  $aa_{\min}$  (triangles). The solid line



**Fig. 6.** Scatter plots of  $aa_{\max}$  against  $aa_{\min}$  (a) and  $T_r$  against  $aa_{\max}$  (b).

indicates the linear fit of  $aa_{\max}$  to  $aa_{\min}$  by

$$aa_{\max} = 10.4 \pm 1.7 + (1.36 \pm 0.14)aa_{\min}, \sigma = 1.7. \quad (9)$$

The correlation coefficient between  $aa_{\max}$  and  $aa_{\min}$  is  $r = 0.95$  (at a confidence level greater than 99%), slightly higher than that,  $r = 0.85$  (0.89), for the correlation between




$aa_{Hmax}$  and  $aa_{Hmin}$  ( $aa_{Lmin}$ ) in Fig. 3 using the 363-day smoothed highest (lowest) 3-hourly  $aa$  index in 3-day-interval.

Substituting  $aa_{min}(25) = 12.78$  into this equation, one can estimate  $aa_{max}(25) = 27.9 \pm 1.7$  (asterisk), about 44% higher than that (19.33) of solar cycle 24. This estimate is similar to the case in Sect. 3.1 that the estimate (91.0) of  $aa_{Hmax}(25)$  from  $aa_{Lmin}$  in Eq. (4) is about 40% higher than that (64.81 nT) of solar cycle 24 using the minimum of 363-day smoothed lowest 3-hourly  $aa$  index in 3-day-interval.

Figure 6(b) shows the scatter plot of the rise time ( $T_r$ ) from  $aa_{min}$  to  $aa_{max}$  against the maximum ( $aa_{max}$ ). The data points are much scattered and  $T_r$  is nearly uncorrelated to the following  $aa_{max}$ ,  $r = -0.09$ . Therefore, this correlation is unable to be used to estimate the rise time of  $aa_{max}$ .

## 5 Discussion and Conclusions

It is well known that the  $aa$  index is positively correlated with the solar activity (as represented by  $R_i$ ), since the latter is the main source of the former (Legrand and Simon, 1981; Feynman, 1982; Echer et al., 2004). In general, the stronger the solar activity, the higher the ( $aa$ ) geomagnetic activity. However the relationship between  $aa$  and  $R_i$  is not a simple linear one (Borello-Filisetti et al., 1992; Mussino et al., 1994; Kishcha et al., 1999; Lockwood et al., 1999; Echer et al., 2004; Tsurutani et al., 2006; Du, 2011a,c, 2020). The  $aa$  index tends to lag behind  $R_i$  about 2–3 years around a solar cycle maximum (Wang et al., 2000; Echer et al., 2004), and (only) about 1 year around a solar cycle minimum (Legrand and Simon, 1981; Wang and Sheeley, 2009; Du, 2011b), as indicated in Fig. 7 for the time difference ( $\Delta T_{max}$ ) of the 13-month smoothed monthly mean  $aa_{max}$  to  $R_m$  (a) and that ( $\Delta T_{min}$ ) of  $aa_{min}$  to  $R_{min}$  (b). The intensity of geomagnetic activity can only be roughly evaluated from that of solar (sunspot) activity, as the linear correlation coefficient between the smoothed monthly mean  $aa$  and  $R_i$  is only 0.61 (Du, 2011c) or even lower (0.43) if using the non-smoothed series (Du, 2011b). In addition, the future solar activity is also unknown at the current



**Fig. 7.** The time difference between  $aa_{max}$  and  $R_m$  (a) and that between  $aa_{min}$  and  $R_{min}$  (b).

time and so it can not be directly used to estimate the future geomagnetic activity.

There are many methods that can be used to predict the maximum amplitude of the sunspot cycle ( $R_m$ ), such as (i) statistical methods, employing the relationship between the inter-cycle parameters (Thompson, 1988; Hathaway et al., 1994) or the early rising rate (Thompson, 1988; Cameron and Schüssler, 2008; Du and Wang, 2012); (ii) the functional methods, using mathematical functions of a few parameters (Hathaway et al., 1994; Du, 2011d) for extrapolating the following monthly values; (iii) the geomagnetic

precursor methods (Brown and Williams, 1969; Ohl and Ohl, 1979; Du et al., 2009), using the geomagnetic activity near the solar minimum; and (iv) the solar precursor ones (Schatten et al., 1978; Pesnell and Schatten, 2018), using the previous cycle's polar field.

5 In contrast, there are less methods found to predict the maximum amplitude of geomagnetic activity index for the 11-year solar cycle. Geomagnetic activity forecast has been over the order of hours or days (McPherron, 1999; Abunina et al., 2013). The annual or monthly prediction on the geomagnetic activity is within a limited accuracy (over 20%) due to the irregular variation in the time series (McPherron, 1999; Gordon,  
10 2015). In the earlier years, Kane (1988) even pointed out that it is impossible to forecast the long-term geomagnetic activity through analyzing the time series of the  $Ap$  and  $aa$  index (refer also to Gordon, 2015). The geomagnetic activity near the solar minimum or at the decreasing phase of the solar cycle has been widely used to predict the maximum amplitude of the sunspot cycle, but was seldom used to predict the  
15 maximum amplitude of the geomagnetic activity itself.

In the current work, we analyzed the highest ( $aa_H$ ) and lowest ( $aa_L$ ) 3-hourly  $aa$  index in each 3 days' interval, smoothed by 363 days (121 points) to analyze their variation with the solar cycle represented by the 13-month smoothed monthly mean  $R_t$ . It is found that the maximum of  $aa_H$  ( $aa_{Hmax}$ ) is well correlated with the preceding minimum of either  $aa_H$  ( $aa_{Hmin}$ ,  $r = 0.85$ ) or  $aa_L$  ( $aa_{Lmin}$ ,  $r = 0.89$ ) for the 11-year solar cycle. So,  
20 these relationships can be used to estimate the maximum intensity of geomagnetic activity for the solar cycle by employing the time series itself,  $aa_{Hmax}(25) = 83.7 \pm 6.9$  (nT). It implies that the maximum intensity of geomagnetic activity for solar cycle 25 would be similar to the average over the past cycles, but higher than that of solar cycle  
25 24 by about 29.2%. Certainly, this estimate may be an upper limit, as  $aa_{Hmin}$  and  $aa_{Lmin}$  may be finally determined a few months after the solar minimum (Fig. 7).

Similar results can also be obtained if using the  $Ap$  index. However, the  $Ap$  index is available only up to April 2018. So, we employed the relationship between  $Ap$  and  $aa$  to estimate the maximum  $Ap$  index for solar cycle 25:  $Ap_{max}(25) = 47.4 \pm 4.4$  (nT). For

the 13-month smoothed monthly mean  $aa$  index, the maximum  $aa$  index ( $aa_{\max}$ ) of the solar cycle is also well correlated to the preceding minimum ( $aa_{\min}$ ), with a correlation coefficient of  $r = 0.95$ .

The well known ‘Waldmeier effect’ (Waldmeier, 1939) that the rise time of a solar cycle is well anti-correlated with the following maximum amplitude has been widely used to estimate the rise or peak time of a solar cycle if the amplitude has been predicted. However, such a correlation is very weak for the geomagnetic activity index. The rise time ( $T_r$ ) from  $aa_{H\min}$  to  $aa_{H\max}$  for the solar cycle is found to be only weakly anti-correlated to the following maximum ( $aa_{H\max}$ ),  $r = -0.42$ . This weak correlation may be related to the fact that the geomagnetic activity maximum (minimum) is not aligned to the solar (sunspot) activity maximum (minimum) in time (Fig. 7). In most cases,  $aa_{\max}$  ( $aa_{\min}$ ) lags behind  $R_m$  ( $R_{\min}$ ). But in some other cases,  $aa_{\max}$  ( $aa_{\min}$ ) precedes  $R_m$  ( $R_{\min}$ ). The weak correlation between the rise time and the following maximum of geomagnetic activity for the solar cycle can hardly be used to estimate the former.

According to the analysis above, the following conclusions may be summarized.

1. The 363-day smoothed highest ( $aa_H$ ) and lowest ( $aa_L$ ) 3-hourly  $aa$  index in 3-day-interval is analyzed, finding that the maximum of  $aa_H$  ( $aa_{H\max}$ ) is well correlated with the preceding minimum of either  $aa_H$  ( $aa_{H\min}$ ,  $r = 0.85$ ) or  $aa_L$  ( $aa_{L\min}$ ,  $r = 0.89$ ) for the 11-year solar cycle. As a result, the maximum  $aa$  index for the current solar cycle 25 is estimated to be  $aa_{H\max}(25) = 83.7 \pm 6.9$  (nT), about 29% higher than that of solar cycle 24. This value is equivalent to the  $Ap$  index of  $Ap_{\max}(25) = 47.4 \pm 4.4$  (nT) if using the relationship between  $Ap$  and  $aa$  (Eq. (8)).
2. The maximum ( $aa_{L\max}$ ) of  $aa_L$  is also found to be well correlated with the preceding  $aa_{H\min}$ ,  $r = 0.80$ . Based on this correlation,  $aa_{L\max}(25)$  is estimated to be  $5.9 \pm 1.2$  (nT), about 12% higher than that of solar cycle 24.
3. The maximum amplitude of the sunspot cycle ( $R_m$ ) is much better correlated with the high geomagnetic activity ( $aa_{H\max}$ ,  $r = 0.79$ ) than with the low one ( $aa_{L\max}$ ,  $r = 0.37$ ).

4. The rise time ( $T_r$ ) from  $aa_{Hmin}$  to  $aa_{Hmax}$  is found to be weakly anti-correlated to the following maximum ( $aa_{Hmax}$ ) for the solar cycle,  $r = -0.42$  at the 84% confidence level.
5. For the 13-month smoothed monthly mean  $aa$  index, the maximum  $aa$  index ( $aa_{max}$ ) of the solar cycle is well correlated with the preceding minimum ( $aa_{min}$ ,  $r = 0.95$ ). The rise time from  $aa_{min}$  to  $aa_{max}$  is nearly uncorrelated to the following maximum ( $r = -0.09$ ).

### Author contribution

This work is completed by Z. L. Du.

### Competing interests

The author declares that he has no conflicts of interest.

*Acknowledgements.* We are grateful to the two anonymous reviewers for valuable suggestions which improved this manuscript. This work is supported by the National Science Foundation of China (NSFC) through grants 11973058 and 11603040.

### References

- Abunina, M., Papaioannou, A., Gerontidou, M., Paschalis, P., Abunin, A., and more: Forecasting Geomagnetic Conditions in near-Earth space, *Journal of Physics: Conference Series*, 409, id. 012197, <https://doi.org/10.1088/1742-6596/409/1/012197>, 2013.
- Bartels, J.: Discussion of time variations of geomagnetic activity indices Kp and Ap 1932-1961, *Ann. Geophys.*, 19, 1–20, <https://ui.adsabs.harvard.edu/abs/1963AnG....19....1B/abstract>, 1963.

- Borello-Filisetti, O., Mussino, V., Parisi, M., and Storini, M.: Long-term variations in the geomagnetic activity level. I. A connection with solar activity, *Ann. Geophys.*, 10, 668–675, <https://ui.adsabs.harvard.edu/abs/1992AnGeo..10.668B/abstract>, 1992.
- 5 Brown, G.M., Williams, W.R.: Some properties of the day-to-day variability of Sq(H), *Planet. Space Sci.*, 17, 455, [https://doi.org/10.1016/0032-0633\(69\)90076-2](https://doi.org/10.1016/0032-0633(69)90076-2), 1969.
- Cameron, R., Schüssler, M.: A robust correlation between growth rate and amplitude of solar cycles: consequences for prediction methods, *Astrophys. J.*, 685, 1291–1296, <https://doi.org/10.1086/591079>, 2008.
- 10 Chen, S., Chai, L., Xu, K., Wei, Y., Rong, Z., and Wan, W.: Estimation of the Occurrence Probability of Extreme Geomagnetic Storms by Applying Extreme Value Theory to Aa Index, *J. Geophys. Res.: Space Physics*, 124, 9943–9952, <https://doi.org/10.1029/2019JA026947>, 2019.
- Clette, F., Cliver, E., Lefèvre, L., Svalgaard, L., Vaquero, J., Leibacher, J.: Preface to topical issue: recalibration of the sunspot number, *Solar. Phys.*, 291, 2479–2486, <https://doi.org/10.1007/s11207-016-1017-8>, 2016.
- 15 Cliver, E.W., Boriakoff, V., and Feynman, J.: Solar variability and climate change: Geomagnetic aa index and global surface temperature, *Geophys. Res. Lett.*, 25, 1035–1038, <https://doi.org/10.1029/98GL00499>, 1998.
- Dobrica, V., Demetrescu, C., Boroneant, C., and Maris, G.: Solar and geomagnetic activity effects on climate at regional and global scales: Case study-Romania, *J. Atmos. Sol. Terr. Phys.*, 71, 1727–1735, <https://doi.org/10.1016/j.jastp.2008.03.022>, 2009.
- 20 Du, Z.L.: The correlation between solar and geomagnetic activity – Part 1: Two-term decomposition of geomagnetic activity, *Ann. Geophys.*, 29, 1331–1340, <https://doi.org/10.5194/angeo-29-1331-2011>, 2011a.
- 25 Du, Z.L.: The correlation between solar and geomagnetic activity – Part 2: Long-term trends, *Ann. Geophys.*, 29, 1341–1348, <https://doi.org/10.5194/angeo-29-1341-2011>, 2011b.
- Du, Z.L.: The correlation between solar and geomagnetic activity – Part 3: An integral response model, *Ann. Geophys.*, 29, 1005–1018, <https://doi.org/10.5194/angeo-29-1005-2011>, 2011c.
- 30 Du, Z.L.: The Shape of Solar Cycle Described by a Modified Gaussian Function, *Solar. Phys.*, 273, 231–253, <https://doi.org/10.1007/s11207-011-9849-8>, 2011d.
- Du, Z.L.: The solar cycle: predicting the peak of solar cycle 25, *Astrophys. Space Sci.*, 365, 104, <https://doi.org/10.1007/s10509-020-03818-1>, 2020.

- Du, Z.L., and Wang, H.N.: The relationships of solar flares with both sunspot and geomagnetic activity, *Research in Astron. Astrophys.*, 12, 400–410, <https://doi.org/10.1088/1674-4527/12/4/004>, 2012.
- 5 Du, Z.L., Li,R., and Wang, H.N.: The Predictive Power of Ohl's Precursor Method, *Astron. J.*, 138, 1998–2001, <https://doi.org/10.1088/0004-6256/138/6/1998>, 2009.
- Du, Z.L., and Wang, H.N.: Predicting the solar maximum with the rising rate, *Science China (Physics, Mechanics & Astronomy)*, 55, 365–370, <https://doi.org/10.1007/s11433-011-4612-6>, 2012.
- 10 Echer, E., Gonzalez, W.D., Gonzalez, A.L.C., Prestes, A., Vieira, L.E.A., and more: Long-term correlation between solar and geomagnetic activity, *J. Atmos. Sol. Terr. Phys.*, 66, 1019–1025, <https://doi.org/10.1016/j.jastp.2004.03.011>, 2004.
- El-Borie, M.A., El-Taher, A.M., Thabet, A.A., and Bishara, A.A.: The impact of asymmetrical distribution of solar activity on geomagnetic indices throughout five solar activity cycles, *Adv. Spa. Res.*, 64, 278–286, <https://doi.org/10.1016/j.asr.2019.03.040>, 2019.
- 15 Feynman, J.: Geomagnetic and solar wind cycles, 1900-1975, *J. Geophys. Res.*, 87, 6153–6162, <https://doi.org/10.1029/JA087iA08p06153>, 1982.
- Gavrilyeva, G.A., Ammosov, P.P., Ammosova, A.M., Koltovskoi, I.I., and Sivtseva, V.I.: Geomagnetic activity signature in seasonal variations of mesopause temperature over Yakutia, *Proceedings of the SPIE*, 10466, id. 1046670, <https://doi.org/10.1117/12.2288710>, 2017.
- 20 Gonzalez, W.D., Joselyn, J.A., Kamide, Y., Kroehl, H.W., Rostoker, G., Tsurutani, B.T., Vasyliunas, V.M.: What is a geomagnetic storm?, *J. Geophys. Res.*, 99, 5771–5792, 1994.
- Gonzalez, W.D., Tsurutani, B.T., Gonzalez, A.L.C., Smith, E.J., Tang, F., and Akasofu, S.-I.: Solar wind-magnetosphere coupling during intense magnetic storms (1978-1979), *J. Geophys. Res.*, 94, 8835–8851, <https://doi.org/10.1029/JA094iA07p08835>, 1989.
- 25 Gordon, R.: Forecasting geomagnetic activity at monthly and annual horizons: Time series models, *J. Atmos. Sol. Terr. Phys.*, 133, 111–120, <https://doi.org/10.1016/j.jastp.2015.08.010>, 2015.
- Hathaway, D.H., Wilson, R.M., and Reichmann, E.J.: The shape of the sunspot cycle, *Solar. Phys.*, 151, 177–190, <https://doi.org/10.1007/BF00654090>, 1994.
- 30 Kane, R.P.: Forecasting geomagnetic activity, *PAGEOPH.*, 126, 85–101, <https://doi.org/10.1007/BF00876916>, 1988.
- Kishcha, P.V., Dmitrieva, I.V., and Obridko, V.N.: Long-term variations of the solar - geomagnetic correlation, total solar irradiance, and northern hemispheric temperature (1868-1997), *J. At-*

- mos. Sol. Terr. Phys., 61, 799–808, [https://doi.org/10.1016/S1364-6826\(99\)00035-8](https://doi.org/10.1016/S1364-6826(99)00035-8), 1999.
- Legrand, J.P., and Simon, P.A.: Ten cycles of solar and geomagnetic activity, Sol. Phys., 70, 173–195, <https://doi.org/10.1007/BF00154399>, 1981.
- Li, Y.: Predictions of the features for sunspot cycle 23, Sol. Phys., 170, 437–445, <https://doi.org/10.1023/A:1004963215247>, 1997.
- Lockwood, M., Stamper, R., and Wild, M.N.: A doubling of the Sun's coronal magnetic field during the past 100 years, Nature, 399, 437–439, <https://doi.org/10.1038/20867>, 1999.
- Lukianova, R., Alekseev, G., and Mursula, K.: Effects of station relocation in the aa index, J. Geophys. Res., 114, A02105, <https://doi.org/10.1029/2008JA013824>, 2009.
- Marat, D., Galina, D., Viktor, D., and Anna, D.: Dependence of the F2-layer critical frequency median at midlatitudes on geomagnetic activity, Sol. Terr. Phys., 3, 67–73, <https://doi.org/10.12737/stp-34201707>, 2017.
- Mayaud, P.: The aa indices: A 100-year series characterizing the magnetic activity, J. Geophys. Res., 77, 6870–6874, <https://doi.org/10.1029/JA077i034p06870>, 1972.
- McPherron, R.L.: Predicting the Ap index from past behavior and solar wind velocity, Physics and Chemistry of the Earth, Part C: Solar, Terrestrial & Planetary Science, 24, 45–56, [https://doi.org/10.1016/S1464-1917\(98\)00006-3](https://doi.org/10.1016/S1464-1917(98)00006-3), 1999.
- Mussino, V., Borello, F.O., Storini, M., and Nevanlinna, H.: Long-term variations in the geomagnetic activity level Part II: Ascending phases of sunspot cycles, Ann. Geophys., 12, 1071–1075, <https://ui.adsabs.harvard.edu/abs/1994AnGeo..12.1071M/abstract>, 1994.
- Ohl, A.I., Ohl, G.I.: A new method of very long-term prediction of solar activity, Solar-Terrest. Predictions Proc. NASA/MSFC, 2, 258–263, <https://ui.adsabs.harvard.edu/abs/1979stp.....2..258O/abstract>, 1979.
- Pesnell, W.D., Schatten, K.H.: An early prediction of the amplitude of Solar Cycle 25, Solar. Phys., 293, 112, <https://doi.org/10.1007/s11207-018-1330-5>, 2018.
- Petrovay, K.: Solar cycle prediction, Living Rev. Sol. Phys., 17, 2, <https://doi.org/10.1007/s41116-020-0022-z>, 2020.
- Prestes, A., Rigozo, N.R., Echer, E., and Vieira, L.E.A.: Spectral analysis of sunspot number and geomagnetic indices (1868-2001), J. Atmos. Sol. Terr. Phys., 68, 182–190, <https://doi.org/10.1016/j.jastp.2005.10.010>, 2006.
- Russell, C.T., and Mulligan, T.: The 22-year variation of geomagnetic activity: Implications for the polar magnetic field of the Sun, Geophys. Res. Lett., 22, 3287–3288, <https://doi.org/10.1029/95GL03086>, 1995.



- Schatten, K.H., Scherrer, P.H., Svalgaard, L., and Wilcox, J. M.: Using dynamo theory to predict the sunspot number during solar cycle 21, *Geophys. Res. Lett.*, 5, 411–414, <https://doi.org/10.1029/GL005i005p00411>, 1978.
- 5 Singh, P.R., Tiwari, C.M., Saxena, A.K., Agrawal, S.L.: Quasi-biennial periodicities and heliospheric modulation of geomagnetic activity during solar cycles 22C24, *Phys. Scr.*, 94, 105005, <https://doi.org/10.1088/1402-4896/ab10b6>, 2019.
- Thompson, R.J.: The rise of solar cycle number 22, *Solar. Phys.*, 117, 279–289, <https://doi.org/10.1007/BF00147249>, 1988.
- 10 Tsurutani, B.T., Gonzalez, W.D., Gonzalez, A. L.C., Guarnieri, F.L., Gopalswamy, N., and more: Corotating solar wind streams and recurrent geomagnetic activity: A review, *J. Geophys. Res.*, 111, A07S01, <https://doi.org/10.1029/2005JA011273>, 2006.
- Waldmeier, M.: Über die Struktur der Sonnenflecken, *Astron. Mitt. Zrich*, 14, 439–450, <https://ui.adsabs.harvard.edu/abs/1939MiZur..14..439W/abstract>, 1939.
- 15 Wang, Y.M., Lean, J., and Sheeley, N.R.: The long-term variation of the Sun's open magnetic flux, *Geophys. Res. Lett.*, 27, 505–508, <https://doi.org/10.1029/1999GL010744>, 2000.
- Wang, Y.M., and Sheeley, N.R.: Understanding the Geomagnetic Precursor of the Solar Cycle, *Astrophys. J.*, 694, L11–L15, <https://doi.org/10.1088/0004-637X/694/1/L11>, 2009.
- Yoshida, A., Yamagishi, H.: Predicting amplitude of solar cycle 24 based on a new precursor method, *Ann. Geophys.*, 28, 417–425, <https://doi.org/10.5194/angeo-28-417-2010>, 2010.