

Response to Referee 1

"A Comparison of Contact Charging and Impact Ionization in Low Velocity Impacts: Implications for Dust Detection in Space" as submitted by Antonsen et al. to ANGEO

We thank the referee for their time to review the manuscript, and for a number of helpful and important comments. In the following, we have tried to respond to all of them, and we have presented the revisions we have carried out in connection to the respective comments. We have not made any major revisions in the overall structure of the paper as both referees seem to agree with the structural setup. Major revisions in the content have been made, and we also refer to the open response to Referee 2. We note, especially, that we have given a better overall description of the mechanisms discussed in the paper. We have also revised the abstract.

Regarding our equations (1) and (2): We have used the wrong citation here. The correct source is John et al (1980), and this has now been corrected. We have also revised the citations in the first paragraph of section 2.2; Here we have now stated that we use the general theory of John et al. (1980) and the results of Wang and John for plastic collisions. The alpha-parameter is, as very correctly stated, not trivial to derive and we have relied on Soo's (1970) utilization of Hertzian deformation. Have added a reference to this work below the eqs.

Regarding the referees second comment (range of velocities in Mocker et al experiments and extrapolation of results): This is a very good point (raised by both referees), and is due to a poor description of our utilization; The reason why we have used Mocker et al. results is that the quality of the data is very good in their range of interest. What is not clearly stated here, is that their result is almost "indestinguishable" (for our purpose; Fe-on-Ag) from the findings of Collette et al 2014, who have investigated impacts done to speeds of 2 km/s – which is well embedded into the range of velocities we need comparison data for in order to compare contact charging and impact ionization. Therefore we have used Mocker's result, however, we could have used Collette et al. with identical conclusions. The data from both Mocker et al and Collette et al for speeds below 10 km/s is furthermore not in the volume ionization regime, which is only expected when impact energies exceed the Fermi-limit (several tens of km/s). Revision: Have tried to specify why we have chosen the Mocker et al results in the start of paragraph 2 of section 3.2.

Regarding the MUDD-results and Figure 7: A very helpful comment. The simulated current is based on high-resolution dust density and size (average size, monodisperse) data as obtained by Havnes et al (2019). This is referenced in the last paragraph of the section. Due to the very low impact velocity, for reasons discussed in section 4.2., we felt the inclusion of the Saha-Langmuir solution in figure 7 was unnecessary. In fact, even the pre-charges on the ice particles alone would completely dominate the current in MUDD in comparison to shock wave ionization, in the case of rocket. We agree with the referee that the text discussing figure 7 can be improved, and have therefore implemented a revision of this. Hopefully, the text reads better.

Minor comments:

For corrections of typos and insertion or removing of certain words, we have not listed the revision below.

- (1) Revised as suggested.
- (3) Yes, underestimate.
- (5) Defined MSP.
- (6) Revised.
- (7) Added explanation and abbreviations.

(9) Yes, there might be an additional factor there, but lowering the filling factor into the lower end around 1%, there is still a significant overestimation present.

(11) We have changed Z to capital Delta.

(25) Added descriptions of pre-charge.

(26) This was chosen due to the “common” practice, as well as it allows for direct comparison between the results. As far as the contact charging model goes, since it is specific yield (C/kg), it does not matter whether or not the incoming (modelled) projectile is 30 or say 100 nm, since – as shown in the appendix – Q_p is proportional to r^3 . Thus, the specific yield is the same for any size. A difficult question is then how the plasticity and other modelling parameters change with velocity, which can become complex to give a thought out answer to.

(28) Rephrased.

(32) The speed range of beta meteoroids is still not very well known due to the fact that we are not certain about where the formation region lies. Even at present, different authors use formation radii in the range 5-20 Solar radii, which would yield very different velocity ranges (assuming only conservation of energy and angular momentum control the orbits of the meteoroids). We hope Parker solar probe and solar orbiter will answer open questions about beta meteoroids and their velocities.

(35) It is raised to the power of 1/3 after, so $4/5 \rightarrow 4/15$.

Response to Referee 2, handle angeo-2020-23-RC2

"A Comparison of Contact Charging and Impact Ionization in Low Velocity Impacts: Implications for Dust Detection in Space" as submitted by Antonsen et al. to ANGEO

We thank the referee for a number of helpful and interesting comments. In the following, we have tried to respond to all of them, and we have presented the revisions we have carried out in connection to the respective comments. We have not made any major revisions in the overall structure of the paper, only major revisions in the content. We have also revised the abstract.

General comment 1:

We understand that the current presentation of the theory is rather complicated, and may benefit from some revision. Firstly, we strongly agree with the referee that the validity of the model should be discussed in a better manner. We believe that this would also clear up some of the confusion the reader might have. We address this issue first, below, and address the point of a high-level description as brought up by the referee after that.

1. The referee brings up an important point here – the discussion of the validity of the model over the range of parameters used. The intention of utilizing a contact charging model is that it is only really representable for the lower part of the velocity spectrum introduced in the paper (0.1 to 10 km/s). This is not the intention either. What we have failed to convey, based on the confusion brought up by the referee, is however that there *is* a change in charging mechanisms between traditional impact ionization (Langmuir-saha) and capacitive charging. To say however, exactly where the overlap between these two regimes finds place is difficult both experimentally and theoretically. In fact, we do not think there exists an answer in the literature currently that explains the sudden change ("discontinuity") between charge mechanisms at $O(1 \text{ km/s})$ impact speeds theoretically, and it is beyond the scope of this work to answer that question.

Revision: It was intended that the discussion in section 4.1 shall address the validity of an impact ionization vs. contact charging model at 'low' speeds, but we can agree that this discussion is presented too late in the work. We have therefore revised the manuscript by adding a paragraph already in the introduction noting that the current work does not address the exact limit (or speed) that a change between mechanisms happen, only that it happens within the range and that we discuss the arguments for validity of one mechanism before the other [P. 2, L. 19]. Furthermore, we feel that the introduction of section 2 introduces the details of the contact charging–impact ionization dichotomy well.

2. We feel that the essence of Section 2 provides a good high-level introduction into the problem at hand and our solution to it, however, we agree with the referee that the current presentation of the material is not optimal. We feel that a better high-level description/introduction can be given by improving Section 2.

Revision: At the very beginning of Section 2, we have added what can be considered a high-level 'simple' explanation, which we hope serves the purpose of guiding non-experts towards an understanding of the deeper topics of the paper.

General comment 2:

The referee raises another well founded and important point, which implies that the presentation or rather distinction between the two addressed modes of charging is not sufficient. To clarify, what was intended understood from the manuscript:

The paper is indeed not comparing apples to apples, as the two discussed mechanisms are fundamentally different. Also addressed in 'comment 1', the two mechanisms in fact compete. To describe the difference at a deeper level than currently, we believe will make the work considerably more complex to comprehend (and probably more speculative). There is no presence of ionization or dissociation at a fundamental level in our contact charging mechanism. Thus, the referee states correctly that the charge production is closely bound to the distribution of fragments. It is also correct that one polarity is completely dominating – which is exactly what has been consistently shown at many instances for rocket experiments and low velocity ice-on-metal collisions in the lab (e.g. referenced works by Havnes, Næsheim, Antonsen et al., Tomsic et al. and so forth). We feel it is also important to point out that the mass spectral characteristica which can be drawn out from conventional impact experiments should not be discussed in connection with our charging model. However, it is merely a tool to better understand when shock wave ionization of certain materials becomes effective/dominant – as we have utilized it in the manuscript.

Revision: In addition to the changes made in connection to general comment 1, which we believe clears some confusion; we have added at the end of the first section of the first paragraph of section 2 [p. 3] an elaboration of the fundamental differences between the discussed mechanisms. This paragraph is subsequently followed by a motivation that we now feel is easier to comprehend and should explain the essence of our modelling efforts better.

General comment 3:

As is hopefully much more clear from the responses above and corresponding revisions in the manuscript, our contact charging model not intrinsically equivalent to ionization as it happens in impacts of small particles with solids. It therefore does not make sense to discuss mass spectral properties of the impact debris of the fragmentation cloud which we provide the model for. There are two important points in the paper where we do in fact touch upon issues related to the role of ions in the impact cloud:

1. In the discussion of the overlap (in produced charge) between impact ionization and contact charging, where it makes sense to note where the literature finds onset of specific ions in impact gas mas spectra. This is interesting, because it is a rough indication of when (shock) impact ionization becomes dominant, and we consequently mention this in section 4.0. We have here relied on work done with the dust accelerator in Heidelberg, and we find the quality and amount of that data to be sufficient for our purpose.

2. In the comparison of our proposed mechanism to the low velocity shock wave ionization solution as presented by Drapatz and Michel. In that regime (i.e. below the limit where mass spectra can detect significant ion partial pressures), particularly volatile ions (e.g. of alkali metals) can diffuse through the molten fragments (or droplets, as Drapatz et al. refers to them) and can be released from their surface. The degree of ionization and produced charge can subsequently be described well with a Saha-Langmuir equation. We address this in depth in section 4.1.

We also want to stress that we do value the tremendous effort behind and results from the Cassini CDA. As the TOF mass spectral properties are not very useful for the direct comparison with our contact charging mechanism, as we have tried to convey in the manuscript, we have not focussed on these. However, from the good advice given by the referee, we understand that a mentioning of such works can easily be justified and have included a reference and description of Hillier et al CDA results in section 4.

General comment 4:

The main issue in this point was addressed in the response to general comment 1, above. We hope it is much clearer as to which regimes of validity we are interested in.

As a side note, the mid-range velocities in which shock wave ionization have been proposed to dominate in (some km/s to a few tens of km/s) have the same velocity-to-charge relationship, or rather power-law, through the whole range, i.e. $Q \sim m^x v^y$. Thus, a somewhat arbitrary upper velocity limit of 10 km/s was chosen. To exaggerate the argument: the limit could have been set to the arguably more arbitrary 9 km/s or 12 km/s – it does not really matter for the presentation of our results. From the discussion in section 4.1, we hope it is clear that we do not propose that contact charging is effective in the upper part of the investigated velocity range. The referee makes a very good point in that the Hertzian deformation theory which we assume cannot be valid at the upper higher speeds investigated here. We have tried to convey this message throughout the paper, but since it may be unclear, we have also emphasised this issue in the presentation of the fragmentation model in section 2.1.

The referee mentions the extreme that at the low end that a grain could behave like a bullet – simply deforming. This is not possible, as the elastic-plastic crossover for grains smaller than several tens of nanometres happens as much lower speeds (even <100 m/s) than interesting for this work [e.g. Rennecke and Weber 2014, Froeschke et al., Tomsic et al.]

Specific comments:

Smaller adjustments; re-phrasing, typos, reusing citations is not addressed below, however, we have tried to follow the recommendations of the referee.

Regarding Saha-solution applied in impact ionization: The referee is correct that a pure Saha-solution is not necessarily valid. What we (and originally Drapatz and Michel) utilize for the low velocity limit of shock wave ionization is the modified Saha-Langmuir equation (now revised in the abstract). This is a

Saha equation that bound electrons in metals are indeed distributed. Such a solution has been shown to hold for impurity ionization in impacts below ~ 10 km/s, and we present the basics in section 2.3.

Regarding Auer 2012: This is a newer/revised edition (?) It is the preferred citation as listed from Springer Online, at least.

Regarding contact charging: We feel this is now sufficiently answered through the revisions in connection to the general comments above.

Regarding section 2.1: A good point. The citations, as we feel can be understood from the text is used to back up the claims about the degree of fragmentation. Other references in the section help create a more rigorous literature background. An important point is also that key parts of the theoretical background/literature cited in connection to the fragmentation is discussed in section 3.1 (in the Results) where we feel it is much more natural to bring up. *Revision:* This is referenced to in section 2.1, however we have adjusted the first paragraph of section 2.1 to help the reader understand the grounds on which we have chosen modelling parameters.

Regarding Hertzian deformation: revised.

Regarding figure 1: Revised overall description in the text and in the figure caption.

Regarding binning: Misprint. Binning is done with 1 Å spacing, but the sentence was removed as it does not matter for the end result; a continuum binning (say 0.000...01 nm) would yield very similar results when the fragmenting particles is several nanometres.

Regarding section 2.3: a very good point, brought up by both referees. The reason why we have utilized Mocker et al. results is that the quality of the data is very good in their range of interest. What is not clearly stated here, is that their result is almost “indestructible” (for our purpose; Fe-on-Ag) from the findings of Collette et al 2014, who have investigated impacts done to speeds of 2 km/s – which is well embedded into the range of velocities we need comparison data for in order to compare contact charging and impact ionization. Therefore we have used Mocker’s result, however, we could have used Collette et al. with identical conclusions. The data from both Mocker et al and Collette et al for speeds below 10 km/s is furthermore not in the volume ionization regime, which is only expected when impact energies exceed the Fermi-limit (several tens of km/s). *Revision:* Have tried to specify why we have chosen the Mocker et al results in the start of paragraph 2 of section 3.2.

Regarding Ztot: Bound on fragments. Revised.

Regarding work functions: This is mentioned in the end of the discussion as one of the more uncertain parameters/assumptions, as the work function for nanoscale particles is in fact size dependent to some extent (ref e.g. Wood 81). From mesospheric studies, e.g. Plane et al., it is not necessarily such a large difference for aerosols at sizes of nanometres to tens of nanometres, that it makes a very large difference. Tens of nanometres is by rule of thumb “bulk” as far as cohesive and surface binding energies goes and thus also work function. However, as has also been mentioned, if all modelling parameters are tuned to the extremes of their tolerances, the resulting charge production could change significantly.

Regarding last comment on figure 4: Another very good point that we have had to think about. As far as the contact charging model goes, since it is specific yield (C/kg), it does not matter whether or not the incoming (modelled) projectile is 30 or say 100 nm, since – as shown in the appendix – Q_p is proportional to r^3 . Thus, the specific yield is the same for any size. A difficult question is then how the plasticity and other modelling parameters change with velocity, which can become complex to give a thought out answer to. We feel it lies a bit outside of the scope of the paper, as the current work is “only” meant to show that we need a better description for the “low velocity” range of impact charging and that our model is a good way to explain some of the observed phenomena connected to such impacts.

Tidbits:

- Charge is not quantized at this point, which may or may not overpredict the contribution from the smaller fragments. Effectively, the charging current can be viewed as a charging probability. A resolution of this ‘issue’ must also discuss in details the role of potential wells on the surface of the smallest fragments – which is probably a phenomenal modelling effort.
- MALDI-type processes are certainly interesting, although may lie a little bit outside our parameter range of interest. In general, the speeds necessary to obtain complete evaporation and have energy available to facilitate MALDI are not obtained at the impact energies we discuss here.

A Comparison of Contact Charging and Impact Ionization in Low Velocity Impacts: Implications for Dust Detection in Space

Tarjei Antonsen¹, Ingrid Mann¹, Jakub Vaverka², Libor Nouzak², and Åshild Fredriksen¹

¹Department of Physics and Technology, UiT- The Arctic University of Norway, 9037 Tromsø, Norway

² Department of Surface and Plasma Science, Faculty of Mathematics and Physics, Charles University, Prague, 180 00, Czech Republic

Correspondence: T. Antonsen (tarjei.antonsen@uit.no)

Abstract. We investigate the generation of charge ~~during collision of~~ due to collision between projectiles with sizes below $\sim 1 \mu\text{m}$ and metal surfaces at speeds ~ 0.1 to 10 kms^{-1} . This corresponds to speeds above the elastic limit and well below speeds where volume ionization can occur. ~~The conditions that we consider apply to dust particles naturally occurring in space and in Earth's upper atmosphere and their direct impacts on rockets, spacecraft, and impacts of secondary ejecta. We introduce~~

5 ~~Impact charge production at these low to intermediate speeds have traditionally been described by invoking the theory of shock wave ionization. By looking at the thermodynamics of the low velocity solution of shock wave ionization, we find that such a mechanism alone is not sufficient to account for the recorded charge production in a number of scenarios in the laboratory and in space. We propose a model of capacitive contact charging that involves no direct ionization, in which we allow for projectile fragmentation upon impact, and. Furthermore, we show that this model describes measurements of metal-metal impacts in the~~

10 ~~laboratory and in-situ measurements of dust in the Earth's atmosphere well. We have considered the utilization of our model for different scenarios in interplanetary space and in Earth's atmosphere. From this discussion we find it likely that our work can be employed in a number of situations where impact velocities are relatively small. Furthermore, we have discussed the thermodynamics of the low velocity solution of shock wave ionization, and conclude that the impurity charging effect utilized in the much used model of Drapatz and Michel (1974) does not sufficiently describe charge generation at impact speeds below~~

15 ~~a few kilometers per second. Consequently, impact charging at low speeds cannot be described with a Saha solution. also address contact charging in the context of ice-on-metal collisions and apply our results to rocket observations of mesospheric dust. In general, we find that contact charging dominates at speeds up to a few kilometres per second, and complements shock wave ionization up to speeds where direct ionization can take place. The conditions that we consider can be applied to dust particles naturally occurring in space and in Earth's upper atmosphere and their direct impacts on rockets, spacecraft, and impacts of secondary ejecta.~~

1 Introduction

The variables in any experiment studying the impact of dust grains – be it of terrestrial, meteoric, interplanetary or interstellar origin – span many orders of magnitude. By *variables* we mean the aggregation of ambient parameters and intrinsic parameters of the projectile dust grains and impact surfaces. The ambient parameters, such as neutral and charged species densities and

temperature can span several orders of magnitude. Combining this fact with the notion that the material properties of the plethora of different dust types that can produce charge in an impact process are also highly variable, there are arguably no single experiments or theoretical considerations that can give a satisfactory explanation of observed phenomena across all possible combinations of *variables*. In this paper, we focus on what we find is a gap in the knowledge about impact charge 5 production at low impact speeds, i.e. $v_p \lesssim 10 \text{ kms}^{-1}$.

Early experimental studies of impact ionization of micrometer sized grains on metal surfaces were applied in designing highly sensitive micrometeor detectors; see e.g. Auer and Sitte (1968) and Adams and Smith (1971). The mode of operation of such detectors is to measure the material specific charge generation, which in general yields a semi-empirical relation on the form: $Q \propto m^\alpha v^\beta$. This allowed for detection of grains down to sizes $\sim 100 \text{ nm}$ at speeds between the order of 1 kms^{-1} and 10 some tens of kms^{-1} . The velocity range is important here, as it has been proposed to be bounded by the limit for production of *ionization* on impact – which we shall understand here as the release of electrons and ions according to a Saha-equation. The pioneering developments in the late 60's and early 70's motivated the employment of impact particle detectors on spacecraft such as HELIOS, Galileo and Cassini which have all successfully detected cosmic dust (see e.g. Auer (2012)).

In a treatment of impact charging at speeds up to some tens of kms^{-1} , Drapatz and Michel (1974) proposed a mechanism for 15 charge generation by shock wave propagation into both projectile and impact surface. This theory is often referred to as shock wave ionization. This mechanism has been widely used in describing the charge generation in dust accelerators and spacecraft dust impacts. For typical impact speeds in many interplanetary spacecraft and laboratory experiments ($\gtrsim 10 \text{ kms}^{-1}$), the shock wave model performs very well. However, as discussed in the current work, we find that the proposed theory in its extrapolation 20 down to lower speeds of a few kms^{-1} does not describe the charge generation at low speeds sufficiently. In general, we find that it underestimates the amount of generated charge. There are several applications in the laboratory and in space (satellites, rockets and even spacecraft) that encounter low impact speeds by particles down to nanoscale size where the shock wave ionization theory thus may not be used.

Motivated by the notions above, we propose in this paper a new charging theory for impact speeds below $\sim 10 \text{ kms}^{-1}$ and 25 projectile sizes down to nanoscale. The theory utilizes the concept of contact (capacitive) charging and furthermore includes the parameterization of fragmentation of the projectile particles on impact into a distribution of smaller grains. The resulting charge generation is then dependent on the degree of fragmentation and affinity to exchange charge capacitively; the latter is dependent on the difference in work function between projectile and impact surface. In this work, we do not address the exact material specific speed limit at which the impact charge mechanism changes from contact charging to shock wave ionization. However, we address in depth the plausible velocity regime in which one or the other mechanism should dominate. A theoretical treatment of the ‘discontinuity’ in charging efficiency at speeds on the order of 1 kms^{-1} is not within the scope of the current work.

The models of charging and fragmentation that constitute our novel approach are presented in sections 2.1 and 2.2. The shock wave ionization theory is presented in 2.3. The results from comparisons of charging models in metal-metal and ice-metal collisions are presented in section 3. The thermodynamics and limitations of the shock wave ionization model at low

speeds is discussed in 4.1. We In section 4.2, we furthermore discuss the application areas of a contact charging model for low velocity impacts of dust on metal surfaces –in spacecraft and rockets. Lastly, a conclusion is presented in section 5.

2 Charging and Fragmentation Models

The current paper concerns itself in essence with the competition between the charging efficiency of two mechanisms; ionization and capacitive charging. The ionization mechanism is here represented by shock wave ionization – that compressibility in solids allow for high enough energy density for charges (electrons) to separate from their respective solids (nanoscale and microscale particles) when the impact speed is high enough. For this mechanism to take place, the particles need to impact a much larger bulk solid at speeds of several km s^{-1} , typically. The capacitive charging mechanism can be explained by its name in that it does not in principle differ from the macroscopic effects observed in a regular capacitor. It states that any particle with a non-zero conductivity can set up a potential difference with a bulk material if they have different work functions. Some confusion might lie in that this mechanism is often called ‘contact charging’, which might indicate that the separation between the particle and impact surface is zero. This is not true, as there would be no capacitive coupling in such a case. Rather the effective separation is around 1 nanometer due to quantum effects. Then, due to different affinities of the “capacitor” surfaces to keep or emit electrons, an effective potential is set up in which electrons trapped in potential wells on the surfaces of the capacitively connected bodies can jump from one to another. We must emphasize that the two mechanisms treated here are fundamentally different: no ionisation or dissociation takes place in contact charging due to broken bonds. The mechanisms can, as is a key result from our efforts, contribute to charge production simultaneously in a certain impact velocity range – *i.e. they compete*. It is also important to underline that we allow incoming particles to shatter – or fragment – in our treatment of contact charging. Fragmentation is simply breaking a projectile particle into a distribution of particles without any ionisation taking place. One implication of our application of fragmentation in connection with contact charging, is that the charge production is closely bounded to the distribution of fragments. Another consequence is that the experimental setups analysed here promote unipolar charge distributions, which is exactly what has been shown from rocket probings of mesospheric ice particles, and in laboratory measurements of ice-on-metal collisions, as elaborated on in section 3.3

In the sections below, we introduce the theoretical framework for our contact charging model based on fragmentation and capacitive charging, as well as the theory of shock wave ionization with special emphasis on the low velocity regime. We utilize our model on two slightly different types of projectile grains.

The motivation behind the model presented below, and approach for its utilization, can be summarized as follows: At speeds comparable to or lower than the ~~critical~~ critical limit for significant deformation or cratering in a grain-surface collision – see e.g. Jones et al. (1996) – there is little to no available material or energy for impact ionization. Moreover, the incoming projectile grains will fragment, as low energy collisions essentially can be viewed as a collision cascade as opposed to a sublimation-like destruction process. The consequence is that impact ionization models such as shock wave ionization ~~overestimate underestimate~~ overestimate underestimate the produced charge for dust-surface interactions at low velocities. Our solution is to invoke a model in which no direct charge (plasma) production takes place, but rather takes advantage of the fact that (semi-) conducting

grains can have a capacitive coupling to surfaces when there is a difference in effective work function between them. In such a scenario, the charge production is in simple terms due to electrons jumping between the surfaces of fragments and target in an effective potential.

2.1 Fragmentation Model

5 At speeds $\gtrsim 100 \text{ ms}^{-1}$, both metal particles (Froeschke et al., 2003) and water ice particles (Tomsic et al., 2003) of sizes $r_d \gtrsim 10 \text{ nm}$ fragment to a high degree. For water ice, molecular dynamics simulations and experimental evidence show a dependence of impact angle on the degree of fragmentation (Tomsic et al., 2001). Moreover, bulk properties can be used as a good approximation for those sizes. The case for smaller particles approaching the sub-nano scale is somewhat more complicated, as the yield stress, cohesive energy and work function can change as one approaches the atomic size limit (Qi 10 and Wang, 2002; Rennecke and Weber, 2014). We must note that the extrapolation of both the theoretical and experimental results cited here to a wide range of velocities and other grain parameters must be done with great caution, as we try to accomplish here. In section 3.1, we motivate our choice of fragment size limits in section 3.1 as in accordance with our current 15 understanding of collisions with our initial parameters. In the current work we employ two slightly different fragmentation models for pure metal particles and ice particles with impurities of meteoric smoke. Meteoric smoke particles (MSP) are coagulates and/or agglomerates of the remains of meteor ablations. However, it should be noted that the size distribution of fragments have the same proportionality with size in the two models, and thus the application of our model in the two cases have certain resemblance.

Figure 1 shows a sketch of the impact geometry. The process can be summarised as follows: an incoming grain (projectile) of size above $\sim 10 \text{ nanometres}$ hits a surface at a speed which allows it to deform plastically. Due to internal shears, a portion 20 of the particle can fragment into a distribution of smaller particles – which can also deform. In this work we have modelled the charge production both for a fragmenting and non-fragmenting particle. The speed at which the fragments interact with the surface is on the order of the initial speed of the main grain. The degree of fragmentation is controlled in our model by a parameter h introduced below. For the impact energies encountered in this paper, i.e. for corresponding impact velocities 25 of a few km s^{-1} , collisions can be assumed to be fully plastic (Rennecke and Weber, 2014), and we thus can utilize Hertzian deformation theory for the main projectile of radius R (with radius r_d in figure 1). The contact area over which a capacitive coupling is established (as described in section 2.2 below), is defined as (Wang and John, 1988) (John et al., 1980):

$$A = \alpha \pi r_p^2 \quad (1)$$

$$\alpha = \left(\frac{5}{4} \pi^2 \rho_p v_p^2 [k_t + k_p] \right)^{\frac{2}{5}} \quad (2)$$

where r_p is the projectile radius, v_p is the projectile impact velocity, ρ_p its density and k_t and k_p the target and projectile 30 elasticities defined by the Young's modulus E according to $k_i \approx 0.89/\pi E_i$. The deformation parameter α can be derived from Hertzian deformation theory, and we have used a result by Soo (1971). We assume that impact duration (\gtrsim picoseconds) is long enough to establish charge equilibrium. Considering the thermal speed of electrons, their mobility is by far high enough to obtain equilibrium for temperatures during impact. We must stress that our assumption of grains obeying Hertzian deformation 35

is only true for the lower part of the investigated velocity range, and that the main point of the current work is to study the speeds at which ionization efficiency is low and capacitive charging dominates. The upper boundary of 10 kms^{-1} is somewhat arbitrary in that it is a limit at which with certainty almost no contact charging takes place, and that shock wave ionization dominates. This is elaborated upon in section 2.3. Furthermore, we have defined a parameter h which gives the height of a cylinder of cross-section A in which all the material is fragmented. Molecular dynamics studies (Tomsic et al., 2003) and rocket results for low impact velocities (presented below) suggest that part of the grain material is decoupled from the rest. Thus only a small part of the projectile ~~contribute~~ contributes to the capacitively generated charge. Such an understanding implies in our model, for velocities well below the volume ionization regime, that only a fraction $3h\alpha/4r_p$ of the original particle is involved in charge production. The rest of the particle is shielded or decoupled from the target surface. As shown in our results below, $h \sim 0.1r_p$ offers a good fit for rocket data. Note that we have disregarded polarization effects, as the characteristic polarization potential switching times for particles of sizes used here, are likely much longer than the collision time (or contact time) (Havnes and Hartquist, 2016).




Figure 1. Contact geometry for the charging model of capacitive contact charging. An incoming grain of size above ~ 10 nanometres can fragment into a distribution of particles (fragments) that can furthermore deform and interact with the contact surface in a similar manner as the incoming main grain. The d subscript denotes *dust* here.

We employ the same parameterization of fragment size distribution in both the fragmentation-at-impact model (iron particles) and fragments-in-projectile model (ice particles containing meteoric smoke particles – MSPs), namely:

$$15 \quad N_f(r_d) = N_0 r_f^{-3} \quad (3)$$

where r_d and r_f is the fragment radius, are the respective incoming dust particle and fragment radius in metres and N_0 is a constant defined by the available volume for fragmentation. This similarity – The distribution gives absolute number densities (non-cumulative). The similarity between the two fragmentation models arises from the fact that grain-grain or grain-surface

collisions (Evans, 1994) and condensation of dust (Antonsen, 2019) yields the same r^{-3} -dependency. The grain-surface collision may be viewed as a collision cascade in free space, which typically yields similar size dependencies. It has also been confirmed that fragments of mesospheric ice containing impurities of meteoric smoke have a size distribution consistent with this model (Antonsen et al., 2017). Motivated by Antonsen and Havnes (2015), we also assume that for ice containing MSPs,
5 the ice will evaporate quickly and moreover have a much lower affinity for charge exchange. Thus it is only the embedded MSPs that contribute to the charge production. It should be pointed out that the two fragmentation models are fundamentally different in that for iron particles the fragments are produced at impact, while for ice particles the fragments (MSPs) keep their original size distribution resulting from condensation and/or coagulation. This latter model is consistent with findings on how ice particles containing impurity fragments are detected in sounding rocket impact probes at speeds $\sim 1000 \text{ ms}^{-1}$. Throughout
10 the discussion below, we assume that the fragments are spherical grains. We also take into account lattice sphere packing, and use a value of 70%; representable for HCP, FCC, CCP and BCC /hexagonal-close-packed), FCC (face-centred cubic), CCP (cubic-close-packed) and BCC (body-centered-cubic) lattice structures. The 70% value implies that 30% of the grain volume
15 is vacuum. The fragment size distributions are binned with a bin width of 0.01 nanometers and use a default size range of 0.5 to 4 nanometers in our numerical simulations. For some materials, such as iron, the lower size limit must be shifted in order to disregard quantum effects, as discussed in section 3. The size distribution of fragments in a complete fragmentation of a 30 nanometer ice particle with embedded MSPs is shown in figure 2. The produced charge Z_{tot} is the charge number bound to
20 fragments, calculated with our capacitive contact charging model. The model do not quantize charges, and only averages over the distribution of fragments, which may overpredict the contribution from the smaller fragments. The value of Z_{tot} is a factor $\sim 5 - 10$ larger than what is usually measured with sounding rockets (Havnes et al., 2014), and is thus a confirmation that only part of the projectile particle contributes to the measured charge. We elaborate on the charge yield scaling for fragmented particles in Appendix A.




Figure 2. Size distribution of ~~parameterized~~ parameterized MSP particles inside an ice particle of size 30 nm. Note that Z_{tot} is the upper bound on charge production for the case of complete charging of fragments. Annotated values show the result of employing our charging model on such a grain.

2.2 Contact Charging

The motivation for the current work is that at low impact speeds $\lesssim 1000 \text{ ms}^{-1}$ the charging of dust can be dominated by a capacitive charging mechanism where the projectile particle and (metal) target surface has an effective co-capacitance dependent on the difference in work function between target and projectile. As described in section 2.1 above, we take into account 5 that incoming projectile particles fragment at the speeds investigated, and we present a theory of contact charging for the individual fragments in the current section. The tensile strength of fragments increase with decreasing radius, and we assume that a capacitive charging model utilized on *fragments* is valid for much higher speeds compared to when utilized on single projectiles. In the following we give a short introduction to the contact charging theory presented by [Wang and John \(1988\)](#) [John et al. \(1980\)](#), and further developed by [Wang and John \(1988\)](#) to describe plastic projectiles, applied on our problem. It is 10 assumed that all impacts arrive perpendicular to the target, i.e. an impact angle $\theta = \pi/2$.

The fundamental mechanism behind contact charging as investigated here, is a capacitive coupling between a particle and a surface over an effective separation $Z_e \sim 10^{-9} \Delta \sim 10^{-9} \text{ m}$ (Dahneke, 1972). The produced charge can be described as a function of time by:

$$Q = CV_c \left(1 - e^{-\frac{\Delta t}{\tau} - \frac{\delta t}{\tau}} \right) \quad (4)$$

15 where V_c is the difference in work function of the contacting materials, $C = \epsilon A Z_e$ the capacitance and τ the charge relaxation time. For conductors, $\tau \sim 10^{-17} - 10^{-19} \text{ s}$, and $1 - \exp(-\Delta t/\tau) \approx \Delta t/\tau$, and it can be shown that eq. (4) reduces to:

$$Q = \chi \pi r_p^2 v_p \frac{\epsilon V_c}{Z_e} \left(\frac{4 \rho_p}{3Y} \right)^{\frac{1}{2}} \quad (5)$$

where we have introduced the yield stress Y of the material that yields first and the permittivity $\epsilon \approx \epsilon_0$. The parameter χ is a 20 constant between 0 and 1 that we have introduced in our parameterization of fragment charging. It can be understood as the proportion of fragments which are charged until equilibrium, or alternatively as the charging probability of a single fragment. The χ -parameter is thus in essence a fitting parameter, and we note that we do not include the possibility of size dependency of it in our model. Note that it was used (set to non-unity) when producing the results in figures 4 and 5. In table 1 we 25 summarize the parameters used to produced the presented results. It must be noted, that even if bulk values can be extrapolated to very small particle sizes, that some of the parameters utilized in the numerical computations here vary with temperature which we do not take into account here: The static, i.e. low frequency equilibrium, relative permittivity of ice increases with [decreasing](#) [decreasing](#) temperature (Auty and Cole, 1952; MacDowell and Vega, 2010). The [Yield](#)-[yield](#) stress and Young's moduli for olivine and ice may also change significantly with temperature (Evans and Goetze, 1979; Nunez-Valdez et al., 2010; Nimmo, 2004). While metals are somewhat more resilient to changes in parameters related to internal stress, their work 30 function increases with decreasing size due to a change in polarizability as particles become small (Wood, 1981).

Table 1. Material properties used in the calculation of contact charging yields for silver (Ag), iron (Fe), water ice, **Stainless** **stainless** steel (**SLSSS**) and a meteoric smoke analogue (MSP).

	Ag	Fe	Ice	SLSSS [†]	MSP ^{††}
$\rho[\text{kgm}^{-3}]$	10500	7874	980	7800	3000
$E[\times 10^9 \text{Nm}^{-2}]$	104	150	9	170	200
$Y[\times 10^6 \text{Nm}^{-2}]$	330	50	10	50	50
$\phi[\text{eV}]$	4.7	4.5	–	4.4	7.3–8.5

[†] Stainless steel 316. ^{††} Meteoric smoke particles; here Olivine at 300 K (Rapp et al., 2012).

2.3 Shock Wave Impact Ionization

It is recognized that impact ionization is a combination of mechanisms, each dominating for certain parts in a wide velocity range. In this paper, we take it as fact that the impact ionization will tend towards a volume ionization mechanism – as a consequence of a Thomas-Fermi model for electronic structure – as impact speeds exceed $\sim 50 \text{ kms}^{-1}$ (see e.g. Auer (2012)).

5 For velocities below such high speeds, the accepted and most widely used model for impact ionization is that of Drapatz and Michel (1974), which describes ionization as a result of shock waves propagating through colliding entities.

The model of shock wave ionization does not, however, describe ionization for the entire velocity range below $\sim 50 \text{ kms}^{-1}$. This was also recognized in the earliest formulations of the theory and discussions on its validity. There are in fact at least two velocity regimes – low ($v_p \lesssim 5 - 10 \text{ kms}^{-1}$) and high ($v_p \gtrsim 10 \text{ kms}^{-1}$) – that display different semi-empirical charge yields.

10 The charge production mechanism is different in the two velocity regimes, and one focus of the present work is to put the theory of shock wave ionization in the low velocity regime under scrutiny. In section 4.1 we discuss in detail the thermodynamics of the low velocity regime shock wave ionization as it was formulated by Drapatz and Michel (1974).

15 For the high velocity regime, the shock wave ionization model assumes that the ionization state freezes at some point during expansion of the **gas** **ball** **impact** **cloud** arising from impact (Raizer, 1960; Kuznetsov and Raizer, 1965). The degree of ionization can then be calculated from a Saha-equation (Dresser, 1968):

$$\frac{n_+ n_-}{n_0} = \frac{2\psi_+}{\psi_0} \left(\frac{2\pi m k_B T}{h^2} \right)^{\frac{3}{2}} \exp \left[-\frac{eV_I^1}{k_B T} \right] \quad (6)$$

where n_+ , n_- and n_0 are the respective number densities of ions, electrons and neutrals; ψ are atomic weights for the ionic and atomic states; T denotes the temperature; k_B denotes Boltzmann's constant; $|e| = 1.6 \times 10^{-19} \text{ Coulombs}$; h denotes Planck's constant, and V_I^1 denotes the first ionization potential of projectile atoms.

20 For the low velocity regime, Drapatz and Michel (1974) pointed out that impurity ionization predominantly from alkali metals in the projectiles was responsible for the charge yield. One must then **utilize** **utilise** that electrons bound in metals follow a distribution (Copley and Phipps (1935); for potassium on tungsten):

$$n_- = 2 \left(\frac{2\pi m k_B T}{h^2} \right)^{\frac{3}{2}} \exp \left[-\frac{e\phi}{k_B T} \right] \quad (7)$$

where ϕ is the work function of the projectile. This ~~ultimate~~ultimately leads, after insertion into eq. (6), to what can be recognized as the Saha-Langmuir equation:

$$\frac{n_+}{n_-} = \frac{\psi_+}{\psi_0} \exp \left[-e \frac{\phi - V_I^1}{k_B T} \right] \quad (8)$$

where for the materials discussed here $\psi_+/\psi_0 \approx 2$.

5 In the following, we consider the situation that both target and projectile are conductive. This means, in simplified terms, that electrons can easily move between potential wells on the surface of the projectile and target and have time to equilibrate the charge within the collision time τ . Then, in all generality, the charge yield can be described by power laws in both velocity and particle size. The charge production is thus often described by the formula:

$$Q[\text{C}] = \gamma m_p [\text{kg}]^\alpha v_p [\text{km s}^{-1}]^\beta \quad (9)$$

10 where the constants γ, α and β are all strongly dependent on material properties and velocity regime (see e.g. Mocker et al. (2013); Collette et al. (2014); Kissel and Krueger (1987)). For the application on micrometeoroid impacts on spacecraft, the ~~velocity exponent~~ most widely cited value for velocity dependence is $\beta \approx 3.5$ adopted from McBride and McDonnell (1999), however values of $2.5 - 6.2$ have been reported for common spacecraft materials (Mann et al., 2019). The exponent α for mass dependence is usually found to be ~ 0.7 for the low velocity regime and close to unity for the high velocity regime. It has
15 already been pointed out, by Kissel and Krueger (1987), that for low velocity impacts ($v_p < 5 \text{ km s}^{-1}$) α should be close to $2/3$. This is to say that the charge yield is proportional to the incoming projectile cross section r_p^2 . The same authors also pointed out that, both in the low and high velocity regime, ionization at the target could also be described by the same power law stated in eq. (9). However, the exponent for mass dependency would be one dimension in particle size lower; i.e. $Q_{\text{target}} \propto r_p^2$ for high projectile velocities and $Q_{\text{target}} \propto r_p$ for low velocities. In this work, we do not discuss the additional effect of direct target
20 ionization. In section 4.1 we elaborate in that low velocity impact charging must probably be described by a different physical mechanism than shock wave ionization as ~~described~~described here.

3 Results

In sections 3.2 and 3.3 below, we present results from calculations of contact charging for projectiles of metal and projectiles of ice on metal surfaces, respectively. The simulations employ the fragmentation and charging models described above. For
25 calculations of ice-on-metal charge yields, we assume ~~for~~ that the ice particles are contaminated with meteoric smoke particles. This model is descriptive for icy dust particles (or aerosols) in the Earth's mesosphere, and we compare our results with in-situ measurements of mesospheric ice.

3.1 A note on projectile size and fragment size sensitivity

The choice of default projectile grain size (30 nm) in the presented model results below, may be motivated by that it is a typical
30 ~~size of~~for mesospheric icy dust grains usually encountered by sounding rockets. It is also among the smaller projectile sizes

(see e.g. figure 2 in Mann et al. (2019)) which can be readily generated in typical dust accelerator experiments. Moreover, the charge produced at a specific projectile size, can be directly scaled to larger sizes according to the scaling relation given in eq. (A5). Normalizing the calculated yield to the projectile mass furthermore allows for direct comparison to semi-empirical laboratory results.

We must also address the choice of limits in fragment size distributions, and sensitivity to changes in the lower cut-off limit. In their treatment of collisional charging of interstellar grains, Draine and Sutin (1987) argued on the basis of results by Omont (1986), that bulk properties would sufficiently describe grains of PAHs [polycyclic aromatic hydrocarbons \(PAH\)](#) down to sizes of only 3 Å – or $\gtrsim 30$ molecules. That is to say that, at least for carbonaceous or PAH dust, one can model particles as conducting spheres and disregard quantum effects when calculating the equilibrium charge due to polarization (image) and capture of charged species. We use this as one reason to model fragments of impacted projectiles as conducting spheres down to sub-nanoscale. In our framework of modelling metal on metal interaction, we must note that the atomic interspacing is ~ 2 Å in a BCC lattice structure. Thus, to obtain a grain that satisfies the constraints used in the references above, we must increase our cut-off to ~ 6 Å for iron fragments. This is also in agreement with the findings of Jones et al. (1996) for minimum fragment size in low velocity impacts. In figure 3 we compare the response of our contact charging model for different low-size cut-offs in the fragment size distributions. The differences are small – an increase from 2 Å to 8 Å in cut-off only decreases yield by around 30%. Thus we increase the low-size cut-off of iron fragments to 7 Å in this work while still keeping the conducting sphere assumption elaborated on above.




Figure 3. Sensitivity of contact charge generation in an Fe-on-Ag collision to different values of lowest allowed fragment sizes. Cut-off values are labelled in the legend.

3.2 Metal–metal collisions

In the following simulations, we have used iron as projectile material and silver as target material. This is due to that experiments with this combination have been done in the past both at the LASP dust accelerator (Collette et al., 2014) and at the Heidelberg dust accelerator facility (Mocker et al., 2013). The Fe–Ag combination may be applicable to dust impacts on space-

5 craft as iron is common in e.g. micrometeorites. It was also used as an example by Drapatz and Michel (1974) in the original formulation of the shock wave ionization theory. Thus, it is possible to compare our results with several others. It should be noted, however, that different experiments may have different complex geometry and working principles. The data produced by the different experiments may therefore have intrinsically systematic differences and direct intercomparison must be done with care.

10 Figure 4 shows a comparison of our calculations of iron projectiles on a silver target to the semi-empirical results obtained by Mocker et al. (2013). ~~Their data (It shows that our model utilized with bulk material properties, and a very high degree of fragmentation (or charging probability) over-predicts the observed charge production, however, with a very similar power law exponent. Although the Mocker et al.-results have not been obtained for the projectile velocities as low as the lowest velocities shown in the figure, we have utilized the semi-empirical law from their work, as their results are virtually indistinguishable~~

15 from the results of Collette et al., which was obtained for velocities down to 2 km s^{-1} . We must however note that since we have extrapolated these results to lower speeds, some caution should be taken when comparing the two in this part of the velocity range. A further deeper discussion of the charge production for the lower velocities, is presented in section 4.2. The Mocker et al.-data (blue curves) show a discontinuity – or sudden increase in yield – at $\sim 11 \text{ km s}^{-1}$. Since the thermodynamics and chemistry of the fragmentation process might not be valid at speeds much higher than this limit, we focus on comparing

20 our curves with the experimental data at speeds below the discontinuity. The lower limit of contact charging, showed as a solid black line, is the case where we consider that the entire original projectile participates in the capacitive coupling. This assumption implies that the entire particle must be bound together, while it still allows for plastic deformation. Such a situation is not probable, but provides a lower boundary on the charge production. The fragmentation model results ~~is-are~~ sketched as a dashed black line. In general, it has a slope very close to the semi-empirical model, but its values are ~~to~~ two orders of

25 magnitude higher. We note that figure 4 shows a default run with default parameters suitable for bulk material, and refinement of these will lead to different results, as shown below. Moreover, as shown in Appendix A, there is a strong dependency on the parameterization of the fragmentation size distribution. Refining the parameterization yields a much better similarity to the experimental results. Although it is not the purpose or motivation of this work to explain the entire charging mechanism at low impact speeds with fragmentational contact charging, we nevertheless have calculated a best fit of our model to experimental

30 data with reasonable parameters. In figure 5 we show the result of a simulation with a set of parameters that produces a 'best fit'. The yield stress was increased by a factor three compared to figure 4. The yield stress is in any case a parameter with a significant uncertainty for nano- and microscale particles. Moreover, the fraction of fragments that become charged was reduced to $\chi = 1\%$. This parameter is difficult to define, and has a large intrinsic uncertainty. A final adjustment was made to the size distribution; where the smallest possible fragment size was changed from the default value of 0.5 nm to 0.7 nm.




Figure 4. Simulation of contact charging of iron projectiles ($r_{p,0} = 30$ nm) on a silver target with (black dashed line) and without (black solid line) a fragmentation model. The power laws plotted in blue were acquired experimentally by Mocker et al. (2013). In this calculation, the yield pressure of iron was set to $Y = 50$ GPa, and the fragment size span was set to $[0.2, 3]$ nanometers. The label '50%' indicates that in this calculation, $\chi = 0.5$ cf. equation (5).

Since many of the parameters used in our charge model are valid for bulk projectiles, the validity of extrapolating the model to sizes $\lesssim 10$ nm can be a topic for discussion. In a more rigorous treatment, one may have to take into account curvature and polarization effects for the smallest fragments. Nevertheless, our model shows that at low speeds, fragments can indeed produce charge in a capacitive coupling very efficiently as opposed to ionization through a Saha-process. We also leave the 5 untreated issue of how much the pre-charge, which can be very large for the large projectiles that dominate the low velocity range, contributes to the yield.




Figure 5. Simulation of contact charging of iron projectiles on a silver target. This simulation was a "best fit" run, in which the fraction of charged fragments, yield pressure, and fragment size span was allowed to change. This shows the case for $\chi = 0.01$ cf. equation (5), $Y = 150$ GPa and fragment size span [0.7, 3] nanometers.

3.3 Ice–metal collisions

It is technically challenging to set up laboratory experiments for studying low impact velocities ($\lesssim 1 \text{ km s}^{-1}$) and small projectiles ($\lesssim 100 \text{ nm}$) simultaneously. Dust accelerators typically use samples of projectile particles which span several orders of magnitude, while the energy is fixed and determined by the strength of a static accelerating potential (see e.g. Thomas 5 et al. (2017) for LASP setup). Such a configuration implies that only very large projectiles will have low impact speeds. The pre-charge collected by the particles in such experiments are usually large – close to field emission limits.

It is however possible to use sounding rockets to obtain a point measurement in the low size and speed range: Typical sounding rockets utilized in upper atmosphere research operate at low speeds $\sim 1 \text{ km s}^{-1}$, and naturally occurring dust particles in the mesosphere ($\sim 50 – 100 \text{ km}$ above sea level) typically have sizes $\sim 1 – 100 \text{ nm}$. Thus, in-situ measurement can 10 be compared to laboratory measurements for certain experimental setups. For the results below, we utilize the rocketborne Faraday impact probe MUDD (MULTiple Dust Detector). Inside it, incoming projectile dust particles hit a slanted stainless steel plane on which they deposit pre-charge and acquired mainly by ambient electron and ion collection and produce contact charge. Photoelectric charging can become important under certain conditions. A detailed technical description, projectile dust dynamics and utilization of the instrument can be found in Havnes et al. (2014); Antonsen and Havnes (2015); Antonsen et al. 15 (2017).

As previously stated, we utilize that dust grains in the mesosphere are contaminated with meteoric smoke – recondensed and agglomerated remnants of meteoric ablation. In figure 6 we show the result of two limiting cases of contact charging of 'dirty ice'. The solid black line describes a situation where no impurities contribute to the produced contact charge (the pre-charge is assumed to be zero). This might be plausible for very low speeds, where ice particles of sizes $\sim 10 \text{ nm}$ experience 20 less fragmentation (see e.g. Tomsic et al. (2003)). For projectile speeds on the order of a few hundred meters per second, the fragmentation model (dashed line) should provide a more physically sound charge yield. For a typical rocket speed of $800\text{-}1000 \text{ ms}^{-1}$ the fragmentation model and single projectile have roughly the same yield. We find that the predicted charge number for this velocity range is consistent with what has been measured with rocketborne Faraday cups (Havnes and Næsheim, 2007; Havnes et al., 2014). The gray shaded area shows a-values of the predicted charge yield where the ice particles have a capacitive 25 coupling, but are allowed to have a non-unity dielectric constant (cf. Wang and John (1988)). This effectively means that the ice particles are insulating, which may be a better description. The true yield of a pure ice projectile should therefore probably lie below the solid black line in figure 6.

In the following we attempt to simulate the current recorded by MUDD during a flight in the MAXIDUSTY campaign (Andøya Space Center, June 30th 2016). We assume the finding of Antonsen (2019); that small ice fragments thermalize 30 and evaporate very quickly and MSP fragments dominate the produced signal. The size distribution of MSPs inside ice was previously found to be $N \propto r_f^{-[2.6, 4.4]}$ (Antonsen et al., 2017), thus an exponent of -3 should fit well. Consequently, we can employ the same fragmentation model as for metal-metal collisions. We use a volume content of MSPs of 1%, which is in the middle of the range of what has been found from rocket measurements and satellite measurements (Hervig et al., 2012). Other




Figure 6. Contact charge yield of 30 nm ice particles with MSP impurities impacting on stainless steel. The shaded area shows possible yields for the case of a mixture of insulating and conducting particles.

parameters used in the calculations are listed in table 1. We also note, that the impact plane in MUDD is slanted, however, we assume that the contact time is on the same order as for head-on collisions and long enough to reach equilibrium.

The rocket traversed a dust layer situated at $\sim 81 - 87$ km at a velocity $\approx 810 \text{ ms}^{-1}$. ~~Accurate~~ ~~We use the accurate and high altitude resolution~~ number densities and sizes of ice particles ~~were as~~ found by a combination of ALOMAR RMR lidar data, in-situ photometer and DUSTY Faraday cup data as described in Havnes et al. (2019). ~~We have assumed, as in the cited paper, that the mesospheric ice particles are monodisperse and we have utilized an average pre-charge on the ice particles. The electron and ion density acquisition densities for this flight was done with Faraday-rotation antennae and capacitive probes, and the acquisition is explained in the same reference. We note that a more rigorous modelling of the MUDD-currents could allow for size- and charge distributions rather than an average of these.~~ Figure 7 shows the comparison between measured MUDD currents and simulated MUDD currents using the described fragmentation model. The two curves display a very high similarity down to the smallest scales, and only differ significantly at the upper ~ 1 km of the dust layer. Combined with the fact that the fragments – i.e. the embedded MSPs – do not carry significant pre-charge, our results presents a convincing case for contact charging being the dominant charging process for the speeds and particle sizes encountered when probing the dust in the Earth’s mesosphere. ~~For reasons elaborated on in section 4.2 below, the expected contribution from shock wave ionization as calculated with Drapatz’ and Michel’s low velocity solution is negligible in comparison with contact charging and even direct pre-charge current in MUDD, at the rocket’s velocity, Thus we have not included a comparison simulation for impact shock wave ionization here.~~




Figure 7. Measurements from the impact Faraday cup MUDD flown on the MXD-1 sounding rocket payload (red) and a best fit from simulation of contact charging (grey) using the fragmentation model described in 2.1. $V_c = 0.5$ eV was found to be the best fit for fragments of density $\rho = 3000$ kgm $^{-3}$ and yield pressure 50 MPa. The minimum fragment size threshold was set to 0.3 nm.

4 Discussion

As presented in section 2.3, it was pointed out by Kissel and Krueger (1987) that the Saha-Langmuir solution (SLS) from Drapatz and Michel (1974) underestimates impact charge generation for speeds $\lesssim 5 \text{ kms}^{-1}$. The low velocity solution of their theory assumes that charge is generated mainly by impurity diffusion through the molten projectile material. Although Mocker et al. (2013), whose results we have used for comparison, conclude with an agreement with the SLS, it must be noted that they find that the appearance of Fe in impact time-of-flight mass spectra occurs at much lower speeds (3.6 kms^{-1}) than the SLS predicts. Hydrocode simulations aiming at predicting the threshold of impact plasma generation in iron-on-metal collisions have found a threshold of 8 kms^{-1} (Ratcliff et al., 1997). [Results from the Cassini Cosmic Dust Analyzer have confirmed occurrence of metal ions in time-of-flight mass spectra at speeds \$\gtrsim 10 \text{ kms}^{-1}\$ \(hil, 2007\)](#). The takeaway from these notions, is that both direct impact plasma generation and the SLS probably is insufficient in explaining charge generation in low velocity impacts.

To investigate the applicability of an SLS at speeds on the order of 1 kms^{-1} , we look closer at the thermodynamics of the process in section 4.1. In section 4.2 we discuss possible areas of application of our contact charging model, with emphasis on spacecraft and sounding rocket observations.

15 4.1 Thermodynamics of Low Velocity Limit of Shock Wave Ionization

A first order estimate of the mean diffusion distance of an impurity ion inside a cooling – i.e. solidifying – metal grain, can be found by recognizing that the diffused area must be $D(T)\tau_s$, where $D(T)$ is the temperature dependent diffusivity over a solidification time τ_s . We have that the one-dimensional mean diffusion distance is

$$\delta r = |(D(T)\tau_s)^{\frac{1}{2}}| \quad (10)$$

20 This can be interpreted as the thickness of a shell from which impurities can reach the surface of a cooling grain. For the purpose of comparing diffusion of alkali impurities through iron particles with the results of Drapatz and Michel (1974), we utilize the mean diffusivity $\overline{D(T)} = 5 \cdot 10^{-9} \exp(-5000/\overline{T}) \text{ m}^2 \text{s}^{-1}$.

The available amount of substance from which ions can be released from a grain of radius r_f has a volume of – assuming We assume the particle has bulk properties, which is suitable for particles clusters of size on the order of $\sim 10 \text{ nm}$ ten nanometres. 25 The available volume from which ions can be released is then (denoting the grain radius r_f):

$$\delta V(\tau_s) = \frac{4\pi}{3} [r_f^3 - (r_f - \delta r)^3]. \quad (11)$$

Then the upper bound on the number of impurity ions released from a single grain becomes:

$$N_{im} = \underline{KK} \frac{4\pi \rho_p \xi}{3M_p} [r_f^3 - (r_f - \delta r)^3] \quad (12)$$

where ξ is the impurity content by volume and ρ_p and M_p are the mass density and molecular mass of the grain ('projectile') material respectively. We use here that $\xi \sim 1\%$, which is representable for alkali metal content in Earth's crust and in raw

smelted iron and steel. $\frac{K}{K}$ is the atomic packing factor, which is set to 0.7 in our calculations. The resulting single charge impurity ionization predicted by the Saha-Langmuir equation then becomes

$$Z_{im} = N_{im} \frac{n_{im}}{n_{im} + n_{Fe}} \approx 2N_{im} e^{-\frac{e\phi - V_I^1}{k_B T}} \quad (13)$$

where e is the elementary charge, ϕ is the work function of the impurity material and V_I^1 is the first ionization potential. The 5 difference between these for potassium (K), which is used in this work as a dominant impurity, is $\phi - V_I^1 \approx 1.8$ eV. The factor 2 arises from the statistical weights in equation (8).

It is clear that we also require a parameterization of the temperature of inside the expanding shock. For this purpose we utilize that the relationship between the shock front velocity u and the projectile velocity v_p is governed to first order by the ratio of the difference in mass density between the projectile and target through energy conservation. We have for specific 10 energy

$$\varepsilon = \frac{u^2}{2} = \frac{v_p^2}{2} \left/ \left(\frac{\rho_p}{\rho_t} \right)^{\frac{1}{2}} + 1 \right. \quad (14)$$

which for our example case of iron projectiles on a silver surface yields $u \approx 0.62v_p$. This is to say that $\sim 40\%$ of the initial energy goes into expansion of the shock. It can be shown than the temperature behind the shock for monoatomic gases ($\gamma = 5/3$) is (Zel'Dovich and Raizer, 1967):

$$15 \quad \frac{T}{T_0} = \frac{5}{16} \text{Ma}^2 \quad (15)$$

where $\text{Ma} = u/v_{th,N}$ is the shock front Mach-number and T_0 is the pre-collision temperature.

In the following calculations of impurity ionization production, we have used that the solidification temperature of nanoscale iron particles is 1000 K (Fedorov et al., 2017). Furthermore, we employ a cooling rate of 10^{12} Ks $^{-1}$, which has been found from molecular dynamics simulations to be representable for nanoscale metal particles (Shibuta and Suzuki, 2011). This gives 20 typical solidification times of $\tau_s \sim 10^{-9}$ seconds, which is two orders of magnitude smaller than the cooling time used by Drapatz and Michel (1974). In the current model, we restrain the diffusivity coefficient with a hard stop at the solidification temperature, and do not parameterize solidification/crystallization effects. Moreover, we do not discuss here the evaporation of impurities from the surface of the main particle; we simply assume all impurities are removed, and thus present an upper 25 bound on impurity charge production. By the set of equations above, we find that the limit for impurity production for a 3 nm iron grain is a Mach-number of 3.3, corresponding to a velocity $v_p \sim 1$ kms $^{-1}$.

In figure 8 we have not parameterized the temperature decrease in the expanding **gas**-volume behind the shock. However, depending on whether or not thermodynamic equilibrium can be reached or not, this effect might be a significant inhibitor of thermal ionization described by the S-L equation. For an adiabatic expansion, we have that

$$\frac{T(\tau_s)}{T_0} = \left(\frac{V_0}{V(\tau_s)} \right)^{\gamma-1}. \quad (16)$$

30 Thus, in the case of a 30 nm projectile particle and a solidification time of $\tau_s = 10^{-9}$ seconds, it is found that the limiting expansion velocity to accomodate diffusion should be on the order of $u \sim 100$ ms $^{-1}$, which is clearly never the case. In this




Figure 8. Results from calculation of impurity (1% Potassium) charging using the Saha-Langmuir equation (blue, dashed) and fragmentation model described in this work. The solid red line ~~shows~~ shows the number of released K-atoms as a function of velocity, and therefore constitutes a theoretical upper bound on the charge number (for singly charged ions).

regard, we note that the assumption of thermodynamic equilibrium may not be suitable for the set of parameters encountered in the current work. Moreover, we must note that the emissivity of nanoscale dust grains is strongly dependent on size and material properties, so the cooling time may also need refinement (Rizk et al., 1991).

We summarize our result of low velocity impact charging in figure 9. The SLS is vanishing below $\sim 2 \text{ kms}^{-1}$. The contact charge solution overestimates the experimentally acquired yield in the entire range. The two curves have almost the velocity and mass dependence, and possible downshifts of the contact charge solution were discussed in section 3.2. Notably, we have used $\chi = 1$ here, which provides an upper boundary on the charge yield. We must note that in the mechanism proposed in the current work, we have not taken into account charge production at the surface. Such an effect may be expected to scale with the particle radius rather than cross-section at low speeds, as mentioned in section 2.3. To first order, one may therefore disregard such additional charging, as ~~is-it~~ introduces another layer of complexity into modelling efforts. There is also a possibility of impurities on the surface – which are arguably always present in metals – that can produce additional charging. Extending our theory to include the contributions from impurities may require a treatment of surface chemistry and evaporation ~~gas~~ microphysics which is beyond the goal of our study. Nevertheless, our results must always be read with the ulterior notion that volatile impurities such as alkali metals can introduce additional charge.

15 4.2 Relevance for Dust Detection on Spacecraft and Rockets

In section 3.3 we demonstrated the applicability of our model to rocket measurements of dust (or aerosols) in the upper mesosphere of Earth. Other related types of dust, namely those originating in the ablation of meteors in the altitude range ~ 70




Figure 9. Comparison of specific yields from our contact charging model (dashed) to the Saha-Langmuir solution from Drapatz and Michel (1974) (solid blue) and the semi-empirical law obtained by Mocker et al. (2013) (solid red). All lines represent Fe-on-Ag impacts, and in the Saha-Langmuir solution we have used potassium as the impurity. The label '100%' in the legend corresponds to $\chi = 1$ and indicates that all produced fragments collide with the surface.

to 140 km, are candidates for comparison with a contact charging model. Free dust grains of meteoric origin have recently been observed by sounding rockets (Havnes et al., 2018), and other novel experiments have been aimed at investigating such particles (see e.g. Strelnikov et al. (2018)).

The number of *catalogued* man made space debris objects in the near-Earth space is already on the order of 10^4 , with the number of objects increasing inversely with size (Klinkrad, 2010). Estimates of the number of objects smaller than $100 \mu\text{m}$ have large uncertainties, but have been cited as in the order of tens of billions (Schildknecht, 2007). The probability of any satellite encountering a space debris object during its lifetime is therefore nonvanishing. Besides this, typical orbital speeds of Low Earth Orbiting satellites are $\lesssim 8 \text{ km s}^{-1}$, and decreasing with increasing orbital altitude. Thus, even without considering whether the grains of debris are pro- or retrograde, debris-satellite interactions may readily occur at speeds relevant for contact charging. Additional charging or upsets due to contact charging is arguably undesirable for e.g. satellites measuring plasma parameters.

Based on recent observations by the Parker Solar Probe (PSP), Szalay et al. (2020) concluded that β -meteoroids (β describing the ratio of the radiation pressure force to gravity) dominate the recorded dust flux. They found that such grains typically have larger impact velocities than circularly bounded dust, but for $\beta \lesssim 0.5$, there may be a nonvanishing flux of β -meteoroids with impact speeds in the upper limit of the velocity range investigated in the present work for contact charging. From their results of modelling the dynamics of dust in bounded circular orbits, based on a model of Pokorný and Kuchner (2019), it is clear that there may well be a smaller number of impacts on PSP that can be traced to such dust grains.

Page et al. (2020) reported that although β -meteoroids can produce dust impact fluxes as measured on PSP near perihelion, some of the directionality in the dust flux data can be consistent with prograde circular orbit dust. At perihelion, the impact

velocity of these is still to large to apply a contact charge model ($\sim 20 \text{ kms}^{-1}$), however, it might be a possibility that one can use such a model on prograde dust further away from the Sun with lower impact velocities. In regards to further explorations of the utilization of our model on spacecraft data, dependence of impact inclination in contact charge production should also be studied.

5 Another possible candidate for employment of our model on spacecraft data, is secondary ejecta. Secondary ejecta, which is material from craters generated by dust impacts on the spacecraft body, have energies much lower than the impacting grains. Such secondary grains have been observed as stray light in optical images from e.g. STEREO (St. Cyr et al., 2009). Szalay et al. (2020) also noted that such secondary particles were observed with the WISPR experiment on PSP (see e.g. Vourlidas et al. (2016)), and that the ejecta correlated well with antenna measurements of dust impacts.

10 One impediment to utilizing our model on dust in space, is that it may be difficult to determine its composition and structure. In consequence, the work function of the projectile material may be unknown and moreover size dependent (Wood, 1981). In some cases, where the projectile grains have very low conductivity, it may be required to either: 1. Assign an effective work function (Matsusaka et al., 2010) or 2. Extend our theory to insulating particles. The latter can in brief be described as letting the the ratio $\Delta t/\tau \rightarrow 0$ in eq. (4). The result is a slightly lower velocity dependence in the charge production; $Q_c \propto v_p^{3/5}$. The 15 charge production will also be significantly weaker than for conducting or semi-conducting grains (John et al., 1980; Wang and John, 1988).

The ESA Solar Orbiter (ESO) was launched in February 2020. Its orbit is different from PSP in that its perihelia are larger than $\sim 0.28 \text{ AU}$ throughout its lifetime – versus $\sim 0.046 \text{ AU}$ for PSP. In addition, [Solar Orbiter ESO](#) has a planned 25 degree inclination in its nominal mission. The orbital parameters will ensure that ESO will encounter prospective bounded dust grains 20 and β -meteoroids with generally lower velocities than for PSP. It is not in the scope of this paper to analyze the expected dust flux of ESO, but it can be expected that a larger number of dust impacts can involve a contact charging mechanism compared to PSP.

5 Conclusions

In this work we have investigated the production of charge in impacts of projectiles of iron and agglomerates of ice and 25 meteoric smoke on a metal surface at speeds $\lesssim 10 \text{ kms}^{-1}$. We introduce a novel model of contact charging due to a capacitive coupling between metal surfaces and fragments of projectile grains. Here we show that our model is consistent with laboratory measurements of Fe-on-Ag collisions as well as rocket measurements of icy dust particles on stainless steel. Our method can be utilized with a large range of projectile dust types, where the intrinsic properties of the grains are known. We also find that our theory may be used to explain certain observations of dust by the recently launched spacecraft NASA Parker Solar Probe 30 and ESA Solar Orbiter. We moreover find that the currently accepted theory for impact charging at the speeds of interest here, namely the shock wave ionization theory of Drapatz and Michel (1974), is insufficient in explaining laboratory observations of charge generation in metal-on-metal impacts [alone](#). Consequently, we suggest that at low speeds, there must be a significant

contribution to the produced charge by contact charging — i.e. the two discussed mechanisms both contribute significantly to impact charge production at speeds from ~ 3 up to the limit where direct ionization becomes important.

Appendix A: Scaling relation for charge yield of fragmented particles

In this appendix we give a scaling relation for charge production by capacitive charging when employing a fragmentation

5 model.

The available material from which fragments can form is given by the Hertzian deformation presented in section 2.1 above, and is:

$$V_c = \alpha h \pi r_p^2 \propto r_p^3 \sim Q_p. \quad (\text{A1})$$

The largest possible spherical fragment (of volume V_s) that can be formed from this material, using $V_s(r_f) = V_c(r_p)$, has
10 radius:

$$r_{\max} = \left(\frac{3h\alpha r_p^2}{4} \right)^{\frac{1}{3}} \propto r_p. \quad (\text{A2})$$

As contact charging scales with the cross-section of fragments, we calculate the total surface area of all (discretely distributed) fragments:

$$S'_{\text{tot}} = \sum_{\forall i} \pi \left(N_0 r_{f,i}^{-3} \right) r_{f,i}^2 \quad (\text{A3})$$

15 where we have used that the fragments are distributed in size according to $N_f \propto r_f^{-3}$. If we assume that the size distribution is continuous, we can moreover find that

$$S_{\text{tot}} = \pi N_0 \int_{r_{\min}}^{r_{\max}} dr_f \cdot r_f^{-1} = \pi N_0 \ln \left(\frac{r_{\max}}{r_{\min}} \right) \quad (\text{A4})$$

where r_{\min} is the smallest possible fragment radius.

Now we recall the scaling $Q_p \sim r_p^3$ from eq. (A1), which constraints the amount of material available for fragmentation.
20 Inserting the result from eq. (A2) into eq. (A4) we finally obtain that the charge production of a fragmented projectile particle scales with size (and velocity according to its dependence in α):

$$Q_p \propto r_p^3 \ln \left(\frac{(3h\alpha r_p^2)^{1/3}}{4^{1/3} r_{\min}} \right) \propto r_p^3 \ln \left(\frac{r_p v_p^{4/15}}{r_{\min}} \right). \quad (\text{A5})$$

This result is also intuitively reasonable; that since there are many more small particles than large ones, the surface area of the small particles contributes more to the total area and thus charge production. Moreover, we note that the sensitivity to
25 the parameter r_{\min} becomes even more important for size distributions with steeper inverse power laws than the one chosen here. This solution will never become unphysical (singularity as $r_{\min} \rightarrow 0$), as r_{\min} has a natural lower bound. In this paper

its dependence on velocity, which must be assumed that is has. However, we assume that that a feasible value would be on the order of ~ 1 Ångstrøm ($= 0.1$ nanometer), which is the order of the length of a single atom or molecule. As seen in section 3, our charge production model is relatively sensitive to this parameter.

Acknowledgements. This work was supported by the Research Council of Norway through grant number 262941. The rocket campaign 5 and the construction of the rocket instrumentation was supported by grants from the Norwegian Space Centre (VIT.04.14.7, VIT.02.14.1, VIT.03.15.7, VIT.03.16.7) and the Research Council of Norway, grant 240065. J.V. and L.N. were supported by the Czech Science Foundation under Project 20-13616Y. The publication charges for this article have been funded by a grant from the publication fund of UiT The Arctic University of Norway.

Data availability. The data to reproduce the rocket measurements in figure 7 can be obtained from the UiT Open reseach Repository at 10 <https://doi.org/10.18710/N8GF1U>. The relevant data set is tagged 'M1BP_m2'.

Competing interests. The authors declare that they have no conflicts of interest.

References

Interplanetary dust detected by the Cassini CDA Chemical Analyser, *Icarus*, 190, 643 – 654, <https://doi.org/10.1016/j.icarus.2007.03.024>, deep Impact Mission to Comet 9P/Tempel 1, Part 2, 2007.

Adams, N. and Smith, D.: Studies of microparticle impact phenomena leading to the development of a highly sensitive micrometeoroid 5 detector, *Planetary and Space Science*, 19, 195 – 204, [https://doi.org/10.1016/0032-0633\(71\)90199-1](https://doi.org/10.1016/0032-0633(71)90199-1), 1971.

Antonsen, T.: In-situ Measurements of Mesospheric Aerosols – On the observable characteristics of nanoscale ice and meteoric smoke particles, Ph.D. thesis, UiT The Arctic University of Norway, Tromsø, 2019.

Antonsen, T. and Havnes, O.: On the detection of mesospheric meteoric smoke particles embedded in noctilucent cloud particles with rocket- 10 borne dust probes, *Review of Scientific Instruments*, 86, 033 305, <https://doi.org/10.1063/1.4914394>, 2015.

10 Antonsen, T., Havnes, O., and Mann, I.: Estimates of the Size Distribution of Meteoric Smoke Particles From Rocket-Borne Impact Probes, *Journal of Geophysical Research: Atmospheres*, <https://doi.org/10.1002/2017JD027220>, 2017.

Auer, A. and Sitte, K.: Detection technique for micrometeoroids using impact ionization, *Earth and Planetary Science Letters*, 4, 178 – 183, 15 [https://doi.org/10.1016/0012-821X\(68\)90013-7](https://doi.org/10.1016/0012-821X(68)90013-7), 1968.

Auer, S.: Chapter 5 in Interplanetary dust, Edited by: Grün, E., Gustafson, B., Dermott, S. and Fechtig, H., Springer-Verlag Berlin Heidelberg, 15 <https://doi.org/10.1007/978-3-642-56428-4>, 2012.

Auty, R. P. and Cole, R. H.: Dielectric Properties of Ice and Solid D2O, *The Journal of Chemical Physics*, 20, 1309–1314, <https://doi.org/10.1063/1.1700726>, 1952.

Collette, A., Grün, E., Malaspina, D., and Sternovsky, Z.: Micrometeoroid impact charge yield for common spacecraft materials, *Journal of Geophysical Research: Space Physics*, 119, 6019–6026, <https://doi.org/10.1002/2014JA020042>, 2014.

20 Copley, M. J. and Phipps, T. E.: The Surface Ionization of Potassium on Tungsten, *Phys. Rev.*, 48, 960–968, <https://doi.org/10.1103/PhysRev.48.960>, 1935.

Dahneke, B.: The influence of flattening on the adhesion of particles, *Journal of Colloid and Interface Science*, 40, 1 – 13, [https://doi.org/10.1016/0021-9797\(72\)90168-3](https://doi.org/10.1016/0021-9797(72)90168-3), 1972.

Draine, B. T. and Sutin, B.: Collisional Charging of Interstellar Grains, *Astrophysical Journal*, 320, 803, <https://doi.org/10.1086/165596>, 25 1987.

Drapatz, S. and Michel, K.: Theory of shock-wave ionization upon high-velocity impact of micrometeorites, *Zeitschrift für Naturforschung A*, 29, 870–879, <https://doi.org/10.1515/zna-1974-0606>, 1974.

Dresser, M. J.: The Saha-Langmuir Equation and its Application, *Journal of Applied Physics*, 39, 338–339, <https://doi.org/10.1063/1.1655755>, 1968.

30 Evans, A.: The Dusty Universe, Series in astronomy, John Wiley & Sons, 1994.

Evans, B. and Goetze, C.: The temperature variation of hardness of olivine and its implication for polycrystalline yield stress, *Journal of Geophysical Research: Solid Earth*, 84, 5505–5524, <https://doi.org/10.1029/JB084iB10p05505>, 1979.

Fedorov, A. V., Shul'gin, A. V., and Lavruk, S. A.: Investigation of the physical properties of iron nanoparticles in the course of the melting and solidification, *Physics of Metals and Metallography*, 118, 572–578, <https://doi.org/10.1134/S0031918X17040020>, 2017.

35 Froeschke, S., Kohler, S., Weber, A. P., and Kasper, G.: Impact fragmentation of nanoparticle agglomerates, *Journal of Aerosol Science*, 34, 275 – 287, [https://doi.org/10.1016/S0021-8502\(02\)00185-4](https://doi.org/10.1016/S0021-8502(02)00185-4), 2003.

Havnes, O. and Hartquist, T. W.: Nanodust shedding and its potential influence on dust-related phenomena in the mesosphere, *Journal of Geophysical Research: Atmospheres*, pp. 12 363–12 376, <https://doi.org/10.1002/2016JD025037>, 2016.

Havnes, O. and Næsheim, L. I.: On the secondary charging effects and structure of mesospheric dust particles impacting on rocket probes, *Annales Geophysicae*, 25, 623–637, <https://doi.org/10.5194/angeo-25-623-2007>, 2007.

5 Havnes, O., Gumbel, J., Antonson, T., Hedin, J., and Hoz, C. L.: On the size distribution of collision fragments of NLC dust particles and their relevance to meteoric smoke particles, *Journal of Atmospheric and Solar-Terrestrial Physics*, 118, 190–198, <https://doi.org/10.1016/j.jastp.2014.03.008>, 2014.

Havnes, O., Latteck, R., Hartquist, T. W., and Antonson, T.: First simultaneous rocket and radar detections of rare low summer mesospheric clouds., *Geophysical Research Letters*, 45, 5727–5734, <https://doi.org/10.1029/2018GL078218>, 2018.

10 Havnes, O., Antonson, T., Baumgarten, G., Hartquist, T. W., Biebricher, A., Fredriksen, Å., Friedrich, M., and Hedin, J.: A new method of inferring the size, number density, and charge of mesospheric dust from its in situ collection by the DUSTY probe, *Atmospheric Measurement Techniques*, 12, 1673–1683, <https://doi.org/10.5194/amt-12-1673-2019>, 2019.

Hervig, M. E., Deaver, L. E., Bardeen, C. G., III, J. M. R., Bailey, S. M., and Gordley, L. L.: The content and composition of meteoric smoke in mesospheric ice particles from SOFIE observations, *Journal of Atmospheric and Solar-Terrestrial Physics*, 84-85, 1 – 6, <https://doi.org/10.1016/j.jastp.2012.04.005>, 2012.

15 John, W., Reischl, G., and Devor, W.: Charge transfer to metal surfaces from bouncing aerosol particles, *Journal of Aerosol Science*, 11, 115 – 138, [https://doi.org/10.1016/0021-8502\(80\)90029-4](https://doi.org/10.1016/0021-8502(80)90029-4), 1980.

Jones, A. P., Tielens, A. G. G. M., and Hollenbach, D.: Grain Shattering in Shocks: The Interstellar Grain Size Distribution, *The Astrophysical Journal*, 469, 740–764, <https://doi.org/10.1086/177823>, 1996.

20 Kissel, J. and Krueger, F. R.: Ion formation by impact of fast dust particles and comparison with related techniques, *Applied Physics A*, 42, 69–85, <https://doi.org/10.1007/BF00618161>, 1987.

Klinkrad, H.: Space Debris, John Wiley and Sons, <https://doi.org/10.1002/9780470686652.eae325>, 2010.

Kuznetsov, N. and Raizer, Y. P.: Recombination of electrons in a plasma expanding into a vacuum, *Journal of Applied Mechanics and Technical Physics*, 6, 6–12, 1965.

25 MacDowell, L. G. and Vega, C.: Dielectric Constant of Ice Ih and Ice V: A Computer Simulation Study, *The Journal of Physical Chemistry B*, 114, 6089–6098, <https://doi.org/10.1021/jp100167y>, 2010.

Mann, I., Nouzák, L., Vaverka, J., Antonson, T., Fredriksen, Å., Issautier, K., Malaspina, D., Meyer-Vernet, N., Pavlů, J., Sternovsky, Z., Stude, J., Ye, S., and Zaslavsky, A.: Dust observations with antenna measurements and its prospects for observations with Parker Solar Probe and Solar Orbiter, *Annales Geophysicae*, 37, 1121–1140, <https://doi.org/10.5194/angeo-37-1121-2019>, 2019.

30 Matsusaka, S., Maruyama, H., Matsuyama, T., and Ghadiri, M.: Triboelectric charging of powders: A review, *Chemical Engineering Science*, 65, 5781 – 5807, <https://doi.org/10.1016/j.ces.2010.07.005>, 2010.

McBride, N. and McDonnell, J.: Meteoroid impacts on spacecraft:: sporadics, streams, and the 1999 Leonids, *Planetary and Space Science*, 47, 1005 – 1013, [https://doi.org/10.1016/S0032-0633\(99\)00023-9](https://doi.org/10.1016/S0032-0633(99)00023-9), 1999.

35 Mocker, A., Hornung, K., Grün, E., Kempf, S., Collette, A., Drake, K., Horányi, M., Munsat, T., O'Brien, L., Sternovsky, Z., and Srama, R.: On the application of a linear time-of-flight mass spectrometer for the investigation of hypervelocity impacts of micron and sub-micron sized dust particles, *Planetary and Space Science*, 89, 47 – 57, <https://doi.org/10.1016/j.pss.2013.07.013>, 2013.

Nimmo, F.: What is the Young's Modulus of Ice?, in: *Workshop on Europa's Icy Shell: Past, Present, and Future*, edited by Schenk, P., Nimmo, F., and Prockter, L., p. 7005, 2004.

Nunez-Valdez, M., Umemoto, K., and Wentzcovitch, R. M.: Fundamentals of elasticity of $(\text{Mg}_{1-x}\text{Fe}_x)\text{SiO}_4$ olivine, *Geophysical Research Letters*, 37, L14 308:1–5, <https://doi.org/10.1029/2010GL044205>, 2010.

Omont, A.: Physics and chemistry of interstellar polycyclic aromatic molecules., *Astronomy and Astrophysics*, 164, 159–178, https://doi.org/10.1007/978-94-009-4776-4_35, 1986.

5 Page, B., Bale, S. D., Bonnell, J. W., Goetz, K., Goodrich, K., Harvey, P. R., Larsen, R., MacDowall, R. J., Malaspina, D. M., Pokorný, P., Pulupa, M., and Szalay, J. R.: Examining Dust Directionality with the Parker Solar Probe FIELDS Instrument, *The Astrophysical Journal Supplement Series*, 246, 51, <https://doi.org/10.3847/1538-4365/ab5f6a>, 2020.

Pokorný, P. and Kuchner, M.: Co-orbital Asteroids as the Source of Venus' Zodiacal Dust Ring, *The Astrophysical Journal*, 873, L16, <https://doi.org/10.3847/2041-8213/ab0827>, 2019.

10 Qi, W. and Wang, M.: Size effect on the cohesive energy of nanoparticle, *Journal of Materials Science Letters*, 21, 1743–1745, <https://doi.org/10.1023/A:1020904317133>, 2002.

Raizer, Y. P.: Residual ionization of a gas expanding in vacuum, *Sov. Phys. JETP*, 10, 411–416, http://jetp.ac.ru/cgi-bin/dn/e_010_02_0411.pdf, (Retrieved: 04.12.19), 1960.

15 Rapp, M., Plane, J. M. C., Strelnikov, B., Stober, G., Ernst, S., Hedin, J., Friedrich, M., and Hoppe, U.-P.: In situ observations of meteor smoke particles (MSP) during the Geminids 2010: constraints on MSP size, work function and composition, *Annales Geophysicae*, pp. 1661–1673, 2012.

Ratcliff, P., Reber, M., Cole, M., Murphy, T., and Tsemelis, K.: Velocity thresholds for impact plasma production, *Advances in Space Research*, 20, 1471 – 1476, [https://doi.org/10.1016/S0273-1177\(97\)00419-5](https://doi.org/10.1016/S0273-1177(97)00419-5), hypervelocity Impacts in Space and Planetology, 1997.

20 Rennecke, S. and Weber, A. P.: Charge Transfer to Metal Nanoparticles Bouncing from Conductive Surfaces, *Aerosol Science and Technology*, 48, 1059–1069, <https://doi.org/10.1080/02786826.2014.955566>, 2014.

Rizk, B., Hunten, D. M., and Engel, S.: Effects of size-dependent emissivity on maximum temperatures during micrometeorite entry, *Journal of Geophysical Research: Space Physics*, 96, 1303–1314, <https://doi.org/10.1029/90JA01998>, 1991.

Schildknecht, T.: Optical surveys for space debris, *The Astronomy and Astrophysics Review*, 14, 41–111, 2007.

25 Shibata, Y. and Suzuki, T.: A molecular dynamics study of cooling rate during solidification of metal nanoparticles, *Chemical Physics Letters*, 502, 82–86, <https://doi.org/10.1016/j.cplett.2010.12.020>, 2011.

Soo, S. L.: Dynamics of Charged Suspensions, in: *Topics in Current Aerosol Research*, edited by Hidy and Brock, vol. 11 of *International Reviews in aerosol physics*, p. 61, pergammon, <https://doi.org/10.1016/B978-0-08-016674-2.50011X>, 1971.

30 St. Cyr, O. C., Kaiser, M. L., Meyer-Vernet, N., Howard, R. A., Harrison, R. A., Bale, S. D., Thompson, W. T., Goetz, K., Maksimovic, M., Bougeret, J. L., Wang, D., and Crothers, S.: STEREO SECCHI and S/WAVES Observations of Spacecraft Debris Caused by Micron-Size Interplanetary Dust Impacts, *Solar Physics*, 256, 475–488, <https://doi.org/10.1007/s11207-009-9362-5>, 2009.

Strelnikov, B., Rapp, M., Strelnikova, I., Chau, J. L., Latteck, R., Luebken, F.-J., Friedrich, M., Asmus, H., Renkwitz, T., and Staszak, T.: PMWE-1 sounding rocket campaign, in: 42nd COSPAR Scientific Assembly, vol. 42, 2018.

35 Szalay, J. R., Pokorný, P., Bale, S. D., Christian, E. R., Goetz, K., Goodrich, K., Hill, M. E., Kuchner, M., Larsen, R., Malaspina, D., McComas, D. J., Mitchell, D., Page, B., and Schwadron, N.: The Near-Sun Dust Environment: Initial Observations from Parker Solar Probe, *The Astrophysical Journal Supplement Series*, 246, 27, <https://doi.org/10.3847/1538-4365/ab50c1>, 2020.

Thomas, E., Simolka, J., DeLuca, M., Horányi, M., Janches, D., Marshall, R. A., Munsat, T., Plane, J. M. C., and Sternovsky, Z.: Experimental setup for the laboratory investigation of micrometeoroid ablation using a dust accelerator, *Review of Scientific Instruments*, 88, 034 501, <https://doi.org/10.1063/1.4977832>, 2017.

Tomsic, A., Andersson, P. U., Markovic, N., Piskorz, W., Svanberg, M., and Pettersson, J. B. C.: Molecular dynamics simulations of cluster-surface collisions: Emission of large fragments, *The Journal of Chemical Physics*, 115, 10 509–10 517, <https://doi.org/10.1063/1.1413740>, 2001.

Tomsic, A., Schröder, H., Kompa, K.-L., and Gebhardt, C. R.: Impact dynamics of molecular clusters on surfaces: Fragmentation patterns and anisotropic effects, *The Journal of Chemical Physics*, 119, 6314–6323, <https://doi.org/10.1063/1.1603213>, 2003.

5 Vourlidas, A., Howard, R. A., Plunkett, S. P., Korendyke, C. M., Thernisien, A. F., Wang, D., Rich, N., Carter, M. T., Chua, D. H., Socker, D. G., et al.: The wide-field imager for Solar Probe Plus (WISPR), *Space Science Reviews*, 204, 83–130, <https://doi.org/10.1007/s11214-014-0114-y>, 2016.

10 Wang, H.-C. and John, W.: Dynamic contact charge transfer considering plastic deformation, *Journal of Aerosol Science*, 19, 399 – 411, [https://doi.org/10.1016/0021-8502\(88\)90016-X](https://doi.org/10.1016/0021-8502(88)90016-X), 1988.

Wood, D. M.: Classical Size Dependence of the Work Function of Small Metallic Spheres, *Phys. Rev. Lett.*, 46, 749–749, <https://doi.org/10.1103/PhysRevLett.46.749>, 1981.

Zel'Dovich, Y. and Raizer, Y.: Physics of shock waves and high-temperature hydrodynamic phenomena, Academic Press, <https://doi.org/10.1016/B978-0-12-395672-9.X5001-2>, 1967.