

## ***Interactive comment on “Multipoint Observations of Compressional Pc5 Pulsations in the Dayside Magnetosphere and Corresponding Particle Signatures” by Galina Korotova et al.***

**Galina Korotova et al.**

gkorotov@umd.edu

Received and published: 17 August 2020

We included three new figures from the paper and some description of figure 4.

Figure 4a shows G-13 and -15 observations of the total magnetic field strength from 18:00 UT to 24:00 UT. The spacecraft observed long-duration Pc5 pulsations over a wide longitudinal region in the pre- and post-noon magnetosphere from 10:00 to 15:20 MLT (Figure 2). G-15 observed weak, less than  $\sim 5$  nT amplitude, Pc5 waves from 18:28 UT to 19:04 UT prior to the main event. During the main event from 19:04 to 23:00 UT, the magnetosphere was compressed (Figure 3), magnetic field strengths increased and the amplitude of these waves increased to values ranging from 10 to 16

C1

nT with peak amplitudes prior to local noon. G-13 observed weak Pc5 pulsations with amplitudes of 2-4 nT throughout most of the time interval from 16:40 UT (not shown) to 21:00 UT. During the interval from 19:34 UT ( $\sim 14:45$  MLT) to 20:10 UT ( $\sim 15:20$  MLT), the pulsations reached slightly stronger amplitudes of 5-8 nT. At 23:02 UT all Pc5 wave activity at both GOES stopped. Figure 4b shows the RBSP-A and -B total magnetic field strength from 18:40 UT to 21:10 UT and from 20:40 UT to 23:10 UT, respectively, on January 1, 2016. Taken together, RBSP-A and -B observed Pc5 pulsations that occupied the inner dayside magnetosphere from 5.26 to 5.75 RE and from 09:56 to 12:44 MLT (Figure 2). Prior to the arrival of the strong solar wind dynamic pressure variations from 18:15 to 18:55 UT RBSP-A observed very weak pulsations with Pc5 periods and amplitudes of 1-3 nT (not visible at this scale). After the compression of the magnetosphere just after 19:00 UT, the pulsation amplitude at RBSP-A increased to values ranging from 10 to 15 nT with the peak amplitude occurring prior to local noon (Figure 4b). RBSP-B observed similar compressional Pc5 pulsations from 20:46 UT that ceased simultaneously with the end of the magnetospheric compression at about 23:02 UT.

---

Interactive comment on Ann. Geophys. Discuss., <https://doi.org/10.5194/angeo-2020-32>, 2020.

C2






Figure 1. Bz component of the magnetic field observed at Wind, and geomagnetic activity obtained from the OMNI database (upper panels) from 12:00 UT on December 30 to 00:00 UT on January 1, 2016. The bottom panels show Wind observations of the magnetic field components, total magnetic field strength, pressure, plasma density, and velocity from 16:00 UT on January 1, 2016 to 00:00 UT on January 1, 2016.



Figures 4 (a, b). G-15 and G-13 (a) total magnetic field strength from 18:00 UT to 23:00 UT on January 1, 2016. RBSP-A and -B (b) total magnetic field strength from 18:40 UT to 23:10 UT on January 1, 2016, respectively. Beneath the universal time (UT) and magnetic local time (MLT).



Figures 10 (a, b). RBSP-A observations of electron fluxes (a) in the energy range from 31.5 keV to 1704 keV from 18:30 UT to 21:00 UT and (b) their expanded view from 19:20 UT to 20:00 UT.