

Interactive comment on “Multipoint Observations of Compressional Pc5 Pulsations in the Dayside Magnetosphere and Corresponding Particle Signatures” by Galina Korotova et al.

Galina Korotova et al.

gkorotov@umd.edu

Received and published: 18 August 2020

Dear Referee,

We can not submit a revised manuscript now and think that information on slightly changed Figure 5 may be useful.

To determine the type of the Pc5 waves we converted the magnetic field observations from GSE into field-aligned coordinates (FAC). Here the Z axis lies parallel to the locally-averaged magnetic field. The Y axis points approximately azimuthally eastward and is transverse to B and to the outward radius vector. The X axis completes the

[Printer-friendly version](#)

[Discussion paper](#)




right-handed system and is directed approximately radially outward from Earth. Figure 5 presents RBSP-A and -B magnetic field observations in FAC. The B_z component is the value of the total magnetic field after subtraction of a 16-minute sliding average. The Pc5 pulsations are observed in all three components but the amplitudes of the azimuthal B_y and radial B_x components are rather small and do not exceed 7 nT. The compressional B_z component is much more pronounced for both spacecraft, reaching amplitudes of 14-15 nT before local noon, consequently, the pulsations are primarily compressional. The B_z component oscillated out of phase with the B_x component at RBSP-A and in phase at RBSP-B and in quadrature with the B_y component.

Interactive comment on Ann. Geophys. Discuss., <https://doi.org/10.5194/angeo-2020-32>, 2020.

[Printer-friendly version](#)

[Discussion paper](#)





[Printer-friendly version](#)

[Discussion paper](#)



Figure 5. RBSP-A and -B magnetic field observations in field-aligned coordinates from 18:40

UT to 21:10 UT and from 20:40 UT to 23:10 UT on January 1, 2016, respectively.