

Journal: Annales Geophysicae

Title: Atmospheric drag effects on modelled LEO satellites during the July 2000 Bastille Day event in contrast to an interval of geomagnetically quiet conditions

Author(s): Victor U. J. Nwankwo et al.

Manuscript Number: angeo-2020-33

Manuscript Type: Regular paper

Iteration: Revision

17-10-2020

Dear Sir,

Thank you so much for your time and for organising the effort that is taking this manuscript through the stages making it a better and quality paper. We are also thankful to the reviews for their time and helpful comments and suggestions.

We have now revised our manuscript based on your suggestions, the reviewer comments and the answers we provided. While the revision lasted, we aimed at improving the paper's quality as much as possible in order to meet the standard of your reputed Journal. The point-by-point reply to the Editor and Reviewer's comments are detailed below:

### **EDITOR'S COMMENTS**

Based on the reviewer comments and your answers, I invite you to submit a revised version of your manuscript. Please do take into account the detailed reviewer comments while preparing your revision. In particular, make sure that it is clear which contributions of your paper are new.

#### ***Authors Response***

We have now clarified the contributions in this paper that are new as much as possible. To better track the clarifications, the reviewer comments and required adjustments are also provided (below). However, we note specific portions of the manuscript, which pin-points such contributions

- 1) In the Abstract: Besides the former inclusion (e.g., line 16-19), we buttressed and/or added to the contributions in line 19-21
- 2) In Section 1.2: In addition to the important detail provided under "Relevance of the study and its application" (e.g., line 116-119) we further provided more, as well as the justification for focusing on the Bastille Day event in line 120-133
- 3) In Section 4: line 316-322.

### **COMMENTS BY REFEREE #1**

The paper is dealing with the impact of atmospheric drag on LEO satellites, which is a complex problem. This work does not contain any new ideas and the authors already acknowledge this fact in their abstract! Their methodology and some applications have been already presented in previous work. However, some of the data presented in the paper can be considered as new since the authors examine some specific cases. Overall, the paper does not provide significant scientific contribution, especially when compared with similar work from the literature.

#### ***Authors Response***

Although the reviewer admitted that some of the data we presented in the paper are new, he stopped short of acknowledging the accomplishment of this effort. The referee feels that our paper does not provide

significant scientific contribution. We respect the judgment of the referee and do not take it as being biased. However, we beg to differ on the claim. As the goal of manuscript review (by Referees) are generally to 'provide unbiased and constructive comments aimed, whenever possible, at improving the work,' we feel that a generous suggestion of content-wise inclusion or modification that will increase quality would have been a better conclusion of the referee. To address the comment of the referee, I think elucidating the new scientific contributions of this paper in clear terms (as also suggested by the Editor) is the best thing to do. We therefore highlight them below (as was done earlier via our response to RC1).

(1) This paper emphasized and/or focused on the Bastille Day great geomagnetic storm (and associated phenomena). It is hoped that efforts directed towards assessing, monitoring, modeling and/or prediction of the impacts associated with sudden severe solar energetic transients (like this one) are key to mitigating the potential threat posed by such event in future occurrence. This is the first time we are modeling atmospheric drag effect associated with the Bastille Day event (BDE). Therefore, this paper increases the visibility and better contribute to the scientific body of knowledge surrounding the BDE (as earlier stated). Please see the specific portions above where we provided the detail of this contribution (and others) in the text, as well as the justification for focusing on the BDE.

(2) In our analysis we used new method and indices to describe and estimate drag effects on the satellite trajectory when contrasting between the (i) solar active and the quiescent regimes (ii) active regime and the Bastille day storm, and (iii) the quiescent regime and the Bastille day event/storm. This analysis and the results obtained is now helping us to produce estimation model that compares effects between regimes of varying solar-geomagnetic activity. In addition to examining a specific case (different from previous study), we used a relatively new (or novel) approach/method. As much as I prefer to keep a low profile on this at this stage, I am yet to find similar approach in literature till date.

(3) This work doubles as a strong review paper. We presented extensive details/review on atmospheric drag (and its relevance) in relation to solar activity, against properly referenced background of existing work. If carefully perused, one could see a concise comprehensive connection between atmospheric drag and solar-geomagnetic activity that is particularly unprecedented when compared with our previous work (not overall literature in the area) – thanks to the proficiency of some co-authors!

In summary, our pattern of result presentation (especially in abstract) may look similar but we believe the accomplishments or contributions of this effort are replete and should not be overemphasized (especially when one carefully read beyond the abstract).

## **COMMENTS BY REFEREE #2**

1. The goal of the paper i.e The goal of this effort was to quantify how solar-geomagnetic activity influences atmospheric drag and perturbs satellite orbits, is very clear and worthwhile. The authors focused on the Bastille event because they have done similar work in another paper. I do not agree with the use of word as the authors seem to infer that they have not done any work different from the paper they first published on the topic. I reckon that the authors should have shown distinct comparison between the current paper and the previous paper and strongly justify why they focused on the bastille event.

### ***Authors Response***

We have now revised the manuscript in the manner that buttressed the scientific contribution in this work that are new and also different from previous work (please also see the above highlight). This way the 'distinct comparison between the current paper and the previous paper' can be clearly understood. We have also justified the reason we focused on the Bastille event. The specific portions of the manuscript which pin-points such inclusions are listed below

- 1) In the Abstract: line 19-21 (in addition to line 16-19)
- 2) In Section 1.2: line 120-133 (In addition to line 116-119)
- 3) In Section 4: line 316-322.

2. I am satisfied with the response from the author. I reckon that the authors could include the response, i.e "this work also doubled as a strong review paper because it presented extensive details/review on atmospheric drag (and its relevance) in relation to solar activity, against properly referenced background of existing work. The significant number of readers who have interacted with this manuscript on this platform (and others) certainly did because of its relevance to them. I am also aware of authors have cited this paper in their new manuscript" on the introduction. This will guide readers and clearly underpin the objectives of the present communication.

**Response** Please see line 131-133 for the inclusion. We excluded a few lines since this is a scientific article

It will be good for the authors to spotlight the comparison of the Bastille day event to existing result right from the abstract.

**Response** Please see line 3-4, 18, 19-22

I advise that the part of the abstract that sort of infer that there are no new results in the paper should be taken off as it is grossly misleading if the authors claim that "this analysis motivated the development of new method and indices for description and estimation of drag effects on satellite ephemeris

**Response** The suggested phrase have now been removed from the abstract.

How do you justify the below within in he current paper. You are supposed to convincingly show the strength in this paper for reviewing purposes. "We are now in the process of combining satellite drag model high-fidelity atmospheric specification to produce such realistic estimation model (beginning with the results of this work".

**Response** In addition to examining a specific case (the BDE) that is different from previous study (with new results), one other strength of this work is that we used a relatively new approach that is now helping us to produce estimation model that compares effects between regimes of varying solar-geomagnetic activity – and such formulation is a process!

In conclusion, we have also closely perused the manuscript again to eliminate or reduce typographical errors and expressions that can make the understanding of any portion difficult for the readers (as much as possible). We believe that in its current state, our revised manuscript is now suitable for further consideration by your journal, and sincerely hope that the paper will now be accepted for publication.

Thank you very much.

Victor U. J. Nwankwo

# Atmospheric drag effects on modelled LEO satellites during the July 2000 Bastille Day event in contrast to an interval of geomagnetically quiet conditions

Victor U. J. Nwankwo<sup>1</sup>, William Denig<sup>2</sup>, Sandip K. Chakrabarti<sup>3</sup>, Muyiwa P. Ajakaiye<sup>1</sup>, Johnson Fatokun<sup>1</sup>, Adeniyi W. Akanni<sup>1</sup>, Jean-Pierre Raulin<sup>4</sup>, Emilia Correia<sup>4</sup>, and John E. Enoh<sup>5</sup>

<sup>1</sup>AUL Space Laboratory, Anchor University, Lagos 100278, Nigeria

<sup>2</sup>St. Joseph College of Maine, Standish, ME 04084, U.S.A

<sup>3</sup>Indian Centre for Space Physics, Kolkata 700084, India

<sup>4</sup>Centro de Rádio Astronomia e Astrofísica Mackenzie, Universidade Presbiteriana Mackenzie, São Paulo, Brazil

<sup>5</sup>Interorbital systems, Mojave, CA 93502-0662, U.S.A.

**Correspondence:** Victor U. J. Nwankwo (vnwankwo@aul.edu.ng)

**Abstract.** In this work we simulated atmospheric drag effect on two model SmallSats in Low Earth Orbit (LEO) with different ballistic coefficients during 1-month intervals of solar-geomagnetic quiet and perturbed conditions. The goal of this effort was to quantify how solar-geomagnetic activity influences atmospheric drag and perturbs satellite orbits, **with particular emphasis on the Bastille Day event**. Atmospheric drag compromises satellite operations due to increased ephemeris errors, attitude positional uncertainties and premature satellite re-entry. During a 1-month interval of generally quiescent solar-geomagnetic activity (July 2006) the decay in altitude ( $h$ ) was a modest 0.53 km (0.66 km) for the satellite with the smaller (larger) ballistic coefficient of  $2.2 \times 10^{-3} \text{ m}^2/\text{kg}$  ( $3.03 \times 10^{-3} \text{ m}^2/\text{kg}$ ). The associated Orbital Decay Rates (ODRs) during this quiet interval ranged from 13 m/day to 23 m/day (from 16 m/day to 29 m/day). For the disturbed interval of July 2000 the significantly increased altitude loss and range of ODRs were 2.77 km (3.09 km) and 65 m/day to 120 m/day (78 m/day to 142 m/day), respectively. Within the two periods more detailed analyses over 12-day intervals of extremely quiet and disturbed conditions revealed respective orbital decays of 0.16 km (0.20 km) and 1.14 km (1.27 km) for the satellite with the smaller (larger) ballistic coefficient. In essence, the model results show that there was a 6-7 fold increase in the deleterious impacts of satellite drag between the quiet and disturbed periods. We also estimated the enhanced atmospheric drag effect on the satellites' parameters caused by the July 2000 Bastille Day event (in contrast to the interval of geomagnetically quiet conditions). The additional percentage increase due to the Bastille Day event to the monthly mean values of  $h$  and ODR are 34.69% and 50.13% for Sat-A, and 36.45% and 68.95% for Sat-B. These simulations confirmed; (i) the dependence of atmospheric drag force on a satellite's ballistic coefficient, and (ii) that increased solar-geomagnetic activity substantially raises the degrading effect of satellite drag. In addition, the results indicate that the impact of short-duration geomagnetic transients (**such as the Bastille Day storm**) can have a further deleterious effect on normal satellite operations. **Thus this work increased the visibility and contribute to the scientific knowledge surrounding the Bastille Day event, and also motivated the introduction of new indices used to describe and estimate atmospheric drag effect when comparing between regimes of varying solar-geomagnetic activity.**

We suggest that a model of satellite drag when combined with a high-fidelity atmospheric specification, as was done here, can lead to improved satellite ephemeris estimates.

*Copyright statement.* TEXT


## 25 1 Introduction

Atmospheric drag describes the force exerted on an object moving through the atmospheric medium. The orientation of the drag force is in the reverse direction of relative motion with the resulting effect of impeding the motion of the object. Spacecraft moving through the atmosphere experience the atmospheric drag force which expends energy at the expense of the orbital motion (Wertz and Larson, 1999; Chobotov, 2002; Nwankwo, 2016). Atmospheric drag is the largest force affecting the motion of satellites in Low Earth Orbit (LEO) especially at altitudes below 800 km (Nwankwo et al., 2015). Space weather enhances atmospheric drag on satellites in LEO and the resultant impact can be profound (Nwankwo, 2016). Extreme space weather can cause satellite orbits to unexpectedly degrade making it more difficult to maneuver spacecraft and to identify and track satellites and other space debris (Nwankwo et al. (2015); and references therein). Another detrimental impact of enhanced satellite drag is the unplanned loss of otherwise healthy spacecraft due to premature atmospheric re-entry. Under this scenario a satellite would gradually decay from orbit (losing altitude) and would re-enter the earth's lower atmosphere unless appropriate orbit-raising maneuvers were implemented. Examples of spacecraft that re-entered the atmosphere include Skylab (launched 14 May 1973, re-entered 11 July 1979) and the Russian Radar Ocean Reconnaissance Satellites (RORSATs), Kosmos-954 (launched 18 September 1977, re-entered 24 January 1978) and Kosmos-1402 (launched 30 August 1982, re-entered 07 February 1983) (Nwankwo, 2016).

40

The orbital lifetime of a LEO satellite is subjected to the integrated atmospheric drag force experienced by the satellite over time. Figure 1 depicts the orbital degradation of a hypothetical satellite in a nominally circular orbit which, in this case, degrades over time from a starting position "A" at an altitude of 400 km to a later position "B" at an altitude below 110 km. The drag or negative acceleration,  $f_d$ , (units of  $\text{m/s}^2$ ) experienced by the satellite is given as  $f_d = \frac{1}{2}\rho B v_s^2$  where  $\rho$  (units of  $\text{kg/m}^3$ ) is the altitude-dependent atmospheric density and  $v_s$  (units of  $\text{m/s}$ ) is the satellite speed (King-Hele, 1987). The satellite speed,  $v_s$ , is a consequence of the balance between an inward-directed (towards earth) gravitational force at the satellite altitude and the outward-directed orbital centripetal force. A simplified version of a satellite's ballistic coefficient,  $B$  (units of  $\text{m}^2/\text{kg}$ ), is given as  $B = C_d A_s / m_s$  where  $C_d$  is the unit-less atmospheric drag coefficient,  $A_s$  (units of  $\text{m}^2$ ) is the satellite's projected area in the direction of motion and  $m_s$  (units of  $\text{kg}$ ) is the satellite mass (Bowman, 2002; Bhatnagar et al., 2005). For altitudes representative of most LEO satellites, that being between 140 km to 400-600 km, a constant drag coefficient,  $C_d$ , of 2.2 is appropriate (Cooke, 1965). A drag force,  $F$  (units of  $\text{kg m/s}^2$ ), acting in opposition to the satellite's motion, is given as  $F = -m_s f_d$ . In this work we model changes in the Orbital Decay Rate (ODR, units of  $\text{m/day}$ ) and the monthly mean orbital

50



**Figure 1.** Orbital decay scenario of a LEO satellite due to atmospheric drag. The satellite orbit decreases from altitude “A” to altitude “B” as a consequence of atmospheric drag (Adapted from Nwankwo (2016)).


decay (units of km) experienced by 2 satellites having different ballistic coefficients,  $B$ , under different solar-geomagnetic conditions (Nwankwo et al., 2020).

### 55 1.1 Solar and geomagnetic activities and their implications for atmospheric drag

Solar activity describes changes in the overall energy and mass output from the sun consisting of both long-term trends within the 11-year solar cycle (longer term changes are beyond the scope of this effort) and transient events of increased solar output. Electromagnetic radiation (light) is continuously emitted from the sun across a broad spectral range from energetic gamma rays to radio-waves (Eddy, 2009). Also emitted from the sun are the streams of electrons and protons which comprise the background solar wind and impulsive fluxes of energetic charged particles contained in solar energetic particle (SEP) events (Parker, 1958; Ryan et al., 2000). Charged particle gases from the sun are classified as high-beta plasmas within which the remnants of solar magnetic fields are transported towards Earth and can interact with the geomagnetic field. The transported field is referred to as the Interplanetary Magnetic Field (IMF) and the orientation of this field relative to Earth has a controlling effect on the degree of coupling of the solar wind and/or the transient streams with the magnetosphere; that is, the earth’s outer magnetic shielding layer that acts to protect the terrestrial biosphere from interplanetary energetic charged particles (Schatten, 1971; Yermolaev et al., 2018). However, as the solar streams, with their embedded magnetic fields, impact the magnetosphere they can enhance geomagnetic activity which, in turn, can have a significant effect on the coupled Magnetosphere-Ionosphere-Thermosphere (MIT) system. For example, within interplanetary space, a solar High-Speed Stream (HSS) can overtake a

preceding Low-Speed Stream (LSS) thus forming a dense Corotating Interaction Region (CIR) (Gosling and Pizzo, 1999) that  
70 can profoundly increase the level of geomagnetic activity. The sun also periodically releases large-scale “clouds” of plasma in  
the form of a Coronal Mass Ejection (CMEs), which, when propagated into interplanetary space, is termed an Interplanetary  
CME (ICME) (Gosling et al., 1990). When directed towards earth, CIRs and ICMEs can initiate geomagnetic storms resulting  
in large-scale perturbations of the MIT system lasting up to several days (Borovsky and Denton, 2006). Flares represent another  
class of transient solar phenomena which can affect the MIT system. A solar flare is a large scale (on solar dimensions) recon-  
75 figuration of the photospheric magnetic field resulting in the impulsive release of vast amounts of energy and a redistribution  
of solar mass (Philips, 1991). Electromagnetic radiation and extremely energetic (relativistic) particles released during a solar  
flare event can result in an abrupt increase in the ionospheric density near the subsolar point and within the high-latitude polar  
caps (Sauer and Wilkinson, 2008). A Sudden Ionospheric Disturbance (SID) is the result of the increased solar UltraViolet  
(UV) and X-ray radiative flux released in solar flares (Mitra, 1974) whereas a Polar Cap Absorption (PCA) event is the result  
80 of energetic particles entering the atmosphere along “open” magnetic field lines connect to interplanetary space (Rose and  
Ziauddin, 1962).

The systematic monitoring of sunspots over the last two centuries has shown that solar activity exhibits an approximate  
11-year temporal cycle during which the observed SunSpot Number (SSN) (Clette et al., 2014) varies from a local solar  
85 minimum near 0; that is, no spots observed, to a solar maximum of up to several hundred spots visible on the solar disk. Near  
solar maximum the total radiant energy from the sun reaches a corresponding peak along with a propensity for short-lived  
solar transients of increased radiation and particle emissions. These solar transients are the main drivers of space weather  
(Song et al., 2001; Knipp, 2011). Figure 2 illustrates this cyclic variation in the monthly-averaged SSN along with the related  
solar-geophysical indices for the solar radio flux (F10.7) and the geomagnetic Ap. The F10.7 index (Tapping, 2013) is a local  
90 noontime measurement of the solar radio flux at a wavelength of 10.7 cm, corresponding to a radio-wave frequency of 1400  
MHz. The F10.7 index is often used as a proxy for upper atmospheric heating from solar Extreme UltraViolet (EUV) radiation.  
The F10.7 index is given in solar flux units (sfu;  $1 \text{ sfu} = 10^{-22} \text{ Wm}^{-2} \text{ Hz}^{-1}$ ) and typically ranges from  $<50$  sfu at solar  
minimum to  $>300$  sfu at solar maximum. The daily Ap index (Rostoker, 1972) is derived from the 3-hour Kp index from  
which an additional Joule heating effect associated with geomagnetic activity can be estimated. The Ap index is provided in  
95 units of nanoTeslas (nT) and typically ranges from  $\sim 5$  nT (solar minimum) to  $\sim 40$  nT (solar maximum). Near the time of  
solar maximum the increased frequency of solar transients; i.e. flares and CMEs, can lead to increased geomagnetic activity.  
As energy inputs to the MIT system, the increased solar radiation and enhanced geomagnetic activity heat the thermosphere  
and cause the atmosphere to diffuse outward from lower altitude regions of higher neutral gas pressures (densities) to the more  
tenuous upper atmosphere. The consequential increase in atmospheric drag associated with a more dense atmosphere affects  
100 the motion of a LEO satellite and expends energy at the expense of the orbit. The monthly smoothed values for F10.7 and  
Ap, as plotted in Figure 2, tend to obscure the effects of solar events and geomagnetic transients. The impacts of increased  
solar-geomagnetic activity on the atmosphere and, in turn, the atmospheric drag are discussed in Nwankwo et al. (2015); and  
references therein.



**Figure 2.** Monthly smoothed values of archived Sun Spot Number (SSN), Radio solar flux (F10.7) and geomagnetic Ap from January 1991 - March 2020 (updated predicted data in Nwankwo and Chakrabarti (2013) with actual data).

## 1.2 Relevance of the study and its application

105 Rapid variations in the local thermospheric density increase the risks of satellite collisions due to larger error margins in spacecraft positioning and motion. In 2009 a Russian satellite in orbit (Cosmos 2251) collided with a United States communications satellite (Iridium 33) at an altitude of about 800 km (Jakhu, 2009; Kelso, 2009). Cosmos 2251 was a defunct satellite in orbit whereas Iridium 33 was an operational satellite providing telecommunication services when the accident occurred. In addition to the total destruction of the satellites, this hyper-velocity collision resulted in a large increase in the amount of small, but

110 still potentially lethal, space debris. Assessing atmospheric drag-associated risk is imperative due to the increasing number of both active and expired space missions combined with a less than fully specified debris field (McCrea, private com., 2018). For example, the planned launch of new capabilities, such as SpaceX's Starlink Mega Constellation makes this subject increasingly germane to satellite operators and stakeholders. Space agencies acknowledge the potential threat posed by solar-geomagnetic activity in modulating satellite trajectories and are making strides towards addressing the issue. For example, the European

115 Space Agency (ESA) is currently assessing space-weather related risks within the framework of its Space Situational Awareness (SSA) program (Bobrinsky and Del Monte, 2010). An important mitigation approach (among others) to safeguard satellite operations is the development and implementation of models that can mimic (and assess) the impact of space weather on LEO satellite (Nwankwo et al., 2019). Accordingly, this work is of practical importance as the resulting model and simulation support efforts to increase SSA and improve collision risk mitigation.

120



It has been recommended that mitigation effort be more robust for the solar maximum phase because the rate of impact is higher during this regime (e.g., Nwankwo et al. 2020). Sudden severe solar energetic transients (SSET) that can produce huge impact are highly probable during solar maximum. One example of such SSET is the ‘the great’ geomagnetic storm of 13-14 March 1989. The event caused increase in atmospheric drag for satellites and orbital debris in low Earth orbit leading to ‘temporal loss’ of about 2500 space objects being monitored by tracking systems ((Boteler, 2019); and references therein). Another incidence of SSET is the Bastille Day event of 14-16 July 2000 during which anomalies (or disruption) on several satellites were reported (Watari et. al, 2001). The Bastille Day great geomagnetic storm (and associated phenomena) generated great interest among the scientific community due to its space weather consequences (e.g., Watari et. al (2001); Raeder et al (2001); Webber et al (2002)). Efforts directed towards assessing, monitoring, modeling and/or prediction of the impacts associated with SSET are key to mitigating the potential threat posed by such event in future occurrence. In the light of these, the present work will increase the visibility and better contribute to the scientific body of knowledge surrounding the Bastille Day event (particularly on the impact of atmospheric drag). The current paper also contain extensive details and/or review on solar (and geomagnetic) activity and its implications for atmospheric drag (modulating satellites trajectories) against properly referenced background of existing work, and therefore, doubles as a review paper.

## 2 Data, method and scope of the study

In this work we present model of LEO satellite trajectories during intervals of disturbed and quiet solar-geomagnetic conditions, applied to two satellites with different ballistic coefficients as detailed in Table 1. Sat-A and Sat-B represent typical SmallSats of mass,  $m_s$ , and projected area,  $A_s$ . The selected “real-world” intervals were chosen based of a review of the environmental parameters that describe solar-geomagnetic activity including the solar wind speed ( $V_{sw}$ ) and proton density (PD), the disturbance storm time (Dst) index (Mayaud, 1980), the IMF  $B_y$  and  $B_z$  components, and the auroral electrojet (AE) index (Davis and Sugiura, 1966). The interplanetary parameters ( $V_{sw}$ , PD, IMF  $B_y$  and  $B_z$ ) and the geomagnetic responses in Dst and AE are reflective of the processes by which energy is transferred from the solar wind to the MIT system (Nwankwo (2016); and references therein). Model runs of the atmospheric density profile were made for the quiet environmental interval of July 2006 and for the disturbed conditions of July 2000. **Within the disturbed interval we also estimat the enhanced atmospheric drag effect on the satellites’ parameters caused by the Bastille Day event during 14-16 July 2000. In particular, we simulate the mean altitude decay ( $h$ ) and the orbit decay rates (ODRs) of the satellites during the regimes.**

**Table 1.** Orbital and ballistic parameters used in this study


Satellite	Altitude (km)	$m_s$ (kg)	$A_s$ (m <sup>2</sup> )	$C_d$	$B$ (m <sup>2</sup> /kg)
Sat-A	450	250	0.25	2.2	$2.200 \times 10^{-3}$
Sat-B	450	522	0.72	2.2	$3.034 \times 10^{-3}$

## 2.1 Quiet environmental conditions near solar minimum (1-31 July 2006)

Solar cycle 23 was on its descending phase in 2006 heading towards a solar minimum, which occurred in December 2008. Solar minimum is usually accompanied by a reduction in both solar radiant emissions and the frequency of solar transient events. The monthly averages of F10.7 and Ap for July 2006 were 78.4 sfu and 6.5 nT, respectively. Figure 3 is a plot of the 1-hour averaged variations in  $V_{sw}$ , PD, Dst, IMF  $B_y$  and  $B_z$  and AE for July 2006. The most notable feature, or lack thereof, was the essentially flat Dst index throughout the month that is indicative of no significant geomagnetic storms. However there were a number of interesting features related to the state of the solar-terrestrial environment. In particular, we note the character of the background solar wind speed,  $V_{sw}$ , and density, PD, on 04 July which is indicative of a pressure buildup on the nose of the magnetopause. These data are suggestive of a CIR that was not particularly well coupled to the magnetosphere due to a non-favourable IMF  $B_z$  (Pokhotelov et al., 2009). Supporting evidence of a CIR was the simultaneous detection of an increased flux of energetic protons (data not included here) observed just inside the magnetopause by the Geostationary Operational Environmental Satellite (GOES)(Posner et al., 1999). Similar “CIR-like” features in Figure 3 were the interplanetary parameters for 28 July and 31 July although their strengths were apparently less intense than the feature observed earlier in the month. We suspect that the sources for these CIRs were high-stream flows originating from within solar Coronal Holes (CHs). A review of solar imagery (data not included here) available from the Extreme ultraviolet Imaging Telescope (EIT) on the Solar and Heliospheric Observatory (SOHO) satellite revealed that there was a distinctly visible CH that crossed the solar disk in early July. This CH may well have been the source of a high-stream flow that resulted in the detected CIR on 04 July. During the middle of the month no CHs were apparent excepting a stationary, non-geoeffective, polar CH in the northern solar hemisphere. However, later in the month several CHs could be seen on the solar disk which might have been the sources for the solar wind features detected on 28 July and 31 July. The final characteristic of interest in Figure 3 was the periodic geomagnetic activity observed in the AE substorm index which was suggestive of High-Intensity, Long-Duration, Continuous AE Activity (HILDCAA). To this end, Guarnieri et al. (2006) noted that HILDCAA events can often be associated with CIRs, particularly on the downside of the solar cycle, as was the case here. The atmospheric drag effects modeled for this solar minimum interval of relative quiet will be compared and contrasted to the disturbed period of July 2000.


## 2.2 Disturbed environmental conditions near solar maximum (1-31 July 2000)

Year 2000 witnessed the expected rise in overall solar activity as the sun was progressing towards the maximum of cycle 23 which peaked in November 2001. Figure 4 is a plot of the hourly-averaged interplanetary and geomagnetic parameters for July 2000. The related monthly averages of the F10.7 and Ap indices were 212.2 sfu and 21.4 nT, respectively. Germane to this interval were the solar wind drivers and, more importantly, their significant fluctuations and increases, in the PD on days 01-04 July, 09-15 and 25-29 July. These fluctuations had significant consequences on the MIT system. Of note was the occurrence on 14-15 July of an intense geomagnetic storm (Gonzalez et al., 1994) having a Dst of -301 nT. The apparent source of this, so-called Bastille Day event which nominally spanned the 3-day interval 14-to-16 July, was a geoeffective CME that was first observed erupting from the sun at 10:54 UT on 14 July in association with an X5.7 flare within active region



**Figure 3.** 1-hour averaged variations in  $V_{sw}$ , PD, Dst, IMF  $B_y$  and  $B_z$ , and AE for the geomagnetically quiet, solar minimum interval 1-31 July 2006.

#19077 at solar location N22W07 (Denig et al., 2018). On 15 July a large Sudden Storm Commencement (SSC) of 112 nT at 14:37 UT marked the arrival of the CME at the magnetopause and the start of the geomagnetic storm main phase (Closs, 1967). Given a total transit time of just under 28 hours the estimated CME speed from the sun to earth was a fast  $\sim 1500$  km/s which is consistent with an initially observed speed of 1673 km/s (Denig et al., 2018) and assessments of the probability of a significant geomagnetic response (Srivastava and Venkatakrishnan, 2002). The 40 nT magnitude and initial negative polarity of the IMF  $B_z$  suggests that significant energy was transferred to the MIT system from the solar wind. The related increase in AE corresponding to a substorm occurring within the geomagnetic storm (McPherron et al., 1973; Kepko et al., 2015) indicates enhanced ionospheric currents within the auroral zone due to the strong coupling between the IMF and the MIT (Pudovkin et al., 1995). Clearly, the upper atmosphere was significantly disturbed throughout July 2000 and, in particular, during to the Bastille Day event. The expected consequences of the enhanced solar-geomagnetic activity for July 2000 was increased atmospheric drag and a consequential decrease in the satellite orbital altitude.



**Figure 4.** 1-hour averaged variations in  $V_{sw}$ , PD, Dst, IMF  $B_y$  and  $B_z$ , and AE for the geomagnetically quiet, solar minimum interval 1-31 July 2000.

### 2.3 Modeling atmospheric drag effect on LEO satellites' trajectory

We previously formulated and solved a set of coupled differential equations to obtain the instantaneous position, velocity and acceleration of a typical LEO satellite under the influence of atmospheric drag ( e.g., (Nwankwo and Chakrabarti, 2014, 2015; Nwankwo et al., 2015; Nwankwo, 2018)). We used a spherical coordinate system  $(r, \theta, \phi)$  with origin at the center of the earth and assumed that the polar angle is constant; that is,  $\theta = constant$ . In satellite parlance a constant polar angle is equivalent, in principle, to a constant satellite inclination angle. Orbital decay was determined as a consequence of changes in the radial distance,  $r$ , and the azimuthal angle,  $\phi$ , through the following set of coupled equations:

$$\dot{v}_r = -\dot{\phi}^2 \frac{A_s C_d}{m_s}, \quad (1)$$

$$200 \quad \dot{r} = v_r, \quad (2)$$

$$\ddot{\phi} = -\frac{1}{2}r\rho\dot{\phi}^2\frac{A_s C_d}{m_s}, \quad (3)$$

$$\dot{\phi} = \frac{v_\phi}{r}, \quad (4)$$


where  $v_r$  and  $v_\phi$  are, respectively, the radial and tangential velocity components.  $\dot{\phi}$  and  $\ddot{\phi}$  are, respectively, the azimuthal angular velocity and azimuthal angular acceleration. The parameters  $C_d$ ,  $A_s$ , and  $m_s$  were defined in Section 1 - recall that the expression  $C_d A_s / m_s = B$  is the ballistic coefficient. In the current analysis, the radial velocity,  $v_r$ , is used to calculate the daily ODR whereas the radial distance,  $r$ , is used to model changes in satellite altitude.

## 2.4 Atmospheric density model

The effects of atmospheric drag on LEO satellites and hence the rates at which satellite orbits decay largely depend on the atmospheric density which, in turn, is heavily influenced by solar and geomagnetic activity (Fujiwara et al., 2009). Accurate knowledge of atmospheric drag requires a high-fidelity model of the in-situ neutral-gas density or, more generally, the atmospheric density profile. Supporting information regarding the level of atmospheric heating and, in turn, atmospheric expansion can be gleaned from knowledge of the atmospheric temperature profile. The upper atmosphere, or thermosphere, exhibits large solar-cycle variations in temperature, density, composition and winds (Walterscheid, 1989). A number of high-quality models are available that provide suitable approximations of atmospheric profiles of density,  $\rho$ , and temperature,  $T$  (Picone et al., 2002; Bruinsma et al., 2003; Bowman et al., 2008). For this work we have selected the Naval Research Laboratory Mass Spectrometry and Incoherent Scatter Extended 2000 (NRLMSISE-00) empirical atmospheric model. NRLMSISE-00 consists of parametric and analytic approximations to physical theory for the vertical structure of the atmosphere as a function of time, location, solar and geomagnetic activity. While a global specification was used to extract the density along the satellite flight path, the atmospheric curves used in Figures 5, 7 and 11 (to be discussed) to represent a general atmospheric response used a reference altitude of 450 km. The solar-geomagnetic parameters used for NRLMSISE-00 model are the daily values of F10.7 and  $A_p$  (Nwankwo and Chakrabarti (2018); and references therein).

## 3 Results and discussion

The results of our simulation arising as solutions to the above set of coupled differential equations are presented in this section. The environmentally quiet interval of July 2006 is presented as the baseline for atmospheric drag whereas the disturbed interval of July 2000 illustrates the deleterious impact that solar-geomagnetic activity can have on satellite orbits. Within each of these intervals a 12-day period of environmentally quiet and exceptionally disturbed activity, respectively, is used to highlight the impact of extreme conditions.




**Figure 5.** Mean daily variations in Dst, Bz,  $\rho$ , T, h and ODR for Sat-A (blue) and Sat-B (red) for quiet conditions (July 2006).

### 3.1 Atmospheric drag effects for quiet solar-geomagnetic activity (July 2006)

Figure 5 depicts the mean daily variations in Dst, Bz, atmospheric density ( $\rho$ ) and temperature (T), and the altitude (h) and orbit decay rates (ODRs) of Sat-A and Sat-B under the quiet solar-geomagnetic conditions of July 2006. During this 1-month interval of relatively low environmental stress, the ranges of Dst and Bz (daily mean) are -28.71–7.75 (nT) and -1.18–1.84 (nT), respectively. T varied between 770 °K and 880 °K, whereas  $\rho$  varied between  $0.33 \times 10^{-12} \text{ kg/m}^3$  and  $0.55 \times 10^{-12} \text{ kg/m}^3$ . These atmospheric parameters are consistent with low geomagnetic activity, solar minimum conditions (Fujiwara et al., 2009). The orbital drag calculations indicate that Sat-A decayed by 0.52 km during the month with an ODR ranging from 13-to-23 m/day whereas Sat-B decayed by 0.65 km with an ODR range of 16-to-29 m/day. These modest yet consistent differences in ODR and decay for Sat-A and Sat-B reflect the differences in their respective ballistic coefficients (see Table 1). Figure 6 is a plot of the daily F10.7, SSN and Ap indices. While no significant geomagnetic storms occurred during the entire month we note that the minor increases in the daily Ap for 05 July, 28 July and possibly 31 July corresponded to slight increases in the atmospheric parameters and the peak ODRs of 23 m/day (29 m/day) for Sat-A (Sat-B). The baseline ODRs for July 2006 will be contrasted with the model decay rates for the solar maximum, geomagnetically disturbed interval of July 2000.

### 3.2 Atmospheric drag effects for enhanced solar-geomagnetic activity (July 2000)


Figure 7 depicts the mean daily variations in Dst, Bz,  $\rho$ , T, h, ODR for the environmentally enhanced interval of July 2000. During this month the range of daily values of Dst and Bz are -171.63 – 12.75 nT and -4.84 – 13.30 nT, respectively. The



**Figure 6.** Daily values of Ap, International SSN and F10.7 for July 2006.

modeled temperatures for the month varied between 1156 °K and 1580 °K which were indicative of a generally warm at-  
 245 mosphere near solar maximum (Fujiwara et al., 2009). In response, the thermospheric densities for an expanded atmosphere  
 ranged from  $2.4 \times 10^{-12}$  kg/m<sup>3</sup> to  $5.7 \times 10^{-12}$  kg/m<sup>3</sup> (Fujiwara et al., 2009; Emmert, 2015). Accordingly, Sat-A decayed by  
 about 2.77 km whereas Sat-B decayed by about 3.09 km. Again, as expected, the integrated effect of atmospheric drag on Sat-B  
 was greater than that experienced by Sat-A due to differences in the ballistic coefficients. The range of daily values for Sat-A's  
 ODR was 65 m/day to 120 m/day whereas the range of ODR for Sat-B was 78 m/day to 142 m/day. An interesting trend is  
 250 the general increase then decrease in ODR throughout the month which is consistent with the variations in the modeled  $\rho$  and T.

In order to further investigate this trend we plotted in figure 8 the daily indices for F10.7, SSN and Ap and found that the  
 density and temperature trends were consistent with the solar F10.7 and SSN indices. Again, this is not surprising in that F10.7  
 is a key input to the NRLMSISE-00 model. The apparent similarity in the behaviors of the daily SSN and daily F10.7 index was  
 255 also reasonable given that intense radiant emissions from solar faculae are proportional to the number of sunspots (Ambelu  
 et al., 2011). However, we note that the significant spike in F10.7 (due to the intense flare) did not reflect in the simulated  
 ODR. This outcome is consistent with the findings of Knowles et al. (2001) who stated that "the effect of geomagnetic activity  
 appears to be more important than that of prompt radiation." The model values for ODC, as well as the atmospheric density  
 and temperature, spiked predictably mid-month in response to the additional energy input from the Bastille Day geomagnetic  
 260 storm. The thermospheric T and  $\rho$  on the peak storm day (16 July) were 1580 °K and  $5.7 \times 10^{-12}$  kg/m<sup>3</sup>, and the corresponding  
 values of ODRs for Sat-A and Sat-B were 120 m/day and 142 m/day, respectively. A more detailed plot of the 3-hour magnetic  
 $a_p$  index included in figure 9 for 13-17 July indicates that the geomagnetic storm lasted about one day, starting near noon on



**Figure 7.** Mean daily variations in Dst, Bz,  $\rho$ , T, h and ODR for Sat-A (blue) and Sat-B (red) for quiet conditions (July 2000).


15 July and ending some 24 hours later. We note that the start of the storm was consistent with the previously mentioned SSC that occurred at 14:37 UT on 15 July. Also occurring during the month of July 2000 were a series of minor disturbances (e.g., 265 days 11, 20, 23, 26 and 28-29 as revealed by Fig. 8) which contributed to the enhanced temperatures and densities (beyond solar inputs alone) observed throughout the month.

The results of our simulation showed that the maximum ODRs for Sat-A and Sat-B were in response to the Bastille Day event. We glean from these results that the background atmospheric parameters were responsive to both the slowly-varying solar irradiance inputs during the month and to the impulsive geomagnetic storm inputs. These results confirm that the transient 270 response of satellite drag to impulsive geomagnetic storms lead to the largest uncertainties in orbit dynamics and pose a great risk to efficient satellite operations.


### 3.3 Intervals of exceptionally quiet and disturbed environmental stress

In this section we focus on the 12-day sub-intervals of elevated solar and geomagnetic activity for 09-20 July 2000 and of exceptionally quiet activity for 15-26 July 2006. For these intervals we compare and contrast the activity levels on the satellite 275 trajectory parameters (h, ODR) with the corresponding perturbations in T and  $\rho$ . Figure 10 depicts 1-hour averaged variations






**Figure 8.** Daily values of  $A_p$ , International SSN and F10.7 for July 2000.




**Figure 9.** Three-hour  $A_p$  values for 13-17 July 2000.



**Figure 10.** 1-hour averaged values of  $V_{sw}$ , PD, Dst, IMF  $B_y$  and  $B_z$ , and AE during 9-20 July 2000 (left) and 15-26 July 2006 (right) (Nwankwo et al., 2020)

in  $V_{sw}$ , PD, Dst, IMF  $B_y$  and  $B_z$ , and AE for the intervals of high activity (left) and low activity (right). In Fig 11 we show the corresponding daily variations in Dst,  $B_z$ ,  $\rho$ , T, h and ODR for Sat-A with the lower ballistic coefficient (blue trace) and Sat-B with the higher ballistic coefficient (red trace) within the intervals of elevated activity (left) and quiescent activity (right). During the sub-interval of elevated activity (left) Sat-A orbit decayed by 1.14 km and the ODR increased from 81.46 m/day on 09 July to 120 m/day on 16 July which is just after the peak of the Bastille Day storm. Similarly, Sat-B decayed by about 1.27 km, and the ODR increased from 91.85 m/day on 09 July to 142 m/day on 16 July. Considering the sub-interval of low activity (right) Sat-A (blue trace) decayed by a total of 0.16 km with an ODR ranging from 13.41 to 17.17 m/day whereas Sat-B (red trace) decayed by a total of 0.20 km with an ODR ranging from 16.67 to 21.25 m/day. The salient features for the orbital decay and maximum ODR for both satellites for the active versus quiet conditions are summarized in Table 2. The stark contrast between the two regimes indicates that solar-geomagnetic activity had a more than 6-fold (7-fold in this case) impact on the orbital parameters for the modeled conditions. This dramatic effect makes it imperative that the orbital parameters for relevant space objects in LEO be frequently updated (Knowles et al., 2001).




**Figure 11.** Corresponding mean daily variations in Dst, Bz,  $\rho$ , T, h and ODR for Sat-A and Sat-B during 09-20 July 2000 (left) and 15-26 July 2006 (right)

**Table 2.** Summary of altitude decay and ODR of Sat-A and Sat-B for the extreme (12-day) activity levels

Date Range	Activity Level	Satellite	B ( $\text{m}^2/\text{kg}$ )	Decay (kg)	Max ODR (m/day)
09-20 July 2000	High	Sat-A	$2.200 \times 10^{-3}$	1.14	120.0
09-20 July 2000	High	Sat-B	$3.034 \times 10^{-3}$	1.27	120.0
15-26 July 2000	Low	Sat-A	$2.200 \times 10^{-3}$	0.16	17.2
15-26 July 2000	Low	Sat-B	$3.034 \times 10^{-3}$	0.20	21.2

In Figure 12, we show detailed analysis (and comparison) of how h and ODR of the two satellites varied during the regimes of elevated and exceptionally quiet solar and geomagnetic activity. The goal of this analysis is to demonstrate how enhanced atmospheric drag caused by the July 2000 Bastille Day event affected LEO satellites in contrast to the interval of exceptionally quiet geomagnetically activity conditions during 15-26 July 2006. We describe new indices in Tables 3 and 4 for the analysis. Associating the Tables with the geometry of curves in figure 12 gives a better understanding of the analysis to follow. When compared with the 12-day interval of exceptionally quiet geomagnetic activity, the total decay (h) and ODR increase (from the



**Figure 12.** Contrast between the daily variations in  $h$  and ODR of Sat-A (left) and Sat-B (right) during 09-20 July 2000 and 15-26 July 2006

background or initial values) during the elevated geomagnetic activity are 0.98 km and 80.24 m/day for Sat-A (described by  $dh_{Sat-A}$  and  $\Delta ODR_{Sat-A}$ , respectively), and 1.07 km and 84.12 m/day for Sat-B (described by  $dh_{Sat-B}$  and  $\Delta ODR_{Sat-B}$ ). The decay caused by the Bastille Day event are 0.34 km and 0.39 km for Sat-A and Sat-B, respectively (described by  $h_{BD-A}$  and  $h_{BD-B}$ ). The respective ODR increments (due to Bastille Day event) are 40.23 m/day and 58.00 m/day (described by  $\Delta ODR_{BD-A}$  and  $\Delta ODR_{BD-B}$ ). The corresponding percentage increase of the two parameters ( $h$  and ODR) when contrasted with the interval of exceptionally quiet geomagnetic activity are 74.24% and 66.74% for Sat-A, and 75.35% and 59.18% for Sat-B. However, the additional percentage increase of Bastille Day event to the monthly mean values of July 2000 are 34.69% ( $h$ ) and 50.13% (ODR) for Sat-A, and 36.45% and 68.95% for Sat-B. This result implies that storms of this magnitude can add more than 30% and 50-70% increase to background  $h$  and ODR during the interval.

#### 4 Conclusions

Solar activity in the form of increased solar irradiance and flux of energetic particles form important channels through which the earth's atmosphere is impacted. Atmospheric heating and expansion can significantly increase orbital drag which, in turn,

**Table 3.** Indices used for description effects of the 12-day intervals of elevated and exceptionally quiet geomagnetic activity (09-20 July 2000 and 15-26 July 2006). EQGA connotes exceptionally quiet geomagnetic activity

s/n	Abbrev/Symbol	Definition	Value
1	$h1_{A06}$	Sat-A's height on 15 July 2006	449.74 km
2	$h2_{A06}$	Sat-A's height on 26 July 2006	449.58 km
3	$h1_{A00}$	Sat-A's height on 9 July 2000	449.40 km
4	$h2_{A00}$	Sat-A's height on 20 July 2000	448.28 km
5	$h1_{B06}$	Sat-B's height on 15 July 2006	449.67 km
6	$h2_{B06}$	Sat-B's height on 26 July 2006	449.47 km
7	$h1_{A00}$	Sat-A's height on 9 July 2000	449.32 km
8	$h1_{B00}$	Sat-B's height on 20 July 2000	448.05 km
9	$ODR_{A06}$	orbit decay rate of Sat-A during EQGA corresponding to 15 July 2000	13.79 m/day
10	$ODR_{B06}$	orbit decay rate of Sat-B during EQGA corresponding to 15 July 2000	17.00 m/day
11	$ODR_{A00}$	mean orbit decay rate of Sat-A for July 2000	94.03 m/day
12	$ODR_{B00}$	mean orbit decay rate of Sat-B for July 2000	101.14 m/day
13	$ODR_{BD-A}$	orbit decay rate value of Sat-A on Bastille Day	120.47 m/day
14	$ODR_{BD-B}$	orbit decay rate value of Sat-B on Bastille Day	142.12 m/day

**Table 4.** Indices used to analyse and contrast the effects of intervals of elevated and exceptionally quiet geomagnetic activity with those of the Bastille Day event.

s/n	Abbrev/Symbol	Definition	Value
1	$\Delta h1_{Sat-A}$	$h1_{A06} - h1_{A00}$	0.34 km
2	$\Delta h2_{Sat-A}$	$h2_{A06} - h2_{A00}$	1.32 km
3	$dh_{Sat-A}$	$\Delta h2_{Sat-A} - \Delta h1_{Sat-A}$	0.98 km
4	$\Delta h1_{Sat-B}$	$h1_{B06} - h1_{B00}$	0.36 km
5	$\Delta h2_{Sat-B}$	$h2_{B06} - h2_{B00}$	1.42 km
6	$dh_{Sat-B}$	$\Delta h2_{Sat-B} - \Delta h1_{Sat-B}$	1.07 km
7	$h_{BD-A}$	$h_{Sat-A}$ (13 July 2000) $- h_{Sat-A}$ (16 July 2000)	0.34 km
8	$h_{BD-B}$	$h_{Sat-B}$ (13 July 2000) $- h_{Sat-B}$ (16 July 2000)	0.39 km
9	$\Delta ODR_{Sat-A}$	$ODR_{A00} - ODR_{A06}$	80.24 m/day
10	$\Delta ODR_{BD-A}$	$ODR_{BD-A} - \Delta ODR_{Sat-A}$	40.23 m/day
11	$\Delta ODR_{Sat-B}$	$ODR_{B00} - ODR_{B06}$	84.12 m/day
12	$\Delta ODR_{BD-B}$	$ODR_{BD-B} - \Delta ODR_{Sat-B}$	58.00 m/day

perturbs satellite trajectories and results in accelerated orbital decay. In this work, we simulated the effect of atmospheric drag on two hypothetical SmallSats in LEO with different ballistic coefficients during 1-month long intervals of disturbed and quiet solar-geomagnetic activity. During a 1-month period of enhanced activity (01-31 July 2000) the increased density of the upper

310 atmosphere caused a modeled mean decay of 2.77 km (3.09 km) for the satellite with the smaller (larger) ballistic coefficient. Conversely, for the more quiescent period (01-31 July 2006) the mean decay was only 0.52 km (0.65 km) for the respective satellites. Further analysis and simulation of atmospheric drag for periods of elevated (or extreme) solar-geomagnetic activity during 09-15 July 2000 and exceptionally quiet geomagnetic activity (15-26 July 2006) resulted in Sat-A (Sat-B) modeled orbital decays of 1.14 km (1.27 km) and 0.16 km (0.20 km), respectively. We also estimated the enhanced atmospheric drag  
315 effect on the satellites's parameters caused by the July 2000 Bastille Day event in contrast to the interval of geomagnetically quiet conditions. While the percentage increase of  $h$  and ODR due to elevated geomagnetic activity (of 9-20 July 2000) are 74.24% and 66.74% (75.35% and 59.18%) for Sat-A (Sat-B), the additional (daily) percentage increase due to the Bastille Day event (14-15 July 2000) to the monthly mean values are 34.69% and 50.13% (36.45% and 68.95%) for Sat-A (Sat-B). The results of our simulation confirm the dependencies of the satellite drag on the ballistic coefficient and on the level of solar-  
320 geomagnetic activity. While such dependencies are generally intuitive our model is useful in quantifying these relationships and can thus contribute to an improved situational awareness as well as mitigating the potential threat posed by solar-geomagnetic activity in modulating satellite trajectories. **In addition (to this and other contributions outlined earlier in this paper) this work motivated the development of new method and indices for describing and estimating atmospheric drag effects on satellite ephemeris (when comparing between regimes of varying solar-geomagnetic activity).**

325 *Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* We acknowledge use of NASA/GSFC's Space Physics Data Facility's OMNIWeb service for the 1-hour averaged values of  $V_{sw}$ , PD, Dst, IMF  $B_y$  and  $B_z$ , and AE. The yearly-averaged SSN (version 1.0) was obtained from the World Data Center for the Sunspot Index and Long-term Solar Observations (SILSO), Royal Observatory of Belgium, Brussels. The daily and monthly-averaged values for SSN, F10.7 and  $A_p$  were provided by the GFZ German Research Centre for Geosciences and Space Weather Canada. Other data sources include  
330 the Observatori de l'Ebre (Spain) and the SOHO LASCO CME Catalog. **Authors also acknowledge Mr. Felix Oke and the Language Laboratory, Department of Languages and Linguistics, Anchor University, Lagos for providing us with temporal space to complete this manuscript at a critical moment when our facility suffered power fault.**

## References

- Ambelu, T., Falayi, E.O., Elemo, E.O. and Oladosu, O.: Estimation of total solar irradiance from sunspot number. *Lat. Am. J. Phys. Educ.*, 335 5, 4, pp. 741-745, Available at: [http://www.lajpe.org/dec11/LAJPE\\_576\\_Ambelu\\_Tebabal\\_prreprint\\_corr.pdf](http://www.lajpe.org/dec11/LAJPE_576_Ambelu_Tebabal_prreprint_corr.pdf), 2011. Last accessed: 25 March 2020.
- Bhatnagar, V.P., Germany, G.A. and Tan, A.: Satellite ballistic coefficients and the lower thermosphere. *Geophys. Res. Lett.*, 32, L03109, doi: 10.1029/2004GL021627, 2005.
- Bobrinsky, N and Del Monte, L.: The Space Situational Awareness Program of the European Space Agency, *Cosmic Research*, 48, 5, pp 340 392-398, doi: 10.1134/S0010952510050035, 2010.
- Boteler, D. H.: A 21st century view of the March 1989 magnetic storm. *Space Weather*, 17, 1427-1441, <https://doi.org/10.1029/2019SW002278>, 2019.
- Borovsky, J.E. and Denton, M.H.: Differences between CME-driven storms and CIR-driven storms. *J. Geophys. Res.*, 111, A07S08, doi: 10.1029/2005JA011447, 2006.
- 345 Bowman, B.R.: True Satellite Ballistic Coefficient Determination for HASDM. Paper AIAA 2002-4887, AIAA/AAS Astrodynamics Specialist Conference and Exhibit 5-8 August 2002, Monterey, CA, USA, Available at [http://sol.spacenvironment.net/JB2008/pubs/JB2006\\_AIAA\\_2002\\_4887.pdf](http://sol.spacenvironment.net/JB2008/pubs/JB2006_AIAA_2002_4887.pdf), 2002. Last accessed: 11 Mar 2020.
- Bowman, B., Tobiska, W.K., Marcos, F.A., Huang, C.Y. Lin, C.S. and Burke, W.J.: New Empirical Thermospheric Density Model JB2008 Using New Solar and Geomagnetic Indices. paper 6438 (?), AIAA/AAS Astrodynamics Specialist Conference, 18-21 August 2008, 350 Honolulu, Hawaii, Available: [http://sol.spacenvironment.net/JB2008/pubs/AIAA\\_2008-6438\\_JB2008\\_Model.pdf](http://sol.spacenvironment.net/JB2008/pubs/AIAA_2008-6438_JB2008_Model.pdf), 2008. Last accessed: 05 Mar 2020.
- Bruinsma, S., Thuillier, G. and Barlier, F.: The DTM-2000 empirical thermosphere model with new data assimilation and constraints at lower boundary: accuracy and properties. *J. Atmos. Terr. Phys.*, 65, 9, pp. 1053-1070, doi: [https://doi.org/10.1016/S1364-6826\(03\)00137-8](https://doi.org/10.1016/S1364-6826(03)00137-8), 2003.
- 355 Chobotov V.A.: *Orbital Mechanics*. 3rd ed. AIAA Education series. American Institute of Aeronautics Inc., Virginia, 193-213. 2002.
- Clette, F., Svalgaard, L., Vaquero, J.M. and Cliver, E.W.: Revisiting the Sunspot Number: A 400-Year Perspective on the Solar Cycle. *Space Sci. Rev.*, 186, 35-103, doi: 10.1007/s11214-014-0074-2, 2014.
- Closs, R.L.: A possible explanation of the sudden commencement and initial phase of a magnetic storm in terms of induced ionospheric currents. *J. Geophys. Res.*, 72, 15, 3987-3993, doi: 10.1029/JZ072i015p03987, 1967.
- 360 Cooke, G.E.: Satellite drag coefficients. *Planet. Space Sci.*, 13, 10, pp 929-946, doi: [https://doi.org/10.1016/0032-0633\(65\)90150-9](https://doi.org/10.1016/0032-0633(65)90150-9), 1965.
- Davis, T.N. and Sugiura, M.: Auroral electrojet activity index AE and its universal time variations. *J. Geophys. Res.*, 71(3), 785-801, doi:10.1029/JZ071i003p00785, 1966.
- Denig, W.F., Wilkinson, D.C. and Redmon, R.J.: Extreme Space Weather Events: A GOES Perspective. in "Extreme Events in Geospace - Origins, Predictability and Consequences". ed: N. Buzulukova, 283-347, doi:<https://doi.org/10.1016/B978-0-12-812700-1.00012-1>, 2018.
- 365 Eddy, J.: *The Sun, the Earth, and Near-Earth Space: A Guide to the Sun-Earth System*. 316, U.S. Government Printing Office, ISBN 978-0-16-08308-8, Available: [https://lws-trt.gsfc.nasa.gov/images/pdf/john\\_eddy/SESBook\\_Interactive.pdf](https://lws-trt.gsfc.nasa.gov/images/pdf/john_eddy/SESBook_Interactive.pdf), 2009. Last accessed: 16 Mar 2020.
- Emmert, J.T.: Thermospheric mass density: A review. *Adv. Space Res.*, 56, 5, 773-824, doi:<https://doi.org/10.1016/j.asr.2015.05.038>, 2015.



- Fujiwara, H., Miyoshi, Y., Jin, H., Shinagawa, H., Otsuka, Y., Saito, A. and Ishii, M.: Thermospheric temperature and density variations, Proc. IAU, 5, S264, 310-319, doi:<https://doi.org/10.1017/S1743921309992857>, 2009.
- 370 Gonzalez, W.D., Joselyn, J.A., Kamide, Y., Kroehl, H.W., Rostoker, G., Tsurutani, B.T., and Vasyliunas, V.M.: What is a geomagnetic storm?. J. Geophys. Res., 99( A4), 5771-5792, doi:10.1029/93JA02867, 1994.
- Gosling, J. and Pizzo, V.: Formation and Evolution of Co-rotating Interaction Regions and their Three Dimensional Structure. Space Science Reviews 89, 21-52, doi: <https://doi.org/10.1023/A:1005291711900>, 1999.
- 375 Gosling, J.T., Bame, S.J., McComas, D.J. and Phillips, J.L.: Coronal mass ejections and large geomagnetic storms. Geophys. Res. Lett., 17(7), 901-904, doi:<https://doi.org/10.1029/GL017i007p00901>, 1990.
- Guarnieri, F.L., Tsurutani, B.T., Gonzalez, W.D., Echer, E., Gonzalez, A.L.C.: ICME and CIR storms with particular emphasis on HILDCAA events, ILWS Workshop 2006, Goa, India, 19-20 February 2006. Available: [https://cdaw.gsfc.nasa.gov/publications/ilws\\_goa2006/266Guarnieri.pdf](https://cdaw.gsfc.nasa.gov/publications/ilws_goa2006/266Guarnieri.pdf), 2006. Last accessed: 20 Mar 2020.
- 380 Jakhu, R.S.: Iridium-Cosmos collision and its implications for space operations. In Kai-Uwe Schrogl, et al (eds.), Yearbook on Space Policy 2008/2009, Springer, Wien New York, 2010, 254-275, 2009.
- Kelso, T.S.: Analysis of the Iridium 33Cosmos 2251 Collision. AAS 09-368, 14, Available: <https://celestrak.com/publications/AAS/09-368/AAS-09-368.pdf>, 2009. Last accessed: 16 Mar 2020.
- Kepko, L., McPherron, R.L., Amm, O., Apatenkov, S., Baumjohann, W., Birn, J., Lester, M., Nakamura, R., Pulkkinen, T.I. and Sergeev, V.: 385 Substorm Current Wedge Revisited. Space Sci. Rev., 190, 1-46 doi:<https://doi.org/10.1007/s11214-014-0124-9>, 2015.
- King-Hele, D.: Satellite Orbits in an Atmosphere: Theory and Applications. 291, Blackie Acad. and Prof., New York, 1987.
- Knipp, D.: Understanding Space Weather and the Physics Behind It. Space Technology Series, ISBN: 978-0-07-340890-3, 727, 2011.
- Knowles, S., Picone, J., Thonnard, S. and Nicholas, A.C.: The Effect of Atmospheric Drag on Satellite Orbits During the Bastille Day Event. Sol. Phys., 204, 387-397, <https://doi.org/10.1023/A:1014223807360>, 2001.
- 390 Mayaud, P.N.: Derivation, Meaning, and Use of Geomagnetic Indices. Geophysical Monograph 22, American Geophysical Union, Washington, DC, 1980.
- McCrea I. (Science and Technology Facilities council, RAL Space, Wiltshire, UK). Letter to: Newton International Fellowship (The Royal Society, UK). 2018 Mar. 21. 1 Leaf. Located at: 6-9 Carlton House, Terrace, London, 2018.
- McPherron, R. L., Russell, C.T. and Aubry, M.P.: Satellite studies of magnetospheric substorms on August 15, 1968: 9. Phenomenological 395 model for substorms. J. Geophys. Res., 78, 16, 3131– 3149, doi: 10.1029/JA078i016p03131, 1973.
- Mitra, A.P.: Ionospheric Effects of Solar Flares. Astrophys. Space Sci. Library (ASSL), Vol 46, ISBN 978-94-010-2231-6, XX, doi: <https://doi.org/10.1007/978-94-010-2231-6>, 1974.
- Nwankwo V.U.J.: Effects of Space Weather on Earth’s Ionosphere and Nominal LEO Satellites’ Aerodynamic Drag. (PhD Thesis), University of Calcutta, Kolkata, India, 2016.
- 400 Nwankwo, V.U.J.: Space Weather: Responses of the Atmosphere to Solar Activity and Its Implications for LEO Satellites Aerodynamic Drag. in “Exploring the Universe: From Near Space to Extra-Galactic”. eds: B. Mukhopadhyay and S. Sasmal, Springer Inter. Pub., Switzerland AG, 2018.
- Nwankwo, V.U.J. and Chakrabarti, K.S.: Effects of plasma drag on low earth orbiting satellites due to heating of earth’s atmosphere by coronal mass ejections. arXiv:1305.0233c1 [physics.space-hp]., 2013.
- 405 Nwankwo, V.U.J and Chakrabarti, K.S.: Theoretical modeling of drag force impact on a model international space station (ISS) during variation of solar activity. Trans. JSASS Aerosp. Technol. Jpn. 12, 47-53, 2014.

- Nwankwo, V.U.J. and Chakrabarti, K.S.: Analysis of planetary and solar-induced perturbations on trans-Martian trajectory of Mars missions before and after Mars orbit insertion. *Indian J. Phys.* 89, 1235-1245, doi:10.1007/s12648-015-0705-9, 2015.
- 410 Nwankwo, V.U.J. and Chakrabarti, S.K.: Effects of space weather on the ionosphere and LEO satellites' orbital trajectory in equatorial, low and middle latitude. *Adv. Space Res.*, 61(7), 1880-1889, doi:https://doi.org/10.1016/j.asr.2017.12.034, 2018.
- Nwankwo, V.U.J., Chakrabarti, S.K. and Weigel, R.S.: Effects of plasma drag on low earth orbiting satellites due to solar forcing induced perturbations and heating. *Adv. Space Res.* 56, 47-56. doi:https://doi.org/10.1016/j.asr.2015.03.044, 2015.
- Nwankwo, V.U.J., Jibiri, N.N. and Kio, M.T.: The impact of space radiation environment on satellites operation in near-Earth space. in "Satellites and Innovative Technology", ed: J. Becedas, ISBN 987-1-78985-862-4, InTechOpen Publishing London, 2019.
- 415 Nwankwo V.U.J., Denig W., Ajakaiye M.P. et al., 2020a. Simulation of atmospheric drag effect on low Earth orbit satellites during intervals of perturbed and quiet geomagnetic conditions in the magnetosphere-ionosphere system. 2020 IEEE-ICMCECS, Lagos, Nigeria , pp 1-7, DOI:10.1109/ICMCECS47690.2020.247003.
- Parker, E.N.: Dynamics of the Interplanetary Gas and Magnetic Fields. *Astrophys J*, 128, 664-676, doi:10.1086/146579, 1958.
- Phillips, K.J.H.: Solar flares: A review. *Vistas Astronomy*, 34 Part 2, 353-365, doi:https://doi.org/10.1016/0083-6656(91)90014-J, 1991.
- 420 Picone, J.M., Hedin, A.E., Drobm, D.P. and Aikin, A.C.: NRLMSISE-00 empirical model of the atmosphere: Statistical comparisons and scientific issues. *J. Geophys. Res.*, 107, A12, 468-1484, doi:10.1029/2002JA009430, 2002.
- Pokhotelov, D., Mitchell, C.N., Jayachandran, P.T. MacDougall, J.W. and Denton, M.H.: Ionospheric response to the corotating interaction region-driven geomagnetic storm of October 2002. *J. Geophys. Res.*, 114, A12311, doi:10.1029/2009JA014216, 2009.
- Posner, A., Bothmer, V., Thompson, B.J., Kunow, H., Heber, B., Müller-Mellin, R., Lazarus, A.J., Szabo, A., Mikić, Z., and Linker, J.A.: 425 In-ecliptic CIR-associated energetic particle events and polar coronal hole structures: SOHO/COSTEP observations for the Whole Sun Month Campaign. *J. Geophys. Res.*, 104(A5), 9881-9890, doi:10.1029/98JA02654, 1999.
- Pross, G.W.: Physics of the Earth's space environment. Springer Berlin Heidelberg, Germany, 2004.
- Pudovkin, M.I., Semenov, V.S., Kotikov, A.L. and Shishkina, E.M.: Dynamics of auroral electrojets and energetics of substorms. *J. Atmos. Terr. Phys.*, 57, 2, 187-192, doi: https://doi.org/10.1016/0021-9169(93)E0033-6, 1995.
- 430 Raeder J., Wang Y.L., Fuller-Rowell T.J. and Singer H. J.: Global simulation of magnetospheric space weather effects of the Bastille day storm. *Solar Physics* 204, 325-338, 2001.
- Rose, D.C. and Ziauddin, S.: The polar cap absorption effect. *Space Sci. Rev.*, 1, 115-134, doi:10.1007/BF00174638, 1962.
- Rostoker, G.: Geomagnetic indices. *Rev. Geophys.*, 10, 4, 935-950, doi:10.1029/RG010i004p00935, 1972.
- Ryan, J.M., Lockwood, J.A. and Debrunner, H.: Solar Energetic Particles. *Space Sci. Rev.*, 93, 35-53, 435 https://doi.org/10.1023/A:1026580008909, 2000.
- Sauer, H.H. and Wilkinson, D.C.: Global mapping of ionospheric HF/VHF radio wave absorption due to solar energetic protons. *Space Weather*, 6, S12002, doi: 10.1029/2008SW000399, 2008.
- Schatten, K.: Large Scale Properties of the Interplanetary Magnetic Field. N71-20467, NASA Goddard Space Flight Center, Available: https://ntrs.nasa.gov/archive/nasa/casi.ntrs.nasa.gov/19710010992.pdf, 1971. Last accessed: 16 Mar 2020.
- 440 Shiroma W.A., Martin, L.K., Akagi, J.M., Akagi, J.T., Byron, L.W., Fewell, B.A. and Ohta, A.T.: CubSats: A bright future for nanosatellites. *Central European Journal of Engineering*, 1, 9-15, doi:https://doi.org/10.2478/s13531-011-0007-8, 2011.
- Song, P., Singer, H.J., and Siscoe, G.L.: *Space Weather*. AGU Geophysical Monograph Series, Number 125, Vol 96, 440 pp., ISBN13: 9780875909844, 2001.

- 445 Srivastava, N. and Venkatakrishnan, P.: Relationship between CME Speed and Geomagnetic Storm Intensity. *Geophys. Res. Lett.*, 29(9), doi: 10.1029/2001GL013597, 2002.
- Tapping, K.F.: The 10.7 cm solar radio flux (F10.7). *Space Weather*, 11, 394-406, doi:10.1002/swe.20064, 2013.
- Walterscheid, R.L.: Solar cycle effects on the upper atmosphere - Implications for satellite drag. *J. Spacecraft Rockets, Proc. AIAA*, 26, 439-444, Aerospace Engineering Conference and Show, Los Angeles, CA, 14-16 Feb. 14-16 1989, doi:10.2514/3.26089, 1989.
- 450 Watari S., Kunitake M. and Watanabe T.: The Bastille day (14 July 2000) event in historical large Sun-Earth connection events. *Solar Physics* 204, 423-436, 2001.
- Webber W.R., McDonald F.B., Lockwood J.A. and Heikkila B.: The effect of the July 14, 2000 “Bastille Day” solar flare event on >70 MeV galactic cosmic rays observed at V1 and V2 in the distant heliosphere. *Geophys. Res. Lett.*, 29(10), 1377, doi:10.1029/2002GL014729, 2002
- Wertz J. and Larson, W.J.: *Space Mission Analysis and Design*. third ed. Kluwer Academy, El Segundo, CA, U.S.A., 145, 1999.
- 455 Yermolaev, Y.I., Lodkina, G.I., Nikolaeva, N.S. and Yermolaev, M.Y.: Geoeffectiveness of Solar and Interplanetary Structures and Generation of Strong Geomagnetic Storms, in *Extreme Events in Geospace - Origins, Predictability and Consequences*, ed: N. Buzulukova, 99-113, doi: <https://doi.org/10.1016/B978-0-12-812700-1.00004-2>, 2018.