

# Seasonal evolution of winds, atmospheric tides and Reynolds stress components in the Southern hemisphere mesosphere/lower thermosphere in 2019

Gunter Stober<sup>1</sup>, Diego Janches<sup>2</sup>, Vivien Matthias<sup>3</sup>, Dave Fritts<sup>4,5</sup>, John Marino<sup>6</sup>, Tracy Moffat-Griffin<sup>10</sup>, Kathrin Baumgarten<sup>7</sup>, Wonseok Lee<sup>8</sup>, Damian Murphy<sup>9</sup>, Yong Ha Kim<sup>8</sup>, Nicholas Mitchell<sup>11</sup>, and Scott Palo<sup>6</sup>

<sup>1</sup>Institute of Applied Physics & Oeschger Center for Climate Change Research, Microwave Physics, University of Bern, Bern, Switzerland

<sup>2</sup>ITM Physics Laboratory, Mail Code 675, NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA

<sup>3</sup>German Aerospace Centre (DLR), Institute for Solar-Terrestrial Physics, Neustrelitz, Germany

<sup>4</sup>GATS, Boulder, CO, USA

<sup>5</sup>Center for Space and Atmospheric Research, Embry-Riddle Aeronautical University, Daytona Beach, FL, USA

<sup>6</sup>University of Colorado Boulder, Colorado, USA

<sup>7</sup>Fraunhofer Institute for Computer Graphics Research IGD, Rostock, Germany

<sup>8</sup>Department of Astronomy, Space Science and Geology, Chungnam National University, Daejeon 34134, South Korea

<sup>9</sup>Australian Antarctic Division, Kingston, Tasmania, Australia

<sup>10</sup>British Antarctic Survey, UK

<sup>11</sup>University of Bath, Bath, UK

**Correspondence:** [gunter.stober@iap.unibe.ch](mailto:gunter.stober@iap.unibe.ch)

**Abstract.** In this study we explore the seasonal variability of the mean winds, diurnal, semidiurnal tidal amplitude and phases as well as the Reynolds stress components during 2019, utilizing meteor radars at six southern hemisphere locations ranging from mid- to polar latitudes. These include Tierra del Fuego, King Edward Point on South Georgia island, King Sejong Station, Rothera, Davis and McMurdo stations. The year 2019 was exceptional in the southern hemisphere, due to the occurrence of a rare minor stratospheric warming in September. Our results show a substantial longitudinal and latitudinal seasonal variability of mean winds and tides pointing towards a wobbling and asymmetric polar vortex. Furthermore, the derived momentum fluxes and wind variances, utilizing a recently developed algorithm, reveal a characteristic seasonal pattern at each location included in this study. The longitudinal and latitudinal variability of vertical flux of zonal and meridional momentum is discussed in the context of polar vortex asymmetry, spatial and temporal variability, and the longitude and latitude dependence of the vertical propagation conditions of gravity waves. The horizontal momentum fluxes exhibit a rather consistent seasonal structure between the stations while the wind variances indicate a clear seasonal behaviour and altitude dependence showing the largest values at higher altitudes during the hemispheric winter and two variance minima during the equinoxes. Also the hemispheric summer mesopause and the zonal wind reversal can be identified in the wind variances.

Gravity waves (GW) originating at the lower atmosphere by a number of sources are an essential driver of the mesosphere/lower thermosphere (MLT) dynamics, forcing a meridional flow due to a zonal drag, which drives the mesopause temperature up to 100 K away from the radiative equilibrium (e.g., Lindzen, 1981; Becker, 2012) introducing a residual circulation from the cold summer to the warm winter pole. This important coupling mechanism is caused by GWs carrying energy and momentum from their source regions to the altitude of their breaking, coupling different vertical layers in the atmosphere (Fritts and Alexander, 2003; Ern et al., 2011; Geller et al., 2013). The primary forcing of the MLT at small scales is by gravity waves arising from various tropospheric sources, among them flow over orography (mountain waves), deep convection (convective gravity waves), frontal systems, and jet stream imbalances and shear instabilities (Fritts and Nastrom (1992), also see the review by Fritts and Alexander (2003)) and Plougonven and Zhang (2014). These various GWs typically have horizontal phase speeds comparable to the mean winds at higher altitudes, hence they are strongly influenced by varying winds along their plane of propagation. GWs can propagate upward until they become dynamical unstable or they are filtered by critical levels, where they undergo breaking and dissipation, resulting in local mean flow accelerations that act as sources of non-primary GWs.


GW breaking dynamics occurs on relatively small horizontal scales, 10-100 km; whereas non-primary GW dynamics occurs at larger scales, 100-300 km and arise due to the local, transient mean-flow accelerations accompanying GW momentum transport (Dong et al., 2020; Fritts et al., 2020). Non-primary GWs at larger scales also arise due to interactions among larger-scale GWs in global models unable to resolve GW breaking dynamics (Becker and Vadas, 2018; Vadas and Fritts, 2001; Vadas and Becker, 2018). Importantly, however, non-primary GWs accompanying GW breaking and interactions at lower altitudes require propagation over large depths to become significant, hence, they play more significant roles in the lower thermosphere. Although GWs are such an important driver of the MLT, the number of observations is rather sparse. Very often the GW activity is inferred by subtracting a background from the wind or temperature observations to estimate potential GW energy or wind variations (Ehard et al., 2015; Baumgarten et al., 2017; Chu et al., 2018; Rüfenacht et al., 2018; Stober et al., 2018b; Wilhelm et al., 2019). Satellite observations provide an estimate of absolute momentum fluxes from the troposphere up to the mesosphere and most importantly a global coverage (Ern et al., 2011; Trinh et al., 2018; Hocke et al., 2019). However, satellite observations are lacking the directional information and, thus, there is some ambiguity about the forcing or whether the GW momentum flux is accelerating or decelerating the mean flow.

Vincent and Fritts (1987) introduced, over two decades ago, a radar technique to determine the vertical flux of zonal and meridional momentum utilizing Medium Frequency (MF)-radars using two pairs of co-planar beams. This technique was also applied by Placke et al. (2015a, b) to determine momentum fluxes above Andenes in Northern Norway. However, there are only a few MF-radars worldwide that are able to conduct such measurements. Furthermore, at altitudes above 94 km MF-radars tend to underestimate the wind speeds, which might lead to some systematic bias in the derived momentum flux (Wilhelm et al., 2017). Hocking (2005) presented a method to obtain the Reynolds stress tensor components from the meteor radar observations. Based on this, several studies applied the method to optimize the data analysis as it appeared to be challenging to get the technique implemented (Placke et al., 2011a, b; Andrioli et al., 2013). Fritts et al. (2010b) and Fritts et al. (2012b) presented

a momentum flux meteor radar design to overcome some of the difficulties and evaluated the momentum flux observations using synthetic data (Fritts et al., 2010a), which finally provided evidence that these systems can be used to measure reliably momentum fluxes. This led to several studies using these new generation systems (de Wit et al., 2014, 2016, 2017; Spargo et al., 2019; Vierinen et al., 2019) or more powerful radars such as MU-radar (Riggin et al., 2016).

Climatologies of mean winds and tides are also rather sparse at the southern hemisphere (SH) and are essentially affected by the vertical coupling of upward propagating gravity waves, but also provide a temporal and spatially variable background for the gravity wave propagation itself. Recent studies with general circulation models have shown that mean winds are essential to understand the GW forcing (Liu, 2019; Shibuya and Sato, 2019). This is also the case for tides, which provide essentially temporal variable critical filtering for the vertical propagation of the GWs (Heale et al., 2020). In the past there were several studies about mesospheric winds or tides, which were often limited to a single station or investigated only a certain tidal or wave component (Batista et al., 2004; Fritts et al., 2010b, 2012b) and for tides (Beldon and Mitchell, 2010; Conte et al., 2017). Recently, Liu et al. (2020) used several meteor radars at the southern hemisphere to systematically investigate the 8-hour and 6-hour tides and to obtain a more comprehensive picture of the latitudinal and longitudinal characteristics of these tidal modes. Furthermore, meteor radar observations have turned out to provide a valuable and independent method to validate general circulation models with data assimilation (McCormack et al., 2017) or to investigate the interday variability of tidal amplitudes and phases (Stober et al., 2019).

In this study, we present a cross-comparison of mean winds and the diurnal and semidiurnal tide for six southern hemispheric meteor radars located at mid- and polar latitudes. We present observations from 2019 and investigate the latitudinal and longitudinal differences of these meteorological parameters at each radar site to provide a comprehensive overview and systematic analysis of the wind, and tidal and gravity wave dynamics by applying a unified diagnostic. The meteor radars are located at Tierra del Fuego (TDF), King Edward Point (KEP) on South Georgia island (Jackson et al., 2018), at King Sejong Station (KSS) on King George Island (Lee et al., 2018, 2016), and at Rothera (ROT) Station (Sandford et al., 2010) located on the Antarctic Peninsula, Davis Antarctic Station (DAV) (Holdsworth et al., 2004) and McMurdo (MCM) Antarctic Station. We discuss the presented results within the context of the stratospheric polar vortex for the year 2019. For this purpose, we complement our meteor radar observations with data from the Microwave Limb Sounder (MLS) on board the AURA satellite (Livesey et al., 2006; Schwartz et al., 2008). Furthermore, we utilize a recently developed retrieval algorithm, which builds on the initial momentum flux analysis formulation reported by Hocking (2005). In particular, we introduce a generalized approach to obtain wind variances and momentum fluxes from several meteor radars (many of which are standard low power systems) for the year 2019, which evolved into one of the rare minor stratospheric warming events during September (Yamazaki et al., 2020). We briefly summarize how the Reynolds stress components, also called momentum fluxes and wind variances, are derived from a Reynolds decomposition. The Reynolds decomposition is achieved by utilizing an adaptive spectral filter (ASF), which allows the decomposition of the meteor radar wind time series into mean winds, tidal components and a GW residual (Stober et al., 2017; Baumgarten and Stober, 2019; Stober et al., 2019) similar to the S-transform used in previous studies (Stockwell et al., 1996; Fritts et al., 2010a). The manuscript is structured as follows: In section 2 we present a brief introduction to the wind retrievals, the derivation of the Reynolds stress components and the implemented momentum flux and GW retrieval. Section 3



**Figure 1.** Stereographic-projection of the geographic location of the meteor radars used in this study and a map of the terrain elevation of Antarctica, the Antarctic Peninsula and Southern Argentina to visualize the orography around each radar station. The maps are generated using etopo1 data (Amante and Eakins, 2009).

contains the results of the mean flow terms, which are mean winds, diurnal and semidiurnal tides and their seasonal behaviour,  
 85 as well as the determined momentum fluxes and wind variances in this section. Our results are discussed in section 4 and the  
 conclusions are provided in section 5.

## 2 Observations and methods

### 2.1 Meteor radar observations

In this study, we use observations obtained with 6 meteor radars operating in the SH between  $53^{\circ}\text{S}$  and  $79^{\circ}\text{S}$  in latitude. Four  
 90 of the meteor radars can be grouped into a cluster around the Drake passage consisting of the Southern Argentina Agile Meteor  
 Radar (SAAMER) at Rio Grande, Tierra del Fuego Argentina (hereafter referred to as TDF), King Sejong Station on King  
 George Island (KSS) and Rothera (ROT) on the Antarctic Peninsula and King Edward Point (KEP) on South Georgia island.  
 The other two radars are located almost opposite of the Drake passage at McMurdo (MCM) and Davis Antarctic stations  
 (DAV). Figure 2.1 shows two panels with stereographic projections of the SH where the radar locations are represented by red  
 95 dots (left panel and Table 2.1). The right panel in this figure shows a color contour map of the mean elevations around each  
 radar system to identify potential orographic wave forcing sources underneath the observation volumes.

A technical summary of the radars is provided in Table 2.1. Most of the systems have been in operation for more than a decade  
 and have provided reliable and continuous observations. Although most of these systems have been operated without major  
 parameter changes, both ROT and TDF meteor radars have been upgraded during the observing period of our study. Until  
 100 February 2019, the ROT system used a high pulse repetition frequency (PRF) meteor mode with a PRF of 2144 Hz, a 2 km  
 range sampling and 4 coherent integrations. After this time, the system was upgraded and resumed operation transmitting a

**Table 1.** Technical parameters of the meteor radars

	TDF	KEP	KSS	ROT	DAV	MCM
	Tierra del Fuego	King Edward Point	King Sejong Station	Rothera	Davis	McMurdo
Freq. (MHz)	32.55	35.24	33.2	32.5	33.2	36.170
Power (kW)	64	6	12	6	7	30
PRF (Hz)	625	625	440	2144/625	430	500
coherent integration	1	1	4	4/1	4	1
pulse code	7-bit Barker	7-bit Barker	4-bit complementary	mono/Barker	4-bit complementary	7-bit Barker
sampling (km)	1.5	1.5	1.8	2/1.5	1.8	1.5
location (lat,lon)	53.7°S, 67.7°W	54.3°S, 35.5°W	62.2°S, 58.8°W	67.5°S, 68.0°W	68.6°S, 78.0°E	77.8°S, 166.7°E

7-bit Barker code with 1.5 km range sampling and a PRF of 625 Hz. We also noted a significant noise or interference at ROT before the upgrade in January/February that did not allow to derive trustworthy momentum fluxes. Further, we also restricted our analysis of mean winds and tides to the altitude range between 80-100 km. In addition, in September 2019, the TDF transmitting scheme also changed. The original design of the TDF transmitter (TX) configuration used eight three-element crossed Yagis arranged in a circle of diameter 27.6 m. Each transmitting in opposite phasing of every other Yagi (Janches et al., 2014). In 2019, the system transmission strategy was upgraded with the deployment of a single new TX antenna with the goal of improving the detection rate of meteors at larger zenith angles for astronomical purposes (Janches et al., 2020). By concentrating the full power of TDF in one TX antenna, a more uniform detection pattern is achieved that satisfies this original requirement, but also increases the number of events detected at larger zenith angles. Finally, the MCM radar is the most recent installation which, although it is not the most powerful radar, provides a very good altitude coverage. This is partly explained by the sporadic meteor sources and the southern location of the MCM meteor radar. The helion, antihelion and the south apex meteor source are above the local horizon all the time contributing to the observed sporadic meteor fluxes at MCM, yielding a much weaker seasonality in the altitude variation of the meteor layer (Janches et al., 2004). In addition, meteors arriving from these sources enter the atmosphere at fairly low entry angles ( $<20^\circ$  see Schult et al. (2017) for a northern hemisphere radar), leading to a much smoother ablation profile of the meteoroids and, hence, the released meteoric material is spread over a larger segment of the meteor flight path increasing the detectability. On the other side, the orbit geometry alone provides not yet a sufficient explanation for the better altitude coverage at McMurdo, however, a more detailed investigation is beyond the scope of the paper. Several of the meteor radars used in this study employ the standard meteor radar configuration an array of 5-Yagi antennas for reception with a spacing of 2 and  $2.5 \lambda$  (Jacobs and Ralston, 1981; Jones et al., 1998). MCM was set up in a different configuration with 1.5 and  $2 \lambda$  spacing due to topographic constraints. Similar to other meteor radars most

of them use a single Yagi antenna for transmission. Only TDF employed a beam forming transmission scheme resulting in 8 main beams, as described earlier, but changed to the use of the single crossed Yagi antenna late during the period studied here (Janches et al., 2014; Janches et al., 2020). A more detailed description of the King Sejong Station meteor radar can be found in Lee et al. (2018) and for the DAV meteor radar in Holdsworth et al. (2004).

## 2.2 Retrieval of winds and momentum flux

Meteor radars have been used to measure winds in the mesosphere/lower thermosphere (MLT) for several decades. Typically winds are obtained by least square fits solving for the horizontal wind velocities after binning the data into altitude and time intervals (Hocking et al., 2001; Holdsworth et al., 2004). In this study we retrieve winds using the algorithm presented in Stober et al. (2018a), which includes the treatment of the geometry of the full Earth, based on the WGS84 rotation ellipsoid to provide more precise altitude estimates and geodetic coordinates for each meteor, a spatio-temporal Laplace filter and a non-linear error propagation, which is described in more detail in Gudadze et al. (2019). The wind retrievals are cross-validated against NAVGEM-HA (McCormack et al., 2017; Stober et al., 2019). The results presented in this manuscript are based on winds with a temporal resolution of 1 hour and a vertical resolution of 2 km. The minimum number of meteors per time and altitude bin for a successful fit is 4.

For the case of momentum fluxes, Hocking (2005) proposed a method using typical meteor radars, which was later echoed and reformulated as correlations by Vierinen et al. (2019). In this work, we present a brief derivation of the Reynolds stress tensor and show how the different tensor elements are estimated from the meteor radar observations. The starting point is the well-known radial wind equation. Each meteor will form a trail that will be detected by the radar and will drift with the background wind. The radar will then detect that radial velocity, via Doppler shift in the received signal, and the three components of the background wind can be calculated for each detected meteor using the mathematical convention (reference to east and counterclockwise rotation);

$$v_{\text{rad}} = u \cdot \cos(\phi) \sin(\theta) + v \cdot \sin(\phi) \sin(\theta) + w \cdot \cos(\theta) \quad , \quad (1)$$

where  $u$ ,  $v$ ,  $w$  are the three wind components (zonal, meridional and vertical, respectively),  $\phi$  is the azimuth angle,  $\theta$  is the off-zenith angle and  $v_{\text{rad}}$  is the observed radial wind velocity. Further, it is straight forward to use the standard Reynolds decomposition of the wind, separating the wind components into a mean flow ( $\bar{u}$ ,  $\bar{v}$ ,  $\bar{w}$ ) and wind fluctuations ( $u'$ ,  $v'$ ,  $w'$ );

$$u = \bar{u} + u' \quad (2)$$

$$v = \bar{v} + v'$$

$$w = \bar{w} + w' .$$

As we are mainly interested in the momentum flux associated to GWs, the mean flow terms containing the background wind and the diurnal and semidiurnal tide have to be subtracted/removed from the observed radial velocities for each meteor. Thus, we model the mean flow radial velocity by;

$$\overline{v_{\text{radm}}} = \bar{u} \cdot \cos(\phi) \sin(\theta) + \bar{v} \cdot \sin(\phi) \sin(\theta) + \bar{w} \cdot \cos(\theta) \quad , \quad (3)$$

The radial velocity fluctuations ( $v'_{\text{rad}}$ ), which now only contain GW contributions, are obtained by subtracting the mean flow  
 155 radial velocity ( $\overline{v_{\text{radm}}}$ ) from the observed radial velocity ( $v_{\text{rad}}$ ) measurements;

$$v'_{\text{rad}} = v_{\text{rad}} - \overline{v_{\text{radm}}} . \quad (4)$$

Furthermore, the GW fluctuations can be modelled by;

$$v'_{\text{radm}} = u' \cdot \cos(\phi) \sin(\theta) + v' \cdot \sin(\phi) \sin(\theta) + w' \cdot \cos(\theta) . \quad (5)$$

Considering that these radial velocity fluctuations are mostly driven by GW and, hence, the Reynolds stresses can be computed  
 160 by minimizing the following quantity (Hocking, 2005);

$$\Lambda = \sum \left( (v'_{\text{rad}})^2 - (v'_{\text{radm}})^2 \right)^2 . \quad (6)$$

Inserting eq. 5 into eq. 6 leads to the well-know momentum flux terms;

$$\Lambda = \sum \left( (v'_{\text{rad}})^2 - \left( u'^2 \cdot \cos(\phi)^2 \sin(\theta)^2 + v'^2 \cdot \sin(\phi)^2 \sin(\theta)^2 + w'^2 \cdot \cos(\theta)^2 + 2u'v' \cdot \cos(\phi) \sin(\phi) \sin(\theta)^2 + \right. \right. \quad (7)$$

$$\left. \left. 2u'w' \cdot \cos(\phi) \sin(\theta) \cos(\theta) + 2v'w' \cdot \sin(\phi) \sin(\theta) \cos(\theta) \right) \right)^2 .$$

165 Solving eq. 7 for the unknown Reynolds stresses components is straight forward. Typically, the terms  $u'w'$ ,  $v'w'$  and  $u'v'$  are also called momentum fluxes and the symmetric Reynolds stress tensor is given by;

$$\tau'_{ij} = \rho \overline{u'_i u'_j} = \rho \cdot \begin{pmatrix} u'^2 & u'v' & u'w' \\ u'v' & v'^2 & v'w' \\ u'w' & v'w' & w'^2 \end{pmatrix} , \quad (8)$$

where  $\rho$  is the atmospheric density at the altitude of the measurement and the other terms in the tensor denote the Reynolds  
 stress components (wind variances and momentum fluxes), which have units of squared velocity fluctuations. The Reynolds  
 170 stress are derived from the RANS (Reynolds Averaged Navier Stokes) equations assuming an incompressible Newtonian fluid  
 and that the Reynolds average of the fluctuations vanishes ( $\overline{u'} = 0$ ), which requires that the averaging has to be long enough to  
 cover the inertia GW periods of several hours or longer. The spatial scales can theoretically be estimated by selecting different  
 volumes inside the domain area, however, practically the meteor statistics is often not sufficient to get reliable results.

However, there are some caveats of the theory outlined above, when it comes to implement the algorithm and actually to apply  
 175 it to meteor radar observations. One difficulty is the Reynolds decomposition into the mean flow and the GW fluctuations.  
 Previous studies often limited the analysis to a narrower angular region (Fritts et al., 2010a; Placke et al., 2015a) using only  
 off-zenith angles between 10-50° reducing significantly the number of meteors for the analysis per time and altitude bin, which  
 in turn required longer averaging or was achieved by an active beam forming antenna. Such an antenna directed more energy  
 towards an angular region as for TDF (Fritts et al., 2010b, a) or the meteor radar at Trondheim (de Wit et al., 2014). The  
 180 process by which this much stricter angular selection of meteors improved the momentum flux estimates was the reduction  
 of projection errors due to the Earth's ellipsoid shape, which caused apparent and arbitrary contributions to the fluctuation

terms. Stober et al. (2018a) proposed to minimize this type of uncertainty by computing for each meteor its geodetic position relative to the WGS84 reference ellipsoid, which improves the altitude determination, but also reduces projection errors for the azimuth and off-zenith angle. The benefit of this full Earth geometry correction is that there is no longer a need to reduce  
185 the angular region and all meteors up to off-zenith angles of  $65^\circ$  can be used for the analysis. Typical specular meteor radars (single Yagi-antenna on transmission) detect most meteors at off-zenith angles between  $50^\circ$  and  $70^\circ$ . However, meteors at larger zenith angles are further away from the radar and, thus, are more prone to altitude errors. Considering that the typical angular precision of the employed receiver arrays is approximately  $1.5\text{-}1.7^\circ$  (Jones et al., 1998). A limit of  $65^\circ$  presents a more optimal choice to maximize the number of meteors entering the analysis while keeping a sufficient altitude precision. Another  
190 important aspect related to the momentum flux estimation is the proper removal of the background flow, which was already outlined by Fritts et al. (2010a); Placke et al. (2011b); Andrioli et al. (2013) and later confirmed by de Wit et al. (2014). In particular, tides have large amplitudes in the MLT, causing large vertical and temporal shears within a time and altitude bin. Noting that Hocking (2005) and Placke et al. (2011b) suggested the use of at least 30 meteors for a successful momentum flux fit, which is often achieved by temporal averaging, the importance of the temporal shear becomes evident.

195 In this study we use the adaptive spectral filter (ASF) to perform the Reynolds decomposition to characterize the background flow and the GW fluctuations. A first version of the ASF(1D) (temporal domain) was presented in Stober et al. (2017). Here we make use of the ASF(2D), which employs a vertical regularization constraint for the mean wind and tides assuming a smooth vertical phase progression for each wave without an explicit vertical wavelength threshold (Pokhotelov et al., 2018; Baumgarten and Stober, 2019; Wilhelm et al., 2019). The ASF accounts for the continuous variation of the mean flow as well as for the  
200 intermittent behaviour of the tides. Thus, we obtain hourly resolved background wind fields for each altitude and time bin for the zonal, meridional and vertical wind component, respectively. This background wind field contains the mean flow and the diurnal, semidiurnal and terdiurnal tidal component. However, the terdiurnal tide has usually a much smaller amplitudes (Liu et al., 2020) compared to the diurnal and semidiurnal tides and thus, is not further discussed here. Furthermore, we perform a linear interpolation of the background wind field to the actual occurrence time and altitude for each meteor to estimate  
205 the  $v_{\text{radm}}$ -term minimizing any contribution from the background flow. This procedure is effective in mitigating possible contamination due to tides, and permits to the use of a much longer averaging window. In this study we use 64-hours and a minimum of 100 meteors to determine the Reynolds stress components. However, for the seasonal climatology only solutions with more than 1000 meteors enter the statistics.

The algorithm is implemented similar to that performed for wind retrievals in Stober et al. (2018a). The first guess is  
210 provided by a classical least square fit. Based on this initial iteration, we compute the spatio-temporal Laplace filter, which provides a predictor for each time and altitude bin. This predictor enters as regularization (Tikhonov) all further iterations and is updated each time. The spatio-temporal Laplace filter turns out to be beneficial for ill-conditioned problems due to the random occurrence of meteor detections and asymmetries in the spatial sampling; these can result in difficulties in determining all parameters with similar quality.

215 Furthermore, we perform a non-linear error propagation similar to the one presented in Gudadze et al. (2019). The statistical uncertainties are updated in each iteration step. We also tested barrier functions<sup>2</sup> to penalize negative values of  $u'^2$ ,  $v'^2$  and  $w'^2$ ,



respectively. Such negative values were reported in Placke et al. (2011a), but this appears to be a minor issue in our retrievals. Only a negligible number of fits resulted in negative values for just some of the radar systems utilized in this work.

220 In addition, we performed some test retrievals to account for the vertical velocity bias intrinsic to the meteor radar observations. Specular meteors have trail lengths of up to several kilometers where the radio waves are scattered and, thus, meteors entering the Earth's atmosphere at steep entry angles can encounter strong vertical wind shears, which lead to a rotation of the trail causing systematic errors. In particular, during the local summer months this can lead to a systematic deviation of a few cm/s for mid-latitude stations.

225 Very often wind fits are performed by assuming  $w = 0$  m/s (Hocking et al., 2001; Holdsworth et al., 2004). However, here we use the retrievals as presented in Stober et al. (2018a) who used the vertical wind velocity as quality control. Typically, we obtain daily mean values of the order of  $\pm 0.25$  m/s, which is more than an order of magnitude less than reported by Egito et al. (2016). However, the remaining bias due the vertical winds, which potentially has the wrong sign, had no impact on the retrieved Reynolds stresses. Finally, in order to get confidence in the retrievals, we performed several test cases similar to the ones presented in Fritts et al. (2010a). Therefore, we extracted the observed meteor detections from TDF and synthesized wind 230 fields including altitude dependent mean winds and tides with a vertical wavelength of 80 km and various altitude dependent GW fields to optimize the retrieval setting with respect to the regularization strength, the required statistics and the applied averaging. Performing these tests we find minor deviations from the synthetic wind and GW fields only at the upper and lower edges of the meteor layer. The tidal amplitudes were retrieved within  $\pm 2$  m/s compared to the synthetic data. The momentum fluxes agreed for the 30-day median remarkably well. We also tested the possibility to retrieve the vertical wind fluctuation 235 amplitudes and found mean deviations of  $\pm 0.01$  m/s residual bias for the synthetic fields and about  $\pm 0.25$  m/s bias in our observations.

### 2.3 MLS satellite observations

To bring the local radar observations into the global context we calculate the geostrophic zonal wind as described in Matthias and Ern (2018) from geopotential height (GPH) data from the Microwave Limb Sounder (MLS) on board the Aura satellite 240 (Waters et al., 2006; Livesey et al., 2015). MLS has a global coverage from  $82^\circ\text{S}$  to  $82^\circ\text{N}$  on each orbit, and a usable height range from approximately 11 to 97 km ( $261 - 0.001$  hPa) with a vertical resolution of  $\sim 4$  km in the stratosphere and  $\sim 14$  km at the mesopause. The temporal resolution is 1 day at each location, and data are available from August 2004 until the present (Livesey et al., 2015). Version 4 MLS data were used in this paper along with the application of the most recent recommended quality screening procedures from Livesey et al. (2015). For our analyses the original orbital MLS data are accumulated in grid 245 boxes with  $10^\circ$  grid spacing in longitude and  $5^\circ$  in latitude. Afterwards they are averaged at every grid box and for every day, generally resulting in a global grid with values at every grid point.

### 3 Results

#### 3.1 Mean winds 2019


As pointed out in the previous section, we perform a Reynolds decomposition in order to separate a mean flow from the GW  
250 fluctuations. Thus, we analyze the data with the adaptive spectral filter (ASF) technique (Baumgarten and Stober, 2019) to  
obtain daily mean winds, as well as diurnal and semidiurnal tides.

We first compare the seasonal zonal and meridional winds of all 6 locations to identify any seasonal and local differences. Fig-  
ure 2 shows the seasonal zonal and meridional wind pattern during 2019 obtained from the daily mean zonal and meridional  
winds after applying a 30-day running median shifted by one day. This reveals any seasonal variability by removing atmo-  
255 spheric waves with shorter periods. A similar analysis was applied in Wilhelm et al. (2019) for meteor radars in the Northern  
hemisphere in order to derive mean wind climatologies. Significant differences between the locations can be observed from this  
figure, in particular during the SH winter seasons (JJA). Both, TDF and KEP observations, operated at almost the same latitude  
at  $54^{\circ}\text{S}$ , show a similar morphology for the zonal winds and only the meridional winds deviate during June and July from each  
other. At the tip of South America, TDF shows that the meridional winds experience a sign reversal around June/July, which is  
260 not present over KEP. The meridional winds seem to also have a semiannual oscillation at both locations.

Further polewards at KSS and ROT ( $62^{\circ}\text{S}$  and  $67^{\circ}\text{S}$ , respectively), which are at a similar longitude as TDF, the zonal winds  
reflect a similar seasonal behaviour compared to KEP and TDF, but with a slightly weaker wind magnitude. However, the  
meridional winds are fairly consistent during the summer months compared to the mid-latitude radars, but deviate consid-  
erably during the winter season. There is even a noticeable difference between KSS and ROT, even though the systems are  
265 located rather close together. At ROT and KSS the meridional winds show only during April, May and June a typical winter  
behaviour and approximately northward winds for the other months above 80-85 km. Only during September and at altitudes  
above 90 km above KSS a short southward wind patch occurs.

Comparing the observed wind fields measured at ROT and DAV, which are only separated by 2 degrees in latitude, but by  $170^{\circ}$   
in longitude further emphasizes the existence of a significant asymmetry in the southern hemispheric wind systems. As expected,  
270 looking at the general morphology, the seasonal zonal wind pattern for 2019 is remarkably similar between both locations.  
There are only marginal differences in the zonal magnitudes considering the overall agreement of the zonal wind structures.  
This is also the case for the meridional winds during the summer months (DJF). However, during the winter season the merid-  
ional wind structure is significantly different between both stations. The morphology at DAV appears to be less asymmetric  
with a tendency to show increased southward wind magnitudes towards the end of the winter season, whereas at ROT the  
275 highest southward winds are registered at the begin of the winter season 2019.

The southernmost location in our analysis is MCM at  $78^{\circ}\text{S}$ . The seasonal zonal wind morphology compares well with those  
measured at DAV and ROT, but shows much weaker wind magnitudes. Similar to observations in the Northern hemisphere the  
summer zonal wind reversal altitude also increases with increasing southern latitude. Compared to DAV, the meridional winds  
are intensified during the summer and winter seasons. Furthermore, the asymmetry during the winter months is also present  
280 at MCM, which shows, similarly to DAV, the highest southward meridional winds towards the end of the winter season as a



**Figure 2.** Comparison of zonal and meridional mean winds for each station during the year 2019. The left column shows the zonal component and the right column the meridional wind. All stations are sorted according to their latitude from TDF, KEP, KSS, ROT, DAV to McM.


double structure. In fact, the southward meridional winds at MCM during July and September 2019 are the strongest of all locations.

### 3.2 Diurnal tidal amplitudes and phases measured during 2019


Atmospheric tides provide a time variable background filter for the vertical propagation of GWs, which can, depending on the tidal phase and the propagation direction of the GW, lead to GW breaking and dissipation. These breaking events might trigger/foster the generation of secondary or non-primary waves (Heale et al., 2020). Thus, tides are essentially contributing to the Reynolds decomposition. In particular, the day-to-day variability is crucial for the momentum flux analysis. Typically, atmospheric tides are derived assuming phase stability over a certain period of time, which can be several days, weeks or months (Murphy et al., 2006; Hoffmann et al., 2010; Conte et al., 2017; He et al., 2018; Pancheva et al., 2020). More recent studies favor much shorter windows of 24 to 48 hours to account for the intermittent behaviour of tides (Stober et al., 2017; Wu et al., 2019; de Araújo et al., 2020; Das et al., 2020), in particular, phases of atmospheric tides that appear not to be constant with time (Ward et al., 2010; Baumgarten and Stober, 2019; Stober et al., 2019). In this study, all tidal amplitudes and phases were determined with the ASF which, similar to wavelet spectra, adapts the window length to the period of the fitted frequencies. The obtained daily tidal amplitudes are vector averaged using 30-day medians centered at the respective day to derive the seasonal variation.

Figure 3 presents the seasonal variation of the diurnal tidal amplitudes measured at each station. Although the daily mean winds showed significant differences between TDF, KEP, KSS and ROT, the seasonal behaviour of the diurnal tide is rather consistent between all four locations. There is a pronounced summer maximum in the zonal and meridional amplitudes from January to February at altitudes from 78-106 km. The meridional tidal amplitudes tend to exceed the zonal amplitude by up to 10 m/s. At altitudes above 100 km the diurnal tide remained of significant magnitude until May 2019. Apparently, for the other months the diurnal tidal amplitudes remained fairly ( $< 10$  m/s) at altitudes between 80-100 km during 2019. KSS and ROT indicate a small diurnal tidal enhancement for July/August and in December below 80 km and above 100 km altitude. The December enhancements are also found at TDF, but almost disappear at KEP. DAV measurements show basically the same seasonal diurnal tide behaviour, but with weaker amplitudes. The winter diurnal tidal enhancement in June/July appears to be more pronounced. However, the southernmost meteor radar at MCM observes a significantly different seasonal diurnal tidal pattern. The summer maximum is much more pronounced compared to the other stations and shows amplitudes of 20 m/s from January to April at 90 km and above and again from October to December. There is also a noticeable difference between the zonal and the meridional diurnal tidal amplitude. The zonal component indicates a winter minimum, whereas the meridional component shows a tidal enhancement.

Diurnal tidal phases are shown in Figure 4. The tidal phases are given in UTC, hence, longitudinal differences are present as phase shifts. As expected the diurnal phases are much more variable during time with low tidal amplitudes for TDF, KEP, KSS and ROT. During the summer months of January and February 2019 the diurnal phases are more stable and indicate rather long vertical wavelengths, but with significant differences between the zonal and the meridional components. However, the phase plots indicate a distinct seasonal pattern showing phase drifts of several hours at the same altitude over the course of the year.



**Figure 3.** Same as Figure 2 but for the diurnal tidal amplitudes.



**Figure 4.** Same as Figure 2 but for the diurnal tidal phases.

315 The further more south the location of a meteor radar is, the less characteristic is the seasonal behaviour. Measurements at DAV and MCM indicate a decreased variability of the diurnal tidal phases throughout the year. At DAV there are periods suggesting almost phase stability over several weeks, instead of the typical continuous variation reflected by the other stations.

### 3.3 Semidiurnal tidal amplitudes and phases measured during 2019


At mid- and high latitudes, semidiurnal tides are the dominating tidal wave during the course of the year (Hagan and Forbes, 2002, 2003). Figure 5 shows the vector averaged semidiurnal tidal amplitudes measured by all six meteor radars using again a 30-day median shifted by one day in analogy to the mean winds and diurnal tides.

The seasonal structure of the semidiurnal tide reveals a rather interesting pattern for the SH. Semidiurnal tides measured at TDF, KEP, KSS and ROT show some similarities for the zonal component, resulting in amplitudes with values below  $< 10$  m/s during the summer months January to mid-March. From April to June all 4 stations show a strong semidiurnal tidal activity with amplitudes up to 40 m/s, another minimum of the tidal activity in July and a secondary maximum from August to the end of the year. Furthermore, the tidal amplitudes show a decrease with increasing polar latitude, which is also observed at the northern hemisphere. However, the meridional semidiurnal tide shows a clear longitude dependence and asymmetry compared to the zonal tidal amplitudes, which was not reported previously (Conte et al., 2017). At the longitude of TDF and ROT the meridional tidal component is much weaker during April to June compared to the zonal. At KSS, which is further East, similar amplitudes for the zonal and meridional component are observed. This was also found at KSS in a previous study reporting the tidal amplitudes under solar maximum conditions, which resulted in larger amplitudes of the semidiurnal tide (Lee et al., 2013). At KEP, which is  $25^\circ$  eastward, shows the opposite behaviour and the meridional component of the semidiurnal tide reaches the highest amplitudes in April to June. Such differences with longitude might be related to the superposition of migrating and non-migrating tides (Murphy et al., 2006).

335 The semidiurnal tidal seasonal behaviour observed at DAV looks quite different from the stations that are located further to the north. The amplitudes are much weaker and barely reach values of 25 m/s and there are 4 periods during which an increased activity is observed, which are during January-February, May, August-September and in December. The largest tidal amplitudes are observed during May 2019.


Further to the south, at MCM Station, the semidiurnal tide exhibits only a very faint seasonal structure. Most of the time the amplitudes are below 10 m/s. Only during March, May and November-December and below 90 km altitude there are periods where amplitudes exceed 10 m/s. This is surprising when we compare these values with measurements performed at geographically conjugate northern hemisphere latitude. For instance, at Svalbard ( $78.17^\circ$  N,  $15.99^\circ$  E) the semidiurnal tide still reflects a similar seasonal activity as other polar and mid-latitude locations (Wilhelm et al., 2019; Pancheva et al., 2020). This is obviously not the case in the SH and represents a remarkable interhemispheric difference.

345 Semidiurnal tidal phases are displayed in Figure 6 where it can be seen that the semidiurnal tidal phases reflect similar features than those present in the amplitudes. TDF, KEP, KSS and ROT show a very similar seasonal structure indicating continuous changes of the tidal phases throughout 2019 at all altitudes. At DAV and MCM, on the other hand, the observed phases indicate an even more pronounced seasonal structure and faster gradual phase drifts. In particular, at MCM the phases appear to be




**Figure 5.** Same as Figure 2 but for the semidiurnal tidal amplitudes.





**Figure 6.** Same as Figure 2 but for the semi-diurnal tidal phases.



**Figure 7.** Comparison of semidiurnal tidal vertical wavelengths of a) TDF, b) KEP, c) KSS, d) ROT, e) DAV and f) McM.

more variable, which is likely due to the generally weaker amplitudes pointing towards a much weaker and more intermittent excitation of the tides. Comparing the seasonal phase behaviour of the SH to those measured at conjugate latitudes in the northern hemisphere indicates that there are some differences. In the northern hemisphere, from mid- to high latitudes, the semidiurnal tides show a seasonal asymmetry between the winter to summer transition and the fall transition (Portnyagin et al., 2004; Wilhelm et al., 2019; Stober et al., 2019). The fall transition is accompanied by a significant phase change from September to November, whereas in the SH this feature is very weak at TDF and ROT (March to May), and almost negligible for KEP and KSS.

Finally, we briefly discuss the presence of a potential lunar tide. Sandford et al. (2006, 2007) estimated the lunar tide amplitude from two northern hemispheric meteor radars and Davis MF-radar in the SH and found values of 1-2 m/s, which is negligible compared to the typical GW amplitudes of about 20-30 m/s for the resolved waves. However, Forbes and Zhang (2012) investigated a potential lunar tide amplification due to the Pekeris resonance. They found favorable conditions to shift the Pekeris peak towards the lunar tide periods M2 (12.42 h) and N2 (12.66 h), during the time of major sudden stratospheric warming in 2009 in the northern hemisphere, as only during the time of the wind reversal the vertical temperature and wind structure satisfies the resonance condition. Later, Zhang and Forbes (2014) argued that the Pekeris resonance peak is rather broad and, thus, more or less each sudden stratospheric warming can cause a lunar tide amplification.

These reports triggered several studies investigating the lunar tide and its relevance for the mesosphere dynamics. However, most of the observational diagnostics (wavelet or harmonic fitting) separating the lunar tide from the semidiurnal tide applied long windows of 21-days or even longer periods up to several months assuming phase stability of the semidiurnal tide (Forbes and Zhang, 2012; Chau et al., 2015; Conte et al., 2017; He et al., 2018; Siddiqui et al., 2018). However, as shown in Figure 6, the semidiurnal tidal phase shows considerable variability and seasonal changes and, thus, the assumption of phase stability for the semidiurnal tide is not valid. Therefore, we performed a holographic analysis to test whether a temporally variable semidiurnal tidal phase could be misinterpreted as a lunar tide (see appendix A1) (Stober et al., 2019). In fact, the holograms often exhibit a shift towards the M2 frequency (12.42 h) uncorrelated with the lunar orbit. Given these results and considering that there was only a minor stratospheric warming in September 2019 (Yamazaki et al., 2020), we consider the lunar tide as a minor wave with a negligible amplitude compared to GWs and, we did not make an attempt to remove this tidal component in our Reynolds decomposition.

For the sake of completeness, we also estimated the vertical wavelengths of the semidiurnal tide, which is presented in Figure 7. The vertical wavelength provides a good overview to identify potential changes in the Hough modes of the tide, which are solutions of the Laplace tidal differential equation (Lindzen and Chapman, 1969; Wang et al., 2016). The vertical wavelengths show a similar latitude and longitude dependence as already discussed for the semidiurnal tidal amplitudes and phases. The observations at TDF, KEP, KSS and ROT indicate almost the same vertical wavelengths from March to October 2019 of about 70-100 km. This corresponds to the time with the largest semidiurnal tidal amplitudes. However, the seasonal summer months January-February and November-December show a longitudinal difference. KEP and KSS observe much longer vertical wavelengths of up to 1000 km during these months, compared to the stations located to the west. These very long vertical wavelengths are associated to times with a small semidiurnal tidal amplitude. The results obtained at DAV reflects

a slightly different seasonal behaviour. There, the longest vertical wavelengths are observed in March-April followed by a  
385 stable hemispheric winter season until August and a gradual decreasing vertical wavelengths towards the end of the year. The  
results at MCM show an even more complicated picture due to the almost vanishing semidiurnal tidal amplitudes. Only during  
the local summer months of January/February and November/December are meaningful vertical wavelengths derivable, with  
vertical wavelengths of about 70-100 km. It is also worth mentioning that the agreement between the zonal and meridional  
wavelengths is remarkable and provides further confidence in the applied ASF technique used for the Reynolds decomposition.


### 390 **3.4 Reynolds stress components**

Gravity waves are an essential driver of the MLT dynamics and variability carrying energy and momentum from their source  
region to the altitude of their deposition. The breaking of GWs can trigger the generation of non-primary GWs, which again  
can propagate upwards (Becker and Vadas, 2018; Vadas and Becker, 2018) causing a complex interaction chain for the GW  
activity and the resulting forcing at the MLT. The acceleration/deceleration of the mean flow due to momentum and energy  
395 transfer by breaking GW can be estimated from the vertical gradient of gravity wave momentum flux (Ern et al., 2011).


From our Reynolds decomposition and the retrieval, we determine three momentum fluxes, which are often referred to as the  
vertical flux of zonal momentum  $\langle uw \rangle$ , the vertical flux of meridional momentum  $\langle vw \rangle$  and the horizontal momentum  
flux  $\langle uv \rangle$ , where the  $\langle \rangle$  denotes temporal averaging.

Figure 8 shows all three momentum flux components as a 30-day median shifted by one day for the year 2019. There are three  
400 groups of panels presenting the vertical flux of zonal momentum (panel a), the vertical flux of meridional momentum (panel b)  
and the horizontal momentum flux (panel c). The stations are sorted according to their latitude within each panel to allow an  
easier comparison between the sites. The results shown in this figure indicate that, for all six meteor radar observations, there is  
a characteristic seasonal pattern with noticeable differences between the different locations. The vertical flux of zonal momen-  
tum  $\langle uw \rangle$  is rather variable with longitude and latitude. Observations at TDF and ROT show some similarities regarding  
405 the seasonal structure. During the local summer both indicate positive zonal momentum fluxes at the altitude of the zonal  
wind reversal. At higher altitudes, above 95-100 km, the zonal momentum flux reverses to negative values. The winter season  
appears to be more variable, which might be related to the minor warming in September 2019 and the wave activity before. To  
the east, at KEP and KSS, positive zonal momentum fluxes at the higher altitudes (95-105 km) are observed throughout the  
year, but a rather different behaviour during the local winter season at the altitudes below. In particular, at KSS a variable zonal  
410 momentum flux is measured that seems to be in better agreement with TDF and ROT results. Further to the South, at DAV and  
MCM, the seasonal behaviour of the zonal momentum flux seems to reflect the features that are already found at KEP, but with  
different magnitudes.

The vertical flux of horizontal momentum presented in Figure 8 exhibits some longitudinal dependence. Observations at TDF  
and KEP show approximately the opposite vertical structure of the meridional momentum flux pointing out that the meridional  
415 drag (acceleration/deceleration) reverses between their longitudes. Furthermore, results from KEP, KSS and ROT show a good  
agreement of the vertical structure and seasonality of the meridional momentum flux throughout the year. Measurements at  
DAV still show some features of the seasonal meridional momentum flux behaviour, but with decreasing magnitude, while at



**Figure 8.** Comparison of vertical flux of zonal and meridional momentum and horizontal momentum flux for each station during the year 2019 for a) zonal momentum flux b) meridional momentum flux c) horizontal momentum flux.



**Figure 9.** Same as Figure 8, but for the a) zonal, b) meridional and c) vertical wind variances.

MCM, the results show once again a seasonal dependency comparable to those obtained at KEP.


420 Only the horizontal momentum flux  $\langle uv \rangle$  shows a similar seasonal behaviour at TDF, KEP, KSS, ROT and DAV with negative values of  $-50 \text{ m}^2/\text{s}^2$  during the seasonal summer and positive values in winter from April to October. The local winter shows more variability and a semiannual structure at some sites, similar to the mean zonal and meridional winds. At MCM this seasonal variation is still visible, but with a much weaker magnitude.

The seasonal dependence Reynolds stress on the main diagonal of the tensor are also investigated. These terms are also  
425 often called zonal, meridional and vertical wind variances. Figure 9 shows all three variances for each station. Note that the color scale for the vertical variances is 5 times smaller compared to the horizontal wind fluctuations. The zonal and meridional variances exhibit a seasonal structure and a rather obvious altitude dependence. The highest variances are observed at the highest altitudes, which is expected considering that the Reynolds stresses are weighted by the atmospheric density, which decreases exponentially with altitude. It is also a common feature for all sites that the meridional velocity variances exceed the  
430 zonal fluctuations. The seasonal behaviour of the zonal and meridional variances at all stations reflects a semi-annual variation showing minimum variances during the equinoxes, when the mean winds are smallest at altitudes below 95-100 km. Above 100 km the seasonal characteristic appears to be less pronounced. The vertical wind variances are the most challenging values to retrieve. Their seasonal behaviour is less obvious. However, the vertical wind variances also indicate increasing values with decreasing density. Results at MCM are exceptional in this respect and the vertical wind variances exceed the values that are  
435 derived at all other stations. At present we can only speculate on the source of these large values. A possibility is that MCM lies underneath the auroral oval and, thus, the altitudes above 90 km are strongly influenced by precipitating particles and associated effects like Joule heating that might trigger stronger vertical variations (Fong et al., 2014).

To gain confidence in our retrieved horizontal wind variances, we performed a test by estimating the GW wind variances of the resolved GWs directly. It is straightforward to derive a gravity wave residual from the hourly observed wind time series  
440 by subtracting mean winds and the diurnal and semidiurnal tide. Thus, we obtain a hourly time series of the GW residuals, which corresponds to the kinetic energy of the resolved GWs with periods longer than 2 hours and horizontal wavelengths of more than 300 km, whereas the wind variances obtained using Hocking (2005) include the GW variances from all temporal and spatial scales. The GW variances from the residuals are shown in Figure A2.

#### 4 Discussion

445 Meteor radar observations of GW momentum fluxes have now been performed for more than a decade (Hocking, 2005; Fritts et al., 2010a; Placke et al., 2011b; Andrioli et al., 2013; de Wit et al., 2014, 2017). However, the results were not always conclusive and often difficult to interpret. Many of these former studies focus on understanding the method and how to optimize the analysis procedure (Fritts et al., 2010a, 2012b; Placke et al., 2011b; Andrioli et al., 2013; Placke et al., 2015a). Although the Reynolds decomposition appears to be straight forward, it can be challenging to do a proper and robust implementation and  
450 successfully separate the mean flow from the GW fluctuations. Fourier based methods often require long averaging windows



**Figure 10.** Monthly mean geostrophic zonal wind for the austral winter 2019 (a, b, c) and summer 2018/19 (d, e, f) each averaged over the altitude range of its wind maximum (40-60 km for winter, 60-80 km for summer). Black dots mark the positions of the radar stations used here. Data are derived from MLS geopotential height data.



in order to get a proper resolution, but do not capture sufficiently the intermittency of the background (see Figure A3). For shorter windows the irregular sampling of meteors in time and altitude again causes deviations from a regular grid and, additionally, data gaps have to be considered when applying wavelet or Fourier techniques. Another complication of the meteor radar momentum fluxes is that there are no ‘ground truth data’ available to validate the measurements. Satellite observations  
455 provide only a total GW momentum flux without directional information obtained from temperature fluctuations after removing atmospheric tides up to wave number 4 assuming a stationary phase behaviour over a couple of days (Ern et al., 2011; Trinh et al., 2018), and thus confidence on the methodology relies on test with synthetic fields such as those presented in Fritts et al. (2010a) and Fritts et al. (2012a).

As we are mainly interested in the GW momentum flux and wind variances, we have to evaluate carefully the presence of  
460 potential error sources in the Reynolds decomposition methodology. In particular, atmospheric tides show a very intermittent behaviour of the amplitude and phase, which causes some issues in the decomposition when long windows (several days/weeks or months) are used. Thus, we performed some tests to optimize the mean flow, tidal and GW decomposition applying the ASF, a 1-day harmonic fit and a 5-day harmonic fit (see appendix A3). The comparison indicates that the Reynolds decomposition tends to be very sensitive to the applied technique impacting the tidal mean flow and the gravity wave variances. Hence, the  
465 derived momentum fluxes and wind variances can be significantly different, even though the same or similar data sets are used. Previous studies used 4-day fits (de Wit et al., 2014) or S-transforms (de Wit et al., 2017) to decompose the time series. In fact, the harmonic fits provide similar results compared to those obtained using Fourier based techniques such as the S-transform (Stockwell et al., 1996) or wavelets (Torrence and Compo, 1998) for the same averaging length.

Besides these technical aspects of the momentum flux and wind variance retrievals, the year 2019 was exceptional in the SH.  
470 The SH winter season was much more variable in August/September compared to previous years at TDF (climatology from 2008-2018) and disturbed with a rare minor sudden stratospheric warming occurring in September (Yamazaki et al., 2020). This variability is reflected in the mean winds and the momentum fluxes, which show noticeable longitudinal and latitudinal differences pointing towards an unstable and wobbling polar vortex. Figure 10 presents monthly mean geostrophic winds from MLS (Matthias and Ern, 2018) averaged over the altitude range 40-60 km for winter months (a, b, c) and 60-80 km for summer  
475 months (d, e, f), the maximum wind region in each season (not shown). During winter the polar vortex is characterized by a strong longitudinal and latitudinal variation. This characteristic of the SH polar vortex is also found in the climatology of MLS for the SH. The strength of the polar vortex appears to be rather different with longitude and month providing significant differences for the vertical propagation of GW and their encounters with critical levels, fostering wave breaking and the emission of non-primary waves due to localised body forces (Vadas and Fritts, 2001; Becker and Vadas, 2018; Vadas et al., 2018; Dong  
480 et al., 2020; Fritts et al., 2020).

Now we compare the observed momentum fluxes with a theoretical study at the SH in the context of non-primary waves Becker and Vadas (2018). The authors simulated GW drag (GWD) and momentum fluxes for the SH winter and show model outputs for a few days, which are qualitatively comparable to our observations. One major effect of the non-primary waves was a significant increase of the zonal and meridional GWD at 60°W and latitudes between 50-70°S. Our meteor radar observations  
485 reflect this increased zonal wave drag for all stations falling into this geographic region (TDF, KSS, ROT) by a considerably

different meridional mean wind pattern during June-August compared to the other stations (KEP, DAV). Further, the model simulations resulted in a weaker effect for the meridional GWD and, thus, the zonal winds are not as much different between the stations. Comparing the observed momentum fluxes between these simulations and our observations, we found a reasonable agreement for KEP and DAV. The model predicts most of the time an eastward zonal momentum flux and a northward meridional momentum flux for the winter months for all longitudes, except for the mountain wave gravity hot spots around 60°W. At this longitude, the model suggest more variable conditions due to the temporal variable mountain wave excitation, which triggers the body forces at the stratosphere, and, thus the secondary or non-primary wave generation (Vadas and Becker, 2018). As a result stations at Argentina (TDF) and the Antarctic Peninsula (KSS, ROT) show sometimes for the winter months southward meridional momentum fluxes as westward zonal momentum fluxes. We also computed the zonal and meridional GWD from the vertical structure of the retrieved momentum fluxes. The KMCM model reached sometimes values of almost 600 m/s/d for the zonal GWD and about 1000 m/s/d for the meridional GWD (Becker and Vadas, 2018). Our observations resulted in similar values on average, which supports that the model and observations are sensitive to a rather similar part of the relevant gravity wave scales with respect to the periods and horizontal wavelenghts. Recent studies on momentum flux spectra using general circulation models suggest that most of the energy of the GW at the MLT is found at periods between 4-12 hours and at horizontal wavelenghts of  $\lambda_h \approx 1000\text{km}$  (Shibuya and Sato, 2019). These scales are covered by our variance and momentum flux retrievals due to the performed Reynolds decomposition and, thus, our results should be representative as integral over the most relevant gravity wave periods and horizontal scales. However, WACCM simulations suggested a leading order contribution of gravity waves between 20-200 km, which are also present in the obtained momentum fluxes. Considering the modelling results from Vadas and Becker (2018), Becker and Vadas (2018) and Shibuya and Sato (2019) as well as our observations of the SH momentum fluxes for all stations, the leading order effect for the momentum budget is given by larger horizontal scale waves and inertia GW periods, which are characteristic properties of non-primary waves. This polar vortex wobbling in 2019 is essentially modifying the mesospheric gravity wave activity and the resulting momentum flux at the altitude of the wave breaking in the mesosphere and above. Characteristics of GWs in the MLT strongly depend on their vertical propagation path and the background wind field along this path, which efficiently alters the amplitude growth of the gravity waves depending on their phase velocity relative to the mean flow. On the other hand, breaking gravity waves deposit momentum on the mean flow and, thus, enhance/weaken/shift the polar vortex, contributing further to the wobbling especially in the mesosphere. For example, the wind maximum in the meteor radar zonal winds (see Fig. 2) was at lower altitudes in June but at upper altitudes in July and August for most of the stations. One explanation of this phenomenon could be that the polar vortex in the upper stratosphere/lower mesosphere (see Fig. 10) was relatively weak in June for most of the stations while it was considerably stronger especially in July and August. This hypothesis is confirmed by the observations obtained at KEP, where stronger winds are observed in the upper stratosphere and lower mesosphere already in June resulting in a higher wind maximum compared to the other stations. Furthermore, there are considerable longitudinal differences of orographic gravity wave sources in the SH resulting already in some asymmetry at stratospheric altitudes.

During the summer months the differences in the zonal wind between the different stations in the mesosphere are much smaller than in winter (see Fig. 10) resulting in smaller differences between the stations in the meteor radar observation window (see

Fig. 2).

In particular, the SH winter season shows remarkable longitudinal and latitudinal changes in the strength of the zonal wind velocities (see Figure 10). Observations with TDF and the KEP radar, although at the same latitude, show that the polar vortex is rather different at each longitude significantly modifying the conditions for vertical GW propagation, and, thus triggering differences in the altitude of the momentum flux deposition. Furthermore, it is likely that both stations experience the effect of different GW sources. ROT and KSS are also reflecting a significant differences in the momentum fluxes, which are partly explainable by the mesospheric zonal wind field, which shows a strong gradient above the Antarctic Peninsula leading to differences of the polar vortex above KSS and ROT.

The most consistent results among the various locations investigated in this study are obtained in the horizontal momentum fluxes  $\langle uv \rangle$  and the wind variances. TDF, KEP, KSS, ROT and DAV observe a very similar seasonal behaviour and only the strength of the flux differs between the sites. This is also the case for the wind variances, which are very consistent between the meteor radars providing confidence in the retrievals. The orography around the meteor radars plays only an indirect role in the observed mesospheric momentum fluxes and wind variances. The observed total flux above the station is the result of all gravity waves that propagate into the mesosphere independent of their origin (e.g., jet instabilities, convection, orography or non-primary waves). Satellite observations of the total momentum flux showed at the stratosphere a significant GW hot spot around the Andes and Antarctic Peninsula (Ern et al., 2011). However, with increasing altitude this momentum flux forms a plume stretching downwind of the Andes and Antarctic Peninsula, and, at the mesosphere/lower thermosphere one finds more or less a longitudinal band of the momentum flux confined to the latitudinal band between 40°-65°S (Trinh et al., 2018). GCM modelling with a GW resolving model indicated that the zonal and meridional momentum flux shows a latitudinal and longitudinal structure (Becker and Vadas, 2018) with multiple sign reversals of the momentum flux within this latitude band. In particular, the signs of the resulting momentum flux are opposite between the Antarctic Peninsula and the Andes around TDF. Furthermore, the momentum fluxes exhibit a reversal of the GW drag between TDF and KEP at the mesosphere, although both locations are at the same latitude.

## 5 Conclusions

This study presents an overview of gravity wave momentum fluxes and wind variances at the MLT in the SH from the mid-latitudes at TDF, Argentina and KEP to the polar latitude of DAV and MCM Antarctic Stations as well as King Sejong Station and ROT at the Antarctic Peninsula for the year 2019. The year 2019 was exceptional and, in particular the hemispheric winter season appears to be more disturbed than previous years resulting in a rare minor stratospheric warming in September.

We performed a detailed analysis of the mean zonal and meridional wind for the year 2019 to explore the longitudinal and latitudinal differences. We noticed significant differences on relatively small regional scales, for instance between KSS and ROT. In addition, we found a strong dependence of the zonal and meridional wind pattern during the SH winter season indicating an asymmetric structure of the polar vortex at the MLT. This asymmetry was verified by MLS geostrophic zonal wind observations at the stratosphere and mesosphere, which revealed longitudinal differences of the intensity as well as an altitude

dependence of the polar vortex leading to temporal and spatially variable filter conditions for the vertical propagation of gravity waves. These results are consistent to the predictions made from model simulations about the secondary or non-primary wave generation due to breaking mountain waves above the gravity hot spots in Argentina and the Antarctic Peninsula.

The derived mean daily wind climatologies for the year 2019 provide convincing evidence that the meridional wind and to a weaker extent the zonal winds are disturbed during the winter season due to orographically generated mountain waves, causing non-primary wave emissions at the stratosphere. This asymmetry in the zonal and meridional winds sustains a polar vortex wobbling in the SH that can be found consistently in the climatologies from MLS and at TDF and DAV. Furthermore, we investigated the diurnal and semidiurnal tidal seasonal variation of the amplitudes and phases for all six stations to assess longitudinal and latitude differences similar to the mean winds. The diurnal tide showed a consistent behaviour of the amplitude and phases measured at TDF, KEP, KSS, ROT and DAV and entirely different seasonal response over MCM. Diurnal tidal phases appeared to be most variable during the local winter season, where the smallest amplitudes are observed. Semidiurnal tides indicated a more complex seasonal structure exhibiting a strong difference in amplitude between the zonal and meridional component for TDF, KEP and ROT, which was not reflected by the semidiurnal tidal phases. In addition, with increasing southern latitudes the amplitude of the semidiurnal tide decreases and shows a different seasonal structure at DAV and basically vanishes over MCM. The seasonal characteristic of vertical wavelengths observed by the meteor radars between 80-100 km for the hemispheric winter season from mid-February to October 2019 at TDF, KEP, KSS and RO is very consistent and takes values of 80-100 km. Moreover, there was a tendency of increased vertical wavelengths during times with very small semidiurnal tidal amplitudes.


Finally, we retrieved the wind variances at all stations. These wind variances exhibit a seasonal behaviour with minimum variances during the equinoxes. In general, the meridional wind variances exceed the zonal components. Besides some differences in the absolute values of the wind variances all observations feature remarkable similarities throughout the year 2019.

*Data availability.* The meteor radar data used in this study from ROT and KEP is from Mitchell, N. (2019): University of Bath Skymet meteor radar data collection. Centre for Environmental Data Analysis, 2020. <https://catalogue.ceda.ac.uk/uuid/836daab8d626442ea9b8d0474125a446>. The DAV radar data are available upon request from Damian Murphy (damian.murphy@aad.gov.au). They are described at [https://data.aad.gov.au/metadata/records/Davis\\_33MHz\\_Meteor\\_Radar](https://data.aad.gov.au/metadata/records/Davis_33MHz_Meteor_Radar). The TDF meteor radar data can be requested from DJ (diego.janches@nasa.gov). The KSS meteor radar data is available on request from YHK (yhkim@cnu.ac.kr).

## Appendix A

### A1 Holographic analysis

The holograms shown in Figure A1 are computed from the semidiurnal tidal phases in the complex domain to account for phase wrapping. As a time dependent phase corresponds to a frequency shift, it is straight forward to estimate the potential Doppler shift of the tidal frequency. Colors towards the red indicate a shift towards lower frequencies, whereas a shift towards the blue



**Figure A1.** Holographic analysis of the semidiurnal tidal variability and lunar elevation angle (thin black line) for each station during the year 2019 for a) TDF, b) KEP, c) King Sejong Station, d) ROT, e) Davis and f) MCM.

585 indicate higher frequencies in analogy to optical Doppler measurements. The method is presented in more detail in Stober et al. (2019). It is evident from this figure that the natural variability of the semidiurnal tide already covers the frequency of the predicted lunar tide during a Pekeris resonance (12.42 hours) (Forbes and Zhang, 2012; Zhang and Forbes, 2014). However, these frequent phase shifts appear to be uncorrelated with the lunar orbit and can occur throughout the year at all meteor radar locations without satisfying the vertical conditions of the winds and temperature fields required for the Pekeris resonance.

## 590 **A2 Gravity wave residual kinetic energies**

The Reynolds decomposition with the ASF can also be used to separate the resolved gravity waves from tides and the daily mean winds. Figure A2 shows the gravity wave residuals after subtracting the daily mean zonal and meridional wind, as well as the diurnal and semidiurnal tides. This gravity wave residual essentially contains the inertia gravity waves with periods up to 16 hours that are not tides. The separation between tide and gravity wave is done by the vertical wavelength.

## 595 **A3 Reynolds decomposition comparison of ASF and harmonic fitting**

The Reynolds decomposition is a very important element in the momentum flux and wind variance measurements. Therefore, we tested different methods to perform the Reynolds decomposition. Figure A3 shows a comparison of the ASF (upper two panels), a harmonic fit with 24 hours window (central panels) and the harmonic fit with 5 day window (lower two panels). The left column always shows the reconstructed time series using a daily mean winds and the diurnal, semidiurnal and terdiurnal  
600 tide. The right column shows the resulting gravity wave residuals. The 2D ASF performs well in recovering a very smooth tidal field and even changes in the tidal phases with altitude. The main difference compared to the 24-hour fit is given in a cleaner removal of a potential inertia gravity wave contamination of the fitted tidal fields due to gravity waves with rather short vertical wavelengths. A 5-day harmonic fit increases the gravity wave residuals significantly, as phase variations of the tide within the long window are no longer sufficiently captured.

605 *Author contributions.* The conceptional idea of this study was developed by GS, DJ and DF in the frame of the SOUTHTRAC campaign. VM and KB substantially helped with the data analysis of MLS and the development of the ASF. JM contributed to the data analysis at MCM. All authors reviewed and edited the manuscript. TMG, NM, DM, DJ, JM, SP, WL and YHK provided meteor radar data.

*Competing interests.* The authors declare that there are no competing interests.

*Acknowledgements.* DJ was supported by by the NASA NASA Heliophysics ISFM Program. TDF's operation is supported by NASA SSO,  
610 NESAC assessment TI-17- 01204, and NSF grant AGS-1647354. For NM and TMG this work was supported by the Natural Environment Research Council [grant numbers: NE/R001391/1, NE/R001235/1]. YHK and WL are financially supported by Korea Polar Research Institute.

Operation of the Davis meteor radar was supported under Australian Antarctic Science project 4445. The authors appreciate the invaluable support of the EARG personnel with the operation of TDF. The momentum flux retrievals were developed as part of the ARISE design study (<http://arise-project.eu>) funded by the European Union's 7th Framework Programme for Research and Techno-logical Development. GS is  
615 a member of the Oeschger Center for Climate Change Research. GS acknowledges the helpful discussions within the DFG research unit MS-GWaves.

## References

- Amante, C. and Eakins, B.: ETOPO1 1 Arc-Minute Global Relief Model: Procedures, Data Sources and Analysis. NOAA Technical Memorandum NESDIS NGDC-24. National Geophysical Data Center, NOAA, <https://doi.org/doi:10.7289/V5C8276M> 2020/09/23, 2009.
- 620 Andrioli, V. F., Fritts, D. C., Batista, P. P., and Clemesha, B. R.: Improved analysis of all-sky meteor radar measurements of gravity wave variances and momentum fluxes, *Annales Geophysicae*, 31, 889–908, <https://doi.org/10.5194/angeo-31-889-2013>, <http://www.ann-geophys.net/31/889/2013/>, 2013.
- Batista, P., Clemesha, B., Tokumoto, A., and Lima, L.: Structure of the mean winds and tides in the meteor region over Cachoeira Paulista, Brazil (22.7°S,45°W) and its comparison with models, *Journal of Atmospheric and Solar-Terrestrial Physics*, 66, 623  
625 – 636, <https://doi.org/https://doi.org/10.1016/j.jastp.2004.01.014>, <http://www.sciencedirect.com/science/article/pii/S1364682604000264>, dynamics and Chemistry of the MLT Region - PSMOS 2002 International Symposium, 2004.
- Baumgarten, K. and Stober, G.: On the evaluation of the phase relation between temperature and wind tides based on ground-based measurements and reanalysis data in the middle atmosphere, *Annales Geophysicae*, 37, 581–602, <https://doi.org/10.5194/angeo-37-581-2019>, <https://www.ann-geophys.net/37/581/2019/>, 2019.
- 630 Baumgarten, K., Gerding, M., and Lübken, F.-J.: Seasonal variation of gravity wave parameters using different filter methods with daylight lidar measurements at midlatitudes, *Journal of Geophysical Research: Atmospheres*, 122, 2683–2695, <https://doi.org/10.1002/2016JD025916>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2016JD025916>, 2017.
- Becker, E.: Dynamical Control of the Middle Atmosphere, *Space Science Reviews*, 168, 283–314, <https://doi.org/10.1007/s11214-011-9841-5>, <http://dx.doi.org/10.1007/s11214-011-9841-5>, 2012.
- 635 Becker, E. and Vadas, S. L.: Secondary Gravity Waves in the Winter Mesosphere: Results From a High-Resolution Global Circulation Model, *Journal of Geophysical Research: Atmospheres*, 123, 2605–2627, <https://doi.org/10.1002/2017JD027460>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2017JD027460>, 2018.
- Beldon, C. L. and Mitchell, N. J.: Gravity wave–tidal interactions in the mesosphere and lower thermosphere over Rothera, Antarctica (68°S, 68°W), *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/10.1029/2009JD013617>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2009JD013617>, 2010.
- 640 Chau, J. L., Hoffmann, P., Pedatella, N. M., Matthias, V., and Stober, G.: Upper mesospheric lunar tides over middle and high latitudes during sudden stratospheric warming events, *Journal of Geophysical Research: Space Physics*, 120, 3084–3096, <https://doi.org/10.1002/2015JA020998>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2015JA020998>, 2015.
- Chu, X., Zhao, J., Lu, X., Harvey, V. L., Jones, R. M., Becker, E., Chen, C., Fong, W., Yu, Z., Roberts, B. R., and Dörnbrack, A.: Lidar Observations of Stratospheric Gravity Waves From 2011 to 2015 at McMurdo (77.84°S, 166.69°E), Antarctica: 2. Potential Energy Densities, Lognormal Distributions, and Seasonal Variations, *Journal of Geophysical Research: Atmospheres*, 123, 7910–7934, <https://doi.org/10.1029/2017JD027386>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2017JD027386>, 2018.
- Conte, J. F., Chau, J. L., Stober, G., Pedatella, N., Maute, A., Hoffmann, P., Janches, D., Fritts, D., and Murphy, D. J.: Climatology of semi-diurnal lunar and solar tides at middle and high latitudes: Interhemispheric comparison, *Journal of Geophysical Research: Space Physics*,  
650 122, 7750–7760, <https://doi.org/10.1002/2017JA024396>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2017JA024396>, 2017.
- Das, U., Ward, W. E., Pan, C. J., and Das, S. K.: Migrating and non-migrating tides observed in the stratosphere from FORMOSAT-3/COSMIC temperature retrievals, *Annales Geophysicae*, 38, 421–435, <https://doi.org/10.5194/angeo-38-421-2020>, <https://angeo.copernicus.org/articles/38/421/2020/>, 2020.



- de Araújo, L. R., Lima, L. M., Batista, P. P., and Jacobi, C.: Behaviour of monthly tides from meteor radar winds at 22.7°S during declining phases of 23 and 24 solar cycles, *Journal of Atmospheric and Solar-Terrestrial Physics*, 205, 105 298, <https://doi.org/https://doi.org/10.1016/j.jastp.2020.105298>, <http://www.sciencedirect.com/science/article/pii/S1364682620301140>, 2020.
- de Wit, R. J., Hibbins, R. E., Espy, P. J., Orsolini, Y. J., Limpasuvan, V., and Kinnison, D. E.: Observations of gravity wave forcing of the mesopause region during the January 2013 major Sudden Stratospheric Warming, *Geophysical Research Letters*, 41, 4745–4752, <https://doi.org/10.1002/2014GL060501>, <http://dx.doi.org/10.1002/2014GL060501>, 2014.
- de Wit, R. J., Janches, D., Fritts, D. C., and Hibbins, R. E.: QBO modulation of the mesopause gravity wave momentum flux over Tierra del Fuego, *Geophysical Research Letters*, 43, 4049–4055, <https://doi.org/10.1002/2016GL068599>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2016GL068599>, 2016.
- de Wit, R. J., Janches, D., Fritts, D. C., Stockwell, R. G., and Coy, L.: Unexpected climatological behavior of MLT gravity wave momentum flux in the lee of the Southern Andes hot spot, *Geophysical Research Letters*, 44, 1182–1191, <https://doi.org/10.1002/2016GL072311>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2016GL072311>, 2017.
- Dong, W., Fritts, D. C., Lund, T. S., Wieland, S. A., and Zhang, S.: Self-Acceleration and Instability of Gravity Wave Packets: 2. Two-Dimensional Packet Propagation, Instability Dynamics, and Transient Flow Responses, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD030 691, <https://doi.org/10.1029/2019JD030691>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019JD030691>, e2019JD030691 2019JD030691, 2020.
- Egito, F., Andrioli, V., and Batista, P.: Vertical winds and momentum fluxes due to equatorial planetary scale waves using all-sky meteor radar over Brazilian region, *Journal of Atmospheric and Solar-Terrestrial Physics*, 149, 108 – 119, <https://doi.org/https://doi.org/10.1016/j.jastp.2016.10.005>, <http://www.sciencedirect.com/science/article/pii/S1364682616303133>, 2016.
- Ehard, B., Kaifler, B., Kaifler, N., and Rapp, M.: Evaluation of methods for gravity wave extraction from middle-atmospheric lidar temperature measurements, *Atmospheric Measurement Techniques*, 8, 4645–4655, <https://doi.org/10.5194/amt-8-4645-2015>, <https://www.atmos-meas-tech.net/8/4645/2015/>, 2015.
- Ern, M., Preusse, P., Gille, J. C., Hepplewhite, C. L., Mlynczak, M. G., Russell III, J. M., and Riese, M.: Implications for atmospheric dynamics derived from global observations of gravity wave momentum flux in stratosphere and mesosphere, *Journal of Geophysical Research: Atmospheres*, 116, <https://doi.org/10.1029/2011JD015821>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2011JD015821>, 2011.
- Fong, W., Lu, X., Chu, X., Fuller-Rowell, T. J., Yu, Z., Roberts, B. R., Chen, C., Gardner, C. S., and McDonald, A. J.: Winter temperature tides from 30 to 110 km at McMurdo (77.8°S, 166.7°E), Antarctica: Lidar observations and comparisons with WAM, *Journal of Geophysical Research: Atmospheres*, 119, 2846–2863, <https://doi.org/10.1002/2013JD020784>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2013JD020784>, 2014.
- Forbes, J. M. and Zhang, X.: Lunar tide amplification during the January 2009 stratosphere warming event: Observations and theory, *Journal of Geophysical Research: Space Physics*, 117, <https://doi.org/10.1029/2012JA017963>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2012JA017963>, 2012.
- Fritts, D. and Alexander, M. J.: Gravity wave dynamics and effects in the middle atmosphere, *Reviews of Geophysics*, 41, 1–64, <https://doi.org/doi:10.1029/2001RG000106>, 2003.
- Fritts, D. C. and Nastrom, G. D.: Sources of Mesoscale Variability of Gravity Waves. Part II: Frontal, Convective, and Jet Stream Excitation, *Journal of the Atmospheric Sciences*, 49, 111–127, [https://doi.org/10.1175/1520-0469\(1992\)049<0111:SOMVOG>2.0.CO;2](https://doi.org/10.1175/1520-0469(1992)049<0111:SOMVOG>2.0.CO;2), [https://doi.org/10.1175/1520-0469\(1992\)049<0111:SOMVOG>2.0.CO;2](https://doi.org/10.1175/1520-0469(1992)049<0111:SOMVOG>2.0.CO;2), 1992.

- Fritts, D. C., Janches, D., and Hocking, W. K.: Southern Argentina Agile Meteor Radar: Initial assessment of gravity wave momentum fluxes, *Journal of Geophysical Research: Atmospheres*, 115, n/a–n/a, <https://doi.org/10.1029/2010JD013891>, <http://dx.doi.org/10.1029/2010JD013891>, d19123, 2010a.
- 695 Fritts, D. C., Janches, D., Iimura, H., Hocking, W. K., Mitchell, N. J., Stockwell, R. G., Fuller, B., Vandeppeer, B., Hormaechea, J., Brunini, C., and Levato, H.: Southern Argentina Agile Meteor Radar: System design and initial measurements of large-scale winds and tides, *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/10.1029/2010JD013850>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2010JD013850>, 2010b.
- Fritts, D. C., Janches, D., Hocking, W. K., Mitchell, N. J., and Taylor, M. J.: Assessment of gravity wave momentum flux measurement capabilities by meteor radars having different transmitter power and antenna configurations, *Journal of Geophysical Research: Atmospheres*, 117, n/a–n/a, <https://doi.org/10.1029/2011JD017174>, <http://dx.doi.org/10.1029/2011JD017174>, d10108, 2012a.
- 700 Fritts, D. C., Janches, D., Iimura, H., Hocking, W. K., Bageston, J. V., and Leme, N. M. P.: Drake Antarctic Agile Meteor Radar first results: Configuration and comparison of mean and tidal wind and gravity wave momentum flux measurements with Southern Argentina Agile Meteor Radar, *Journal of Geophysical Research: Atmospheres*, 117, <https://doi.org/10.1029/2011JD016651>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2011JD016651>, 2012b.
- 705 Fritts, D. C., Dong, W., Lund, T. S., Wieland, S., and Laughman, B.: Self-Acceleration and Instability of Gravity Wave Packets: 3. Three-Dimensional Packet Propagation, Secondary Gravity Waves, Momentum Transport, and Transient Mean Forcing in Tidal Winds, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD030692, <https://doi.org/10.1029/2019JD030692>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019JD030692>, e2019JD030692 2019JD030692, 2020.
- 710 Geller, M. A., Alexander, M. J., Love, P. T., Bacmeister, J., Ern, M., Hertzog, A., Manzini, E., Preusse, P., Scaife, A. A., and Zhou, T.: A Comparison between Gravity Wave Momentum Fluxes in Observations and Climate Models, *Journal of Climate*, 26, 6383–6405, <https://doi.org/10.1175/JCLI-D-12-00545.1>, <https://doi.org/10.1175/JCLI-D-12-00545.1>, 2013.
- Gudadze, N., Stober, G., and Chau, J. L.: Can VHF radars at polar latitudes measure mean vertical winds in the presence of PMSE?, *Atmospheric Chemistry and Physics*, 19, 4485–4497, <https://doi.org/10.5194/acp-19-4485-2019>, <https://www.atmos-chem-phys.net/19/4485/2019/>, 2019.
- 715 Hagan, M. E. and Forbes, J. M.: Migrating and nonmigrating diurnal tides in the middle and upper atmosphere excited by tropospheric latent heat release, *Journal of Geophysical Research: Atmospheres*, 107, ACL 6–1–ACL 6–15, <https://doi.org/10.1029/2001JD001236>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2001JD001236>, 2002.
- Hagan, M. E. and Forbes, J. M.: Migrating and nonmigrating semidiurnal tides in the upper atmosphere excited by tropospheric latent heat release, *Journal of Geophysical Research: Space Physics*, 108, <https://doi.org/10.1029/2002JA009466>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2002JA009466>, 2003.
- 720 He, M., Chau, J. L., Stober, G., Li, G., Ning, B., and Hoffmann, P.: Relations Between Semidiurnal Tidal Variants Through Diagnosing the Zonal Wavenumber Using a Phase Differencing Technique Based on Two Ground-Based Detectors, *Journal of Geophysical Research: Atmospheres*, 123, 4015–4026, <https://doi.org/10.1002/2018JD028400>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2018JD028400>, 2018.
- 725 Heale, C. J., Bossert, K., Vadas, S. L., Hoffmann, L., Dörnbrack, A., Stober, G., Snively, J. B., and Jacobi, C.: Secondary Gravity Waves Generated by Breaking Mountain Waves Over Europe, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD031662, <https://doi.org/10.1029/2019JD031662>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019JD031662>, e2019JD031662 10.1029/2019JD031662, 2020.


- 730 Hocke, K., Hagen, J., Schranz, F., and Bernet, L.: Geographical distributions of mesospheric gravity wave activity before and after major sudden stratospheric warmings observed by Aura/MLS, *Atmospheric Chemistry and Physics Discussions*, 2019, 1–24, <https://doi.org/10.5194/acp-2019-630>, <https://www.atmos-chem-phys-discuss.net/acp-2019-630/>, 2019.
- Hocking, W., Fuller, B., and Vandepeer, B.: Real-time determination of meteor-related parameters utilizing modern digital technology, *Journal of Atmospheric and Solar-Terrestrial Physics*, 63, 155 – 169, [https://doi.org/http://dx.doi.org/10.1016/S1364-6826\(00\)00138-](https://doi.org/http://dx.doi.org/10.1016/S1364-6826(00)00138-3)
- 735 3, <http://www.sciencedirect.com/science/article/pii/S1364682600001383>, radar applications for atmosphere and ionosphere research - PIERS 1999, 2001.
- Hocking, W. K.: A new approach to momentum flux determinations using SKiYMET meteor radars, *Annales Geophysicae*, 23, 2433–2439, <https://doi.org/10.5194/angeo-23-2433-2005>, <http://www.ann-geophys.net/23/2433/2005/>, 2005.
- Hoffmann, P., Becker, E., Singer, W., and Placke, M.: Seasonal variation of mesospheric waves at northern middle and high latitudes, *Journal of Atmospheric and Solar-Terrestrial Physics*, 72, 1068 – 1079, <https://doi.org/https://doi.org/10.1016/j.jastp.2010.07.002>, <http://www.sciencedirect.com/science/article/pii/S1364682610001987>, 2010.
- 740 Holdsworth, D. A., Reid, I. M., and Cervera, M. A.: Buckland Park all-sky interferometric meteor radar, *Radio Science*, 39, n/a–n/a, <https://doi.org/10.1029/2003RS003014>, <http://dx.doi.org/10.1029/2003RS003014>, rS5009, 2004.
- Jackson, D. R., Gadian, A., Hindley, N. P., Hoffmann, L., Hughes, J., King, J., Moffat-Griffin, T., Moss, A. C., Ross, A. N., Vosper, S. B., Wright, C. J., and Mitchell, N. J.: The South Georgia Wave Experiment: A Means for Improved Analysis of Gravity Waves and Low-Level Wind Impacts Generated from Mountainous Islands, *Bulletin of the American Meteorological Society*, 99, 1027–1040, <https://doi.org/10.1175/BAMS-D-16-0151.1>, <https://doi.org/10.1175/BAMS-D-16-0151.1>, 2018.
- 745 Jacobs, E. and Ralston, E. W.: Ambiguity Resolution in Interferometry, *IEEE Transactions on Aerospace and Electronic Systems*, AES-17, 766–780, 1981.
- 750 Janches, D., Palo, S. E., Lau, E. M., Avery, S. K., Avery, J. P., de la Peña, S., and Makarov, N. A.: Diurnal and seasonal variability of the meteoric flux at the South Pole measured with radars, *Geophysical Research Letters*, 31, <https://doi.org/10.1029/2004GL021104>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2004GL021104>, 2004.
- Janches, D., Hocking, W., Pifko, S., Hormaechea, J. L., Fritts, D. C., Brunini, C., Michell, R., and Samara, M.: Interferometric meteor head echo observations using the Southern Argentina Agile Meteor Radar, *J. Geophys. Res. Space Physics*, 119, <https://doi.org/10.1002/2013JA019241>, 2014.
- 755 Janches, D., Bruzzone, J. S., Weryk, R. J., Hormaechea, J. L., Wiegert, P., and Brunini, C.: Observations of an Unexpected Meteor Shower Outburst at High Ecliptic Southern Latitude and Its Potential Origin, *The Astrophysical Journal*, 895, L25, <https://doi.org/10.3847/2041-8213/ab9181>, <https://doi.org/10.3847%2F2041-8213%2Fab9181>, 2020.
- Jones, J., Webster, A. R., and Hocking, W. K.: An improved interferometer design for use with meteor radars, *Radio Science*, 33, 55–65, <https://doi.org/10.1029/97RS03050>, <http://dx.doi.org/10.1029/97RS03050>, 1998.
- 760 Lee, C., Kim, Y. H., Kim, J.-H., Jee, G., Won, Y.-I., and Wu, D. L.: Seasonal variation of wave activities near the mesopause region observed at King Sejong Station (62.22°S, 58.78°W), Antarctica, *Journal of Atmospheric and Solar-Terrestrial Physics*, 105-106, 30 – 38, <https://doi.org/https://doi.org/10.1016/j.jastp.2013.07.006>, <http://www.sciencedirect.com/science/article/pii/S1364682613001971>, 2013.
- 765 Lee, C., Kim, J.-H., Jee, G., Lee, W., and Song, I.-S. H.: New method of estimating temperatures near the mesopause region using meteor radar observations, *Geophysical Research Letters*, 43, 10,580–10,585, <https://doi.org/10.1002/2016GL071082>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2016GL071082>, 2016.

- Lee, W., Kim, Y. H., Lee, C., and Wu, Q.: First Comparison of Mesospheric Winds Measured with a Fabry-Perot Interferometer and Meteor Radar at the King Sejong Station (62.2S, 58.8W), *Journal of Astronomy and Space Sciences*, 35, 235–242, 2018.
- Lindzen, R. S.: Turbulence and stress owing to gravity wave and tidal breakdown, *Journal of Geophysical Research: Oceans*, 86, 9707–9714, 770 <https://doi.org/10.1029/JC086iC10p09707>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/JC086iC10p09707>, 1981.
- Lindzen, R. S. and Chapman, S.: Atmospheric Tides, *Space Sci. Rev.*, 10, 3–188, 1969.
- Liu, G., Janches, D., Lieberman, R. S., Moffat-Griffin, T., Fritts, D. C., and Mitchell, N. J.: Coordinated Observations of 8- and 6-hr Tides in the Mesosphere and Lower Thermosphere by Three Meteor Radars Near 60°S Latitude, *Geophysical Research Letters*, 47, e2019GL086629, <https://doi.org/10.1029/2019GL086629>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019GL086629>, 775 e2019GL086629 2019GL086629, 2020.
- Liu, H.-L.: Quantifying gravity wave forcing using scale invariance, *Nature Communications*, 10, 2019.
- Livesey, N. J., Van Snyder, W., Read, W. G., and Wagner, P. A.: Retrieval algorithms for the EOS Microwave limb sounder (MLS), *IEEE Transactions on Geoscience and Remote Sensing*, 44, 1144–1155, 2006.
- Livesey, N. J., Santee, M. L., and Manney, G. L.: A Match-based approach to the estimation of polar stratospheric ozone loss using Aura 780 Microwave Limb Sounder observations, *Atmospheric Chemistry and Physics*, 15, 9945–9963, <https://doi.org/10.5194/acp-15-9945-2015>, <http://www.atmos-chem-phys.net/15/9945/2015/>, 2015.
- Matthias, V. and Ern, M.: On the origin of the mesospheric quasi-stationary planetary waves in the unusual Arctic winter 2015/2016, *Atmospheric Chemistry and Physics*, 18, 4803–4815, <https://doi.org/10.5194/acp-18-4803-2018>, 2018.
- McCormack, J., Hoppel, K., Kuhl, D., de Wit, R., Stober, G., Espy, P., Baker, N., Brown, P., Fritts, D., Jacobi, C., Janches, D., Mitchell, N., 785 Ruston, B., Swadley, S., Viner, K., Whitcomb, T., and Hibbins, R.: Comparison of mesospheric winds from a high-altitude meteorological analysis system and meteor radar observations during the boreal winters of 2009–2010 and 2012–2013, *Journal of Atmospheric and Solar-Terrestrial Physics*, 154, 132 – 166, <https://doi.org/http://dx.doi.org/10.1016/j.jastp.2016.12.007>, <http://www.sciencedirect.com/science/article/pii/S1364682616303182>, 2017.
- Murphy, D. J., Forbes, J. M., Walterscheid, R. L., Hagan, M. E., Avery, S. K., Aso, T., Fraser, G. J., Fritts, D. C., Jarvis, M. J., McDonald, A. J., 790 Riggan, D. M., Tsutsumi, M., and Vincent, R. A.: A climatology of tides in the Antarctic mesosphere and lower thermosphere, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/10.1029/2005JD006803>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2005JD006803>, 2006.
- Pancheva, D., Mukhtarov, P., Hall, C., Meek, C., Tsutsumi, M., Pedatella, N., Nozawa, S., and Manson, A.: Climatology of the main (24-h and 12-h) tides observed by meteor radars at Svalbard and Tromsø: Comparison with the models CMAM-DAS and WACCM-X, 795 *Journal of Atmospheric and Solar-Terrestrial Physics*, 207, 105 339, <https://doi.org/https://doi.org/10.1016/j.jastp.2020.105339>, <http://www.sciencedirect.com/science/article/pii/S1364682620301528>, 2020.
- Placke, M., Hoffmann, P., Becker, E., Jacobi, C., Singer, W., and Rapp, M.: Gravity wave momentum fluxes in the MLT—Part II: Meteor radar investigations at high and midlatitudes in comparison with modeling studies, *Journal of Atmospheric and Solar-Terrestrial Physics*, 73, 911 – 920, <https://doi.org/https://doi.org/10.1016/j.jastp.2010.05.007>, <http://www.sciencedirect.com/science/article/pii/S1364682610001513>, 800 scientific Results from Networked and Multi-instrument studies based on MST Radar, 2011a.
- Placke, M., Stober, G., and Jacobi, C.: Gravity wave momentum fluxes in the MLT—Part I: Seasonal variation at Collm (51.3°N, 13.0°E), *Journal of Atmospheric and Solar-Terrestrial Physics*, 73, 904 – 910, <https://doi.org/https://doi.org/10.1016/j.jastp.2010.07.012>, <http://www.sciencedirect.com/science/article/pii/S1364682610002087>, scientific Results from Networked and Multi-instrument studies based on MST Radar, 2011b.

- 805 Placke, M., Hoffmann, P., Latteck, R., and Rapp, M.: Gravity wave momentum fluxes from MF and meteor radar measurements in the polar MLT region, *Journal of Geophysical Research: Space Physics*, 120, 736–750, <https://doi.org/10.1002/2014JA020460>, <http://dx.doi.org/10.1002/2014JA020460>, 2014JA020460, 2015a.
- Placke, M., Hoffmann, P., and Rapp, M.: First experimental verification of summertime mesospheric momentum balance based on radar wind measurements at 69° N, *Annales Geophysicae*, 33, 1091–1096, <https://doi.org/10.5194/angeo-33-1091-2015>, <https://angeo.copernicus.org/articles/33/1091/2015/>, 2015b.
- 810 Plougonven, R. and Zhang, F.: Internal gravity waves from atmospheric jets and fronts, *Reviews of Geophysics*, 52, 33–76, <https://doi.org/10.1002/2012RG000419>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2012RG000419>, 2014.
- Pokhotelov, D., Becker, E., Stober, G., and Chau, J. L.: Seasonal variability of atmospheric tides in the mesosphere and lower thermosphere: meteor radar data and simulations, *Annales Geophysicae*, 36, 825–830, <https://doi.org/10.5194/angeo-36-825-2018>, <https://www.ann-geophys.net/36/825/2018/>, 2018.
- 815 Portnyagin, Y. I., Solovjova, T. V., Makarov, N. A., Merzlyakov, E. G., Manson, A. H., Meek, C. E., Hocking, W., Mitchell, N., Pancheva, D., Hoffmann, P., Singer, W., Murayama, Y., Igarashi, K., Forbes, J. M., Palo, S., Hall, C., and Nozawa, S.: Monthly mean climatology of the prevailing winds and tides in the Arctic mesosphere/lower thermosphere, *Annales Geophysicae*, 22, 3395–3410, <https://doi.org/10.5194/angeo-22-3395-2004>, <https://www.ann-geophys.net/22/3395/2004/>, 2004.
- 820 Riggins, D. M., Tsuda, T., and Shinbori, A.: Evaluation of momentum flux with radar, *Journal of Atmospheric and Solar-Terrestrial Physics*, 142, 98 – 107, <https://doi.org/https://doi.org/10.1016/j.jastp.2016.01.013>, <http://www.sciencedirect.com/science/article/pii/S136468261630013X>, 2016.
- Rüfenacht, R., Baumgarten, G., Hildebrand, J., Schranz, F., Matthias, V., Stober, G., Lübken, F.-J., and Kämpfer, N.: Inter-comparison of middle-atmospheric wind in observations and models, *Atmospheric Measurement Techniques*, 11, 1971–1987, <https://doi.org/10.5194/amt-11-1971-2018>, <https://www.atmos-meas-tech.net/11/1971/2018/>, 2018.
- 825 Sandford, D., Mitchell, N., Vincent, R., and Murphy, D.: The lunar tides in the Antarctic mesosphere and lower thermosphere, *Journal of Atmospheric and Solar-Terrestrial Physics*, 69, 2219 – 2237, <https://doi.org/https://doi.org/10.1016/j.jastp.2007.04.010>, <http://www.sciencedirect.com/science/article/pii/S1364682607001137>, vertical Coupling in the Atmosphere/Ionosphere System, 2007.
- Sandford, D. J., Muller, H. G., and Mitchell, N. J.: Observations of lunar tides in the mesosphere and lower thermosphere at Arctic and middle latitudes, *Atmospheric Chemistry and Physics*, 6, 4117–4127, <https://doi.org/10.5194/acp-6-4117-2006>, <https://www.atmos-chem-phys.net/6/4117/2006/>, 2006.
- 830 Sandford, D. J., Beldon, C. L., Hibbins, R. E., and Mitchell, N. J.: Dynamics of the Antarctic and Arctic mesosphere and lower thermosphere – Part 1: Mean winds, *Atmospheric Chemistry and Physics*, 10, 10 273–10 289, <https://doi.org/10.5194/acp-10-10273-2010>, <https://www.atmos-chem-phys.net/10/10273/2010/>, 2010.
- 835 Schult, C., Stober, G., Janches, D., and Chau, J. L.: Results of the first continuous meteor head echo survey at polar latitudes, *Icarus*, 297, 1 – 13, <https://doi.org/https://doi.org/10.1016/j.icarus.2017.06.019>, <http://www.sciencedirect.com/science/article/pii/S0019103516303761>, 2017.
- Schwartz, M., Lambert, A., Manney, G., Read, W., Livesey, N., Froidevaux, L., Ao, C., Bernath, P., Boone, C., Cofield, R., Daffer, W., Drouin, B., Fetzer, E., Fuller, R., Jarnot, R., Jiang, J., Jiang, Y., Knosp, B. W., Krüger, K., Li, J.-L., Mlyneczek, M., Pawson, S., Russell, J., Santee, M., Snyder, W., Stek, P., Thurstans, R., Tompkins, A., Wagner, P., Walker, K., Waters, J. W., and Wu, D. L.: Validation of the Aura Microwave Limb Sounder temperature and geopotential height measurements, *Journal of Geophysical Research: Atmospheres*, 113, <https://doi.org/10.1029/2007JD008783>, <http://dx.doi.org/10.1029/2007JD008783>, d15S11, 2008.
- 840

- Shibuya, R. and Sato, K.: A study of the dynamical characteristics of inertia-gravity waves in the Antarctic mesosphere combining the PANSY radar and a non-hydrostatic general circulation model, *Atmospheric Chemistry and Physics*, 19, 3395–3415, <https://doi.org/10.5194/acp-19-3395-2019>, <https://acp.copernicus.org/articles/19/3395/2019/>, 2019.
- 845 Siddiqui, T. A., Yamazaki, Y., Stolle, C., Lühr, H., Matzka, J., Maute, A., and Pedatella, N.: Dependence of Lunar Tide of the Equatorial Electrojet on the Wintertime Polar Vortex, Solar Flux, and QBO, *Geophysical Research Letters*, 45, 3801–3810, <https://doi.org/10.1029/2018GL077510>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2018GL077510>, 2018.
- Spargo, A. J., Reid, I. M., and MacKinnon, A. D.: Multistatic meteor radar observations of gravity-wave–tidal interaction over southern Australia, *Atmospheric Measurement Techniques*, 12, 4791–4812, <https://doi.org/10.5194/amt-12-4791-2019>, <https://amt.copernicus.org/articles/12/4791/2019/>, 2019.
- 850 Stober, G., Matthias, V., Jacobi, C., Wilhelm, S., Höffner, J., and Chau, J. L.: Exceptionally strong summer-like zonal wind reversal in the upper mesosphere during winter 2015/16, *Annales Geophysicae*, 35, 711–720, <https://doi.org/10.5194/angeo-35-711-2017>, <https://www.ann-geophys.net/35/711/2017/>, 2017.
- 855 Stober, G., Chau, J. L., Vierinen, J., Jacobi, C., and Wilhelm, S.: Retrieving horizontally resolved wind fields using multi-static meteor radar observations, *Atmospheric Measurement Techniques*, 11, 4891–4907, <https://doi.org/10.5194/amt-11-4891-2018>, <https://www.atmos-meas-tech.net/11/4891/2018/>, 2018a.
- Stober, G., Sommer, S., Schult, C., Latteck, R., and Chau, J. L.: Observation of Kelvin–Helmholtz instabilities and gravity waves in the summer mesopause above Andenes in Northern Norway, *Atmospheric Chemistry and Physics*, 18, 6721–6732, <https://doi.org/10.5194/acp-18-6721-2018>, <https://www.atmos-chem-phys.net/18/6721/2018/>, 2018b.
- 860 Stober, G., Baumgarten, K., McCormack, J. P., Brown, P., and Czarnecki, J.: Comparative study between ground-based observations and NAVGEM-HA reanalysis data in the MLT region, *Atmospheric Chemistry and Physics Discussions*, 2019, 1–37, <https://doi.org/10.5194/acp-2019-1006>, <https://www.atmos-chem-phys-discuss.net/acp-2019-1006/>, 2019.
- Stockwell, R. G., Mansinha, L., and Lowe, R. P.: Localization of the complex spectrum: the S transform, *IEEE Transactions on Signal Processing*, 44, 998–1001, <https://doi.org/10.1109/78.492555>, 1996.
- Torrence, C. and Compo, G. P.: A Practical Guide to Wavelet Analysis, *Bull. Amer. Meteor. Soc.*, 79, 61–78, 1998.
- Trinh, Q. T., Ern, M., Doornbos, E., Preusse, P., and Riese, M.: Satellite observations of middle atmosphere–thermosphere vertical coupling by gravity waves, *Annales Geophysicae*, 36, 425–444, <https://doi.org/10.5194/angeo-36-425-2018>, <https://angeo.copernicus.org/articles/36/425/2018/>, 2018.
- 870 Vadas, S. L. and Becker, E.: Numerical Modeling of the Excitation, Propagation, and Dissipation of Primary and Secondary Gravity Waves during Wintertime at McMurdo Station in the Antarctic, *Journal of Geophysical Research: Atmospheres*, 123, 9326–9369, <https://doi.org/10.1029/2017JD027974>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2017JD027974>, 2018.
- Vadas, S. L. and Fritts, D. C.: Gravity Wave Radiation and Mean Responses to Local Body Forces in the Atmosphere, *Journal of the Atmospheric Sciences*, 58, 2249–2279, [https://doi.org/10.1175/1520-0469\(2001\)058<2249:GWRAMR>2.0.CO;2](https://doi.org/10.1175/1520-0469(2001)058<2249:GWRAMR>2.0.CO;2), 2001.
- 875 Vadas, S. L., Zhao, J., Chu, X., and Becker, E.: The Excitation of Secondary Gravity Waves From Local Body Forces: Theory and Observation, *Journal of Geophysical Research: Atmospheres*, 123, 9296–9325, <https://doi.org/10.1029/2017JD027970>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2017JD027970>, 2018.
- Vierinen, J., Chau, J. L., Charuvil, H., Urco, J. M., Clahsen, M., Avsarkisov, V., Marino, R., and Volz, R.: Observing Mesospheric Turbulence With Specular Meteor Radars: A Novel Method for Estimating Second-Order Statistics of Wind Velocity, *Earth and Space Science*, 6, 1171–1195, <https://doi.org/10.1029/2019EA000570>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019EA000570>, 2019.
- 880


- Vincent, R. A. and Fritts, D. C.: A climatology of gravity wave motions in the mesopause region at Adelaide, Australia, *J. Atmos. Sci.*, 44, 748–760, 1987.
- Wang, H., Boyd, J. P., and Akmaev, R. A.: On computation of Hough functions, *Geoscientific Model Development*, 9, 1477–1488, <https://doi.org/10.5194/gmd-9-1477-2016>, <https://gmd.copernicus.org/articles/9/1477/2016/>, 2016.
- 885 Ward, W. E., Oberheide, J., Goncharenko, L. P., Nakamura, T., Hoffmann, P., Singer, W., Chang, L. C., Du, J., Wang, D.-Y., Batista, P., Clemesha, B., Manson, A. H., Riggini, D. M., She, C.-Y., Tsuda, T., and Yuan, T.: On the consistency of model, ground-based, and satellite observations of tidal signatures: Initial results from the CAWSES tidal campaigns, *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/10.1029/2009JD012593>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2009JD012593>, 2010.
- Waters, J. W., Froidevaux, L., Harwood, R. S., Jarnot, R. F., Pickett, H. M., Read, W. G., Siegel, P. H., Cofield, R. E., Filipiak, M. J., Flower, D. A., Holden, J. R., Lau, G. K., Livesey, N. J., Manney, G. L., Pumphrey, H. C., Santee, M. L., Wu, D. L., Cuddy, D. T., Lay, R. R., Loo, M. S., Perun, V. S., Schwartz, M. J., Stek, P. C., Thurstans, R. P., Boyles, M. A., Chandra, K. M., Chavez, M. C., Chen, G.-S., Chudasama, B. V., Dodge, R., Fuller, R. A., Girard, M. A., Jiang, J. H., Jiang, Y., Knosp, B. W., LaBelle, R. C., Lam, J. C., Lee, K. A., Miller, D., Oswald, J. E., Patel, N. C., Pukala, D. M., Quintero, O., Scaff, D. M., Snyder, W. V., Tope, M. C., Wagner, P. A., and Walch, M. J.: The Earth observing system microwave limb sounder (EOS MLS) on the aura Satellite, *IEEE Transactions on Geoscience and Remote*  
890 *Sensing*, 44, 1075–1092, <https://doi.org/10.1109/TGRS.2006.873771>, 2006.
- Wilhelm, S., Stober, G., and Chau, J. L.: A comparison of 11-year mesospheric and lower thermospheric winds determined by meteor and MF radar at 69° N, *Annales Geophysicae*, 35, 893–906, <https://doi.org/10.5194/angeo-35-893-2017>, <https://www.ann-geophys.net/35/893/2017/>, 2017.
- Wilhelm, S., Stober, G., and Brown, P.: Climatologies and long-term changes in mesospheric wind and wave measurements based on radar  
900 observations at high and mid latitudes, *Annales Geophysicae*, 37, 851–875, <https://doi.org/10.5194/angeo-37-851-2019>, <https://www.ann-geophys.net/37/851/2019/>, 2019.
- Wu, Q., Ward, W., Kristoffersen, S., Maute, A., and Liu, J.: Simulation and Observation of Lunar Tide Effect on High-Latitude, Mesospheric and Lower Thermospheric Winds During the 2013 Sudden Stratospheric Warming Event, *Journal of Geophysical Research: Space Physics*, 124, 1283–1291, <https://doi.org/10.1029/2018JA025476>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2018JA025476>, 2019.
- 905 Yamazaki, Y., Matthias, V., Miyoshi, Y., Stolle, C., Siddiqui, T., Kervalishvili, G., Laštovička, J., Kozubek, M., Ward, W., Themens, D. R., Kristoffersen, S., and Alken, P.: September 2019 Antarctic Sudden Stratospheric Warming: Quasi-6-Day Wave Burst and Ionospheric Effects, *Geophysical Research Letters*, 47, e2019GL086577, <https://doi.org/10.1029/2019GL086577>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019GL086577>, e2019GL086577 10.1029/2019GL086577, 2020.
- Zhang, X. and Forbes, J. M.: Lunar tide in the thermosphere and weakening of the northern polar vortex, *Geophysical Research Letters*, 41, 8201–8207, <https://doi.org/10.1002/2014GL062103>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2014GL062103>, 2014.  
910



**Figure A2.** Kinetic gravity wave energy as determined for the resolved waves from the ASF for each station during the year 2019. Same as Figure 2.



## ASF decomposition (2D)



**Figure A3.** Comparison of different Reynolds decomposition for 8 days at TDF (RioGrande) to underline the sensitivity on the applied method to derive the gravity wave fluctuations. Panel a) shows the ASF reconstructed daily mean plus diurnal, semidiurnal and terdiurnal tide time series. Panel b) presents the gravity wave residuals. Panels c) and d) show the same, but for a 24-hour harmonic fit and panels e) and f) for a 5 day harmonic fit.