

## ***Interactive comment on “Testing the Electrodynamic Method to Derive Height-Integrated Ionospheric Conductances” by Daniel Weimer and Thom Edwards***

**Daniel Weimer and Thom Edwards**

dweimer@vt.edu

Received and published: 20 October 2020

Referee #1:

The paper presents a model of height integrated ionospheric conductance that is derived from results of other dedicated empirical models. The manuscript is well written, in most part clear and understandable and potentially suitable for publication at Annales Geophysicae.

There are few comments, that the authors shall addresses:

1. The model is a composite of different empirical models. Quantify the uncertainty

C1

and/or error that accumulates by using the empirical models to build another one.

---

Response:

The results are presented as maps that are calculated for various combinations of preset conditions, such as IMF magnitude, clock angle, and dipole tilt angle. However, the results were never directly referred to as a "model." We took care to not use that word to describe the results. Our view is that an empirical model takes any random combination of input parameters and outputs a result for specified locations and time. We have not created any such program, although the results could, in theory, be used to look-up a map file that most closely matches the desired conditions, as mentioned in response #3. Many more maps of the results are contained in the data archive than presented in the paper. As discussed in the response to item #2, we cannot obtain a good measurement of the uncertainty for each model. However, in comparison with existing conductivity models, our results are known to be within the the expected range of values, so the errors cannot be very great. More recently there have been two papers in the accepted or pre-print stage that have become available to us only after the submission of our manuscript. Our results are within range of the maps shown in these new papers, which provides confidence that our errors are not nearly as large as would be obtained by accumulating the uncertainties (in the range of 20%-50%) that result purely from the high-frequency fluctuations. Carrying out an accumulation of errors with uncertain input values which would be an exercise in futility, particularly due to the unknown influence of the uncertainties in the IMF values, which is discussed in the next point.

Citations of these new papers will be included in the revised paper:

Carter, J. A., S. E. Milan, L. J. Paxton, B. J. Anderson, and J. Gjerloev, Height-integrated ionospheric conductances parameterised by interplanetary magnetic field and substorm phase, J. Geophys. Res., doi:10.1029/2020JA028121, 2020.

C2

Mukhopadhyay, Agnit, Daniel T. Welling, Michael W. Liemohn, Aaron J. Ridley, Shibaji Chakraborty, and Brian J. Anderson, Conductance Model for Extreme Events: Impact of Auroral Conductance on Space Weather Forecasts, Space Weather, <https://arxiv.org/pdf/2008.12276.pdf>, 2020.

Liemohn, Michael W., The case for improving the Robinson formulas, J. Geophys. Res., doi:10.1029/2020JA028332, 2020.

---

Referee #1:

2. This point relates to point 1. The Discussion mentioned a row of error that might be in the model results. Certainly, the small-scale variation cannot be captured, but this is not what would be expected from the empirical model and is quite clear. Also, as the major source of error, an uncertain solar wind measurement is mentioned. However, it needs to be checked how this is valid and that uncertainties from point 1, may not be more important.

---

Response:

We need to make it more clear in a revision that "small-scale variation" includes high-frequency, temporary fluctuations, and such variations prevent an accurate measurement of the standard deviation in a global-scale, spatial mapping. As given in our response to Referee #2, the empirical models all use measurements that were taken as a function of time, while the satellites were moving in space, or in the case of the ground-based magnetometers, fixed in position while moving through local time. The original measurements contain fluctuations at various frequencies, with the most rapid fluctuations, lasting a few seconds or so, often having the largest magnitudes. The end-result models are global-scale, spatial representations of the fields that do not have the temporal fluctuations included. During the fitting process the difference between the

C3

models and input values (the total square error) is minimized, and this error can be calculated. This is where the 20%-50% numbers come from. But since the high-frequency fluctuations are believed to be mostly temporal variations, rather than spatial, the least-square error that is calculated is not a true representation of the accuracy of the spatial models.

In the revision we will add a more detailed discussion about the differences between the IMF values that are measured by sentinels at the L1 orbit, and what impacts the Earth's magnetosphere. The issue is illustrated well in the 2018 paper by Borovsky that was already cited, that this referee is invited to inspect. Further evidence is given in a paper that was published online only recently, having a key point that "the solar wind measured by a single spacecraft at L1 often does not impact Earth in a homogeneous manner":

Burkholder, B. L., Nykyri, K., & Ma, X. (2020). Use of the L1 Constellation as a Multi-spacecraft Solar Wind Monitor. *Journal of Geophysical Research: Space Physics*, 125, e2020JA027978. <https://doi.org/10.1029/2020JA027978>

As the submitted manuscript already indicated, the full, three-dimensional structure of the IMF cannot be determined unless additional sentinels are placed in orbit around L1. On the topic of the variable time delays we are adding these two references in response to Referee #2:

Weimer, D. R., D. M. Ober, N. C. Maynard, W. J. Burke, M. R. Collier, D. J. McComas, T. Nagai, and C. W. Smith. Variable time delays in the propagation of the interplanetary magnetic field. *J. Geophys. Res.*, 107((A8)), 2002.

Weimer, D. R., and J. H. King. Improved calculations of interplanetary magnetic field phase front angles and propagation time delays. *J. Geophys. Res.*, 113, 2008.

---

Referee #1:

C4

3. The conclusion repeats earlier finding from other authors and mention the shortcomings of the model results submitted here, but that the results might be of some value to the community. This again calls for a quantification, if rough, of the errors expected. To which application the value is expected?

---

Response:

The results could be used in AMIE-like assimilation models and numerical simulations of the coupled magnetosphere-ionosphere system. As we have not created an actual empirical model, the results would need to be used by plugging in our derived values for the IMF conditions that most closely match those being simulated, with interpolation and filling in of missing results. As done by Mukhopadhyay et al [2020], validation of the accuracy is obtained by testing whether or not there is an improvement in various metrics and skill scores in the predictions of ground-level magnetic perturbations. This type of test would be the best way to validate the results. Another possible test would be to feed the output of a FAC model or AMPERE measurements into a potential solver that uses the derived conductivity values, and compare the output with the electric potential patterns that are expected.

---

Referee #1:

4. L 23 says that earlier formulas are confusing and that the authors applied a simpler formula. What does justify the simplification?

---

Response:

Several versions of the conductivity formulas have been published, as cited in the paper. These mathematical equations contain different notations for the same calculations,

C5

but with identical results. Some formulas contain symbols for cyclotron frequencies, others use mobility coefficients, and some may explicitly include all of the terms, such as magnetic field strength, that go into the calculation of such quantities. In some texts the summation over different ion species is only implied. It is the differences between the various formulas, and their variations, that can be confusing, not the formulas themselves; the paper revision will need to state that more clearly. Equations (1) through (4) get to the same result as the others, but explicitly shows the summation over the different ion species; the simplification is just that (3) and (4) are used to simply formulas (1) and (2) by replacing multiple symbols with these two.

---

Referee #1:

5. I had sometimes the impression that not the most recent developments of certain areas are referred to, such as for AMIE, SECS method, or NRLMSIS. Please check about newer developments in these fields.

---

Response:

There have been too many variations of the AMIE and SECS methods to keep track of or cite them all, so the earliest or the most well-known papers are cited. The more recent NRLMSIS model, named simply NRLMSIS 2.0, is described in a paper that was only recently submitted and accepted (17 September 2020) so it could not be included in the first draft of this paper, but it can be referenced in the revision.

---

Referee #1:

6. What is the opinion of the authors if direct observations near the E region would enhance their findings, such as is provided by a satellite mission like Daedalus?

C6

(<https://daedalus.earth>)

---

Response:

Daedalus would provide the opportunity to study more direct impacts of plasma flows in the E region, as the referee suggests. For example, this data would likely be interesting as a follow-up to an earlier work on the relationship between FAC and solar index work, particularly the FAC response to the solar indices representing the extreme ultraviolet emissions [Edwards, T. R., D. R. Weimer, W. K. Tobiska, and N. Olsen, Field-aligned current response to solar indices, *J. Geophys. Res. Space Physics*, 122:5798–5815, 2017]. While the Daedalus data would not have a direct impact on our conductivity calculations, it is possible that these data could confirm the existence of features shown in our Figures 6 through 9, and the influence that the IMF clock angle has on the conductivity distribution. Daedalus observations may perhaps be useful in future version of the IRI and/or NRLMSIS models.

---

Interactive comment on Ann. Geophys. Discuss., <https://doi.org/10.5194/angeo-2020-60>, 2020.



## ***Interactive comment on “Testing the Electrodynamic Method to Derive Height-Integrated Ionospheric Conductances” by Daniel Weimer and Thom Edwards***

**Daniel Weimer and Thom Edwards**

dweimer@vt.edu

Received and published: 20 October 2020

Referee #2:

Methodology 1. While the technical approach is largely (see below) well explained the methodology is questionable. When the authors organize their data by IMF, dipole tilt angle, F10.7 and so on the assumption is that these parameters are indeed controlling the ionospheric electrodynamics. There is a causality implied in the approach which opens two fundamental questions: is there a cause-and-effect and if so what is the delay?. I think this may work on the dayside where delays between SW conditions and the response of the ionospheric electrodynamics are likely on the order of minutes

C1

and thus within the 5-min binning. I, however, object to the approach on the nightside. Here the plasma-sheet is involved and the FACs, particle precipitation, convection and thus conductances and ionospheric currents are due to processes within the plasma sheet. These are only probabilistic related to the SW conditions and the delays are largely unknown and certainly much larger than the 5-min binning. An example is in L338 “IMF clock angles influences the conductivity values”. I do not think the word ‘influences’ is right. Another example is in L203. Is the previous 20 used or is it the center of the window? If centered the ionospheric conditions can hardly be associated with future SW conditions? A discussion of this methodology and arguments for why this is appropriate is required.

---

Response:

This reviewer is correct that the dayside and nightside have different response times after the IMF changes, and that there are uncertainties in the delays. Apparently a revision to the paper needs to be made to clarify that the IMF parameters are averaged over a 20-minute time window, not 5 minutes. This averaging accounts for the fact that the magnetosphere and ionosphere as a whole system requires time to reconfigure after a change in the IMF, and that the nightside takes longer than the dayside. An entire section and several figures in the Weimer et al. [2010] paper is devoted to this topic of variable delays and reconfiguration times, as well as a discussion of the various and sometimes conflicting numbers reported in other papers. Complicating the subject is the fact that there are variable time delays between the measurement of the IMF at the L1 orbit and the time it reaches the Earth [Weimer et al, 2002]. Uncertainty in these delays are reduced by using the technique reported by Weimer and King [2008], to propagate the IMF to the Earth’s bow shock. Weimer et al. [2010] had reported on the results of correlations between the ground-level magnetic field response and a function of the IMF (after applying this propagation). Due to the variability in the system it was found that the dayside response was slightly shorter than the nightside

C2

response, but the peaks in the correlation vs. delay curves were very broad. For the purpose of constructing global patterns using spherical harmonics, it was decided that the best compromise was to use a 25-min average of the IMF, after including a 20-min time lag from the computed time of impact at the bow shock. In later papers the averaging period was reduced to 20 minutes. The computation is done so that each measurement (such as magnetic or electric field) is compared with the average of the IMF that had interacted with the magnetosphere over the previous 20-min time period, not a time-centered average. Future solar wind conditions are never used. In order to make the size of the processed IMF database more manageable, the sliding averages were sampled at 5-min intervals, rather than something like 16 seconds. These time-tagged IMF values are assigned to the data samples taken at the times falling within each window, for use in the fitting process.

On the night side the processes mentioned by the reviewer add even more uncertainty, particularly due to the irregular and unpredictable occurrence of substorms. Nevertheless, in all previous modeling work, including that done by several other authors with different data sets, such as SuperDARN, DMSP, AMPERE, and Cluster) it has been found that the ionospheric electric fields and currents have consistent and repeatable patterns in response to the magnitude and orientation of the IMF and dipole tilt angle. Additional references about these consistent patterns could be added to the revision. The 2017 study by Edwards et al. had found that the solar indices do have a correlations with the magnitude of the field-aligned currents, so it was logical to include the F10.7 index that was used with the ground-level magnetic perturbation model. As the existing empirical models of ionospheric conductivity rely only on activity indices while ignoring the IMF, it seemed that this study needed to be done.

In the revision we will work to find an alternative wording for "IMF clock angles influences the conductivity values". We will also add additional references on the topic of propagation delays:

Weimer, D. R., D. M. Ober, N. C. Maynard, W. J. Burke, M. R. Collier, D. J. McComas,  
C3

T. Nagai, and C. W. Smith. Variable time delays in the propagation of the interplanetary magnetic field. *J. Geophys. Res.*, 107((A8)), 2002.

Weimer, D. R., and J. H. King. Improved calculations of interplanetary magnetic field phase front angles and propagation time delays. *J. Geophys. Res.*, 113, 2008.

---

Referee #2:

1) Methodology 2. I worry that the paper uses a circular argument. The SuperDARN convection distributions use a fill-in model which is IMF driven. Thus, the convection has embedded an IMF dependence which will obviously will be seen in the derived distributions.

---

Response:

"SuperDARN" was mentioned only in the context of the earlier results obtained by Green et al [2007]. This study used an electric potential model derived from the Dynamics Explorer-2 and Swarm satellites, (no SuperDARN data are used) along with the IMF measurements. As the intent is to determine how the IMF influences the resulting conductivity mappings, we do not understand the comment about a circular argument.

---

Referee #2:

2) Technique 1. Please provide an argument for why AMPERE and SuperMAG was not used. These have been carefully validated and provide more comprehensive datasets than those used here.

---

Response:

The publications by Weimer et al. [2010] and Weimer [2013] were the results of a project that had started in late 2008, before there were any data available from the SuperMAG web site. So the PI had no other choice but to independently collect and process the magnetometer data from multiple sources, as described in the 2010 paper. After the 2013 magnetic perturbation model was completed there was never any reason or funding to do the same thing all over again with data from SuperMAG. The way that the SuperMAG data are obtained through their web site also makes it nearly impossible to obtain all of the highest resolution data from every Northern hemisphere magnetometer over an eight-year period.

We disagree that the AMPERE dataset is better for this purpose. The magnetometers on the Iridium satellites actually have a very poor resolution and low sampling rate, and their attitude pointing accuracy isn't that great either. The Oersted, CHAMP, and Swarm satellites all have much better magnetometers and attitude control. AMPERE is better suited for event studies, rather than building statistical models, although there are FAC models that have been created from the AMPERE data that strongly resemble ours. The Edwards et al [2020] paper actually includes a comparison of the new FAC model with some AMPERE results, along with a different FAC model.

---

Referee #2:

3) Technique 2. I worry that the ground level perturbations are assumed to be due to an overhead current. This is not the case in the polar cap and equatorward of the auroral oval. Please comment/discuss. Also see L254-255.

---

Response:

The formulas used to obtain the ionospheric equivalent current functions, described by Chapman and Bartels (1940), Haines (1988), and Haines and Torta (1994) do not

C5

assume that the magnetic perturbations are due only to currents that are directly overhead. The effect of all currents are included in the current function at each point, unlike other methods that simply rotate magnetometer measurements by 90 degrees and interpolate. As mentioned in the Weimer [2019] paper, the method also has the undesired side effect of including the effects of magnetospheric and field-aligned currents, which is what the text in L254-255 refers to. The magnetic fields from these external currents produces unrealistic equivalent currents at very low latitudes, often in a direction that is opposite to what is expected from the electric potential model. As indicated in the paper, the conductivity cannot be calculated in these regions.

---

Referee #2:

4) Technique 3. Clearly the solutions are problematic in regions of low E but I also worry about regions in which the direction of E varies sharply. Maintaining the relative alignment of the various parameters in such regions is very difficult and mis-alignments result in erroneous conductances since these essentially are fudge-factors.

---

Response:

No fudge-factors are used in any attempt to align the models, if that is what the referee meant. The original model outputs are used as is. As the results are flagged as invalid in the areas where the electric field is low and where the conductivity is calculated as negative, there shouldn't be a problem.

---

Referee #2:

5) Technique 4. The authors allude in L435 to the Hall vs DF. Can you include a brief discussion on this widely used assumption? Also see L85-86.

C6

---

Response:

We had comments from a theoretician on a prior version of the paper that the use of the Jdf and Jcf current systems is only a mathematical device, so this sentence was added as an attempt to respond to this remark. But since there is no reference that can be quoted here, we think it is best to just remove this comment in the revision. As the referenced paper by Vanhamäki et al. [2012], and others, have detailed discussions of the Jdf and Jcf components, we don't think that there is much that can be added.

---

Referee #2

6) Energy transfer (e.g. L371). How is energy transported over these long distances? L359 states: "their gradients act to transport energy flux" but I don't understand how this actually works.

---

Response:

Line 359 is referring to results in the paper by Vanhamäki et al. [2012]. The energy transfer actually takes place through a horizontal Poynting flux. Paragraph 23 in this paper says "Between these areas the Poynting flux is transported horizontally near the ionospheric plane" and Paragraph 43 says "conductance gradients give rise to divergence-free Pedersen currents that modify the spatial distribution (but not the total amount) of Joule heating and to curl-free Hall currents that modify the spatial distribution (but not the total amount) of vertical Poynting flux."

---

Referee #2:

C7

7) L437-455. Please change from 'scale sizes' to something like 'small- and meso-scale'. Also, I don't know where the 20%-50% comes from. The contribution from these processes is very much a topic of discussion.

---

Response:

The text already uses "small-scale," and we can't find any occurrence of 'scale sizes'.

The empirical models all use measurements that were taken as a function of time, while the satellites were moving in space, or in the case of the ground-based magnetometers, fixed in position while moving through local time. The original measurements contain fluctuations at various frequencies, with the most rapid fluctuations, lasting a few seconds or so, often having the largest magnitudes. The end-result models are global-scale, spatial representations of the fields that do not have the rapid fluctuations included. During the fitting process the difference between the models and input values (the total square error) is minimized, and this error can be calculated. This is where the 20%-50% numbers come from. But since the high-frequency fluctuations are believed to be mostly temporal variations, rather than spatial, the least-square error that is calculated is not a true representation of the accuracy of the spatial models.

The referee comments indicate that we need to explain this part better in the revision.

---

Referee #2:

Remove L75-77. I disagree, considering all the other complications.

---

Response:

These lines will be rewritten to change the wording of this phrase, which in retrospect is

C8

too harsh: "A larger problem is that models based on activity indices have only marginal utility." The point we were trying to get across is that the community is moving away from "index based" model input and toward real data for a reason, particularly because indices have long lag times before they become available. The need for conductivity mappings that are based on incoming IMF/solar wind values, rather than activity indices, is a major focus point of this paper, and our results show the orientation of the IMF is an important parameter.

---

Referee #2:

L150. "after accounting for solar wind propagation delays". Please elaborate. Which technique is used and I assume this is from L1.

---

Response:

The technique used is described in this publication: Weimer, D. R. and J. H. King, Improved calculations of interplanetary magnetic field phase front angles and propagation time delays, *J. Geophys. Res.*, 113, 2008. We will add the reference in the revision. This technique was referenced in all of the original, empirical model papers.

---

Referee #2:

L293. "a wrong result again, but useful to include." Can you please add an argument "a wrong result again, but useful to include, because. . ."

---

Response:

This revision will be included: "... a wrong result again, but useful to include as it shows

C9

why the divergence-free component alone is not sufficient for calculating the Hall conductivity, as might be assumed. Showing these calculations are useful for understanding the contributions that both components have on the final numerical results."

---

Referee #2:

L319. The Green et al paper was a proof of concept or technical paper.

---

Response:

That is absolutely true, and this proof of concept paper had served as the inspiration for trying the technique using our models, for a wide range of conditions. While we had used the word "demonstrated," to describe the Green et al. paper, we will add "proof of concept" to the revision. While it was only a proof of concept, we still think it is useful to include the comparison in L319.

---

Referee #2:

L339: See Ohtani, S., Gjerloev, J. W., Johnsen, M. G., Yamauchi, M., Brändström, U., & Lewis, A. M. (2019). Solar illumination dependence of the auroral electrojet intensity: Interplay between the solar zenith angle and dipole tilt. *Journal of Geophysical Research: Space Physics*, 124, 6636–6653. <https://doi.org/10.1029/2019JA026707>.

---

Response:

The results shown in Figure 6 do agree with the key point in the Ohtani et al paper, that "The nightside westward electrojet (WEJ) is more intense when the ionosphere is dark," so a reference can be added to the revision.

C10

---

Referee #2:

Clean up language. For example: "atmosphere with a high level of precision" what does this mean? "that cause the derived conductivity to seem negative", why seem? "gradients act to transport energy flux from", transport flux? "and potentially useful conductivity values are lost." to who and how?

---

Response:

Thanks for pointing out that these sentences could have been worded better. The revised manuscript will include more clear language in the parts that are indicated.

---

Interactive comment on Ann. Geophys. Discuss., <https://doi.org/10.5194/angeo-2020-60>, 2020.

# Testing the Electrodynamic Method to Derive Height-Integrated Ionospheric Conductances

Daniel Weimer<sup>1,2</sup> and Thom Edwards<sup>3</sup>

<sup>1</sup>Center for Space Science and Engineering Research, Virginia Tech, Blacksburg, Virginia, USA

<sup>2</sup>National Institute of Aerospace, Hampton, Virginia, USA

<sup>3</sup>DTU Space, Technical University of Denmark, Copenhagen, Denmark

**Correspondence:** Daniel Weimer (dweimer@vt.edu)

**Abstract.** We have used empirical models for electric potentials and the magnetic fields in both space and on the ground to obtain maps of the height-integrated Pedersen and Hall ionospheric conductivities at high latitudes. This calculation required use of both “curl-free” and “divergence-free” components of the ionospheric currents, with the former obtained from magnetic fields that are used in a model of the field-aligned currents. The second component is from the equivalent current, usually associated with Hall currents, derived from the ground-level magnetic field. Conductances were calculated for varying combinations of the Interplanetary magnetic field (IMF) magnitude and orientation angle, as well as the dipole tilt angle. The results show that reversing the sign of the Y component of the IMF produces substantially different conductivity patterns. The Hall conductivities are largest on the dawn side in the upward, Region 2 field-aligned currents. Low electric field strengths in the Harang discontinuity lead to inconclusive results near midnight. Calculations of the Joule heating, obtained from the electric field and both components of the ionospheric current, are compared with the Poynting flux in space. The maps show some differences, while their integrated totals match to within 1%. Some of the Poynting flux that enters the polar cap is dissipated as Joule heating within the auroral ovals, where the conductivity is greater.

*Copyright statement.*

## 1 Introduction

The Earth’s ionosphere has a major role in the flow of currents and energy within the magnetosphere, or what is also known as the “geospace environment.” The currents in the ionosphere are responsible for geomagnetic effects seen at the Earth’s surface, and they also have a role in the high-latitude heating of the upper atmosphere. The magnitude of these effects is determined to a large extent by the level of conductivity in the ionosphere, and as such the conductivity needs to be accurately known for reliable geospace modeling. On the other hand, it can be argued that the conductivity values are not known with a high precision, and may be the least well-quantified part of the coupled magnetosphere-ionosphere system.

This problem is not due to a lack of understanding, as the basic equations that define the conductivity values are known. ~~On the other hand the formulas for~~ The reference books by Rees (1989); Prölss and Bird (2004); Brekke (2013) and others provide

formulas for calculating the Pedersen ( $\sigma_P$ ) and Hall ( $\sigma_H$ ) conductivities ~~are often~~. While these formulas are basically the same,  
they are presented in a variety of different ~~and confusing formats, such as in the reference books by~~  
25 ~~Rees (1989); Prölss and Bird (2004); Brekke (2013).~~ These formulas can be reduced to a more simple form formats, symbols,  
and notations, which can lead to confusion. A version that we find more preferable follows:

$$\sigma_P = \frac{n_e |e|}{B} \left[ \frac{r_e}{1 + r_e^2} + \sum_i C_i \frac{r_i}{1 + r_i^2} \right] \quad (1)$$

$$\sigma_H = \frac{n_e |e|}{B} \left[ \frac{1}{1 + r_e^2} - \sum_i C_i \frac{1}{1 + r_i^2} \right] \quad (2)$$

where  $n_e$  is the electron number density,  $B$  is the magnitude of the magnetic field,  $e$  is the fundamental constant for the charge  
30 of an electron, and  $C_i$  is the relative ion concentration for the  $i$ th ion species, that are assumed to have a total number density  
equal to that of the electrons. The ratio  $r_i$  is defined as:

$$r_i = \nu_{in} / \Omega_i = 1 / k_i \quad (3)$$

where  $k_i$  is the “ion mobility coefficient” (Brekke, 2013),  $\nu_{in}$  is the ion-neutral or electron-neutral collision frequency, and  $\Omega_i$   
refers to the cyclotron frequency:

$$35 \quad \Omega_i = |e| B / m_i \quad (4)$$

The  $r_e$  ratio is obtained substituting electrons for ions in Eq. (3) and (4). The absolute value of the electron charge is used in  
the equations above in order to reduce sign ambiguity. Equations (1) and (2) are similar to Eq. (5) and (6) by Mallinckrodt  
(1985) (with a sign correction), and simplified using Eq. (3) and (4).

The height-integrated values of these conductivities are often used, designated with upper-case symbols  $\Sigma_P$  and  $\Sigma_H$ . In  
40 order to calculate these height-integrated values it is necessary to know the magnetic field strength, electron temperature  
and number density, and ion and neutral composition and number densities of each species, at all altitudes within the iono-  
sphere. At low and mid-latitudes these quantities are better known and can be obtained from a reference magnetic field model,  
and familiar empirical models of the ionosphere and neutral atmosphere such as the “International Reference Ionosphere”  
(IRI) (Bilitza, 2001) and the Mass Spectrometer and Incoherent Scatter (MSIS) model (~~Hedin, 1991; Picone et al., 2002~~)  
45 (Hedin, 1991; Picone et al., 2002; Emmert et al., 2020).

These models require calculations within specialized programs to generate the needed quantities, so there have been a num-  
ber of attempts to construct more simple empirical formulas for the conductivity. These are mainly valid for the dayside, where  
solar extreme ultraviolet (EUV) radiation is the main contribution to ionization. The review paper by Brekke and Moen (1993)  
lists 12 different formulas spanning the years 1889 to 1992. More recently conductivity formulas were provided by Richmond  
50 (1995b), Galand and Richmond (2001), and Wiltberger et al. (2004). Assimilation techniques used in the Kamide-Richmond-  
Matsushita (KRM) (Kamide et al., 1981) and the Assimilated Mapping of Ionospheric Electrodynamics (AMIE) (Richmond



and Kamide, 1988) methods also need to use conductivities that are derived from such models. As the solar zenith angle is used in these formulas, they often produce a sharp gradient at the terminator, so Ridley et al. (2004) had added a scattering term to the solar contribution in order to produce a smoother transition over the terminator for a coupled, magnetosphere-ionosphere numerical simulation.

At high latitudes the knowledge of the basic parameters is much less than for the lower latitudes. Due to the auroral ionization and the convection of ionized plasma from the dayside to nightside it is nearly impossible to specify the state of the ionosphere and neutral atmosphere with ~~a high level of precision~~ high accuracy. In fact, the documentation for the IRI model states that “it provides monthly averages in the non-auroral ionosphere for magnetically quiet conditions” (http://ccmc.gsfc.nasa.gov/modelweb/ionos/iri.html). On the night side the ionization due to high-energy auroral particle precipitation contributes most significantly to conductivity enhancements, and this is where there is the greatest uncertainty. As indicated by Liemohn (2020), the space science community needs to reexamine the methods used to calculate the conductivity enhancements due to auroral particle precipitation.

Due to the number of “known unknowns,” the ionospheric conductivity in the high-latitude region remains as one of the least-well quantified parameters in geospace and the study of magnetosphere-ionosphere coupling, yet this is where the most important interactions take place. Global numerical models may estimate the conductivities using formulas that include sunlight ionization rates and ionization production rates from precipitating particles, the recombination rates, and the equilibrium densities, and empirical models and various measurements are often used. For example, Fuller-Rowell and Evans (1987) used electron energy influx and energies from National Oceanic and Atmospheric Administration (NOAA) Television Infrared Observation Satellites (TIROS) to build statistical patterns of these data. These were used in physics-based formulas in order to calculate the Pedersen and Hall conductivities as a function of altitude, as well as the hight-integrated values, which were then used to create maps ordered by an auroral activity index.

A similar to method was used by Hardy et al. (1987) using a statistical model of electron flux from Defense Meteorological Satellite Program (DMSP) measurements sorted by the Kp index. They had used empirical formulas derived from computations by Robinson et al. (1987), relating the conductances to the average energy and energy flux of the electrons. Another statistical technique reported by Ahn et al. (1998) had used radar measurements to derive conductivity, and compared these with ground observations of the magnetic perturbations in order to derive empirical relationships between them. They then used measurements of  $\Delta B$  to obtain global maps of the conductivity, which were compared with the results by Hardy et al. (1987).

While these statistical models have similar features, they are not in complete agreement. ~~A larger problem is that~~ Another issue is that these models use geomagnetic activity indices as the main parameter, and such indices are not readily available in real time. Index-based models ~~based on activity indices have only marginal utility, as they~~ do not take the Interplanetary Magnetic Field (IMF) vector into consideration. ~~It, and it~~ is known that the electric field and field-aligned current (FAC) patterns change significantly as the IMF ~~rotates. As it would be desirable to combine~~ changes orientation. As a conductivity model ~~with an electric field or FAC model~~ is often used with another model or numerical calculation of electric fields and

field-aligned currents, unrealistic results are obtained if the boundaries of these-the models do not properly align, or if they are not from-based on consistent IMF orientations.

An alternative technique for obtaining the conductivity, named the “the elementary current method,” (Amm, 2001) uses multiple satellite and ground magnetometer measurements for deriving the ionospheric currents. This method is based on splitting the ionospheric current vector into divergence-free ( $\mathbf{J}_{df}$ ) and curl-free ( $\mathbf{J}_{cf}$ ) parts. The total height-integrated ionospheric current that is perpendicular to the magnetic field lines is then written as:

$$\mathbf{J}_{\perp} = \mathbf{J}_{df} + \mathbf{J}_{cf} \quad (5)$$

Ground-based magnetometer data are used to derive the “divergence-free” ionospheric currents that are usually associated with Hall currents. The “curl-free” currents are derived from space-based magnetometer measurements that are sensing the field-aligned currents (FAC) that are linked to the divergence of the ionospheric currents. Thus, magnetometer measurements both above the ionosphere and on the ground are required in order to recover the full  $\mathbf{J}_{\perp}$ . More details about the derivations of these currents will follow in a later section.

If there are no neutral winds present, then from Ohm’s law for the ionospheric current sheet,

$$\mathbf{J}_{\perp} = \Sigma_P \mathbf{E}_{\perp} + \Sigma_H (\hat{\mathbf{B}}_{\perp} \times \mathbf{E}_{\perp}) \quad (6)$$

if measurements of the electric field is also available then both the Pedersen and Hall conductances can be obtained from:

$$\Sigma_P^* = \frac{\mathbf{J}_{\perp} \cdot \mathbf{E}_{\perp}}{|\mathbf{E}_{\perp}|^2} \quad (7)$$

$$\Sigma_H^* = \frac{\hat{\mathbf{r}} \cdot (\mathbf{J}_{\perp} \times \mathbf{E}_{\perp})}{|\mathbf{E}_{\perp}|^2} \quad (8)$$

As Amm (2001) had stated, “These equations have been derived under the assumption that the magnetic field lines are directed perpendicular to the ionospheric plane. If they are not, the conductance tensor gets off-diagonal elements, and polarization effects have to be included.” It was also noted that a small error in the direction of the vectors can produce inaccuracies, especially where the magnitude of the electric field is small. Amm (1998) indicated that the assumption that the magnetic field lines are assumed to be radial does not cause significant errors at latitudes above  $45^\circ$ .

We have added the \* superscripts to the conductivities in Eq. (7) and (8) to indicate that these derivations are approximations, particularly since the effect of the neutral winds are not included, and their influence is assumed to be small (Amm, 1995). As clarified by Amm et al. (2008), “In reality the total effective electric field  $\mathbf{E}'_{\perp} = \mathbf{E}_{\perp} + \mathbf{U} \times \mathbf{B}$  should be considered, where  $\mathbf{U}$  is the neutral wind velocity, but the neutral wind velocity is highly height dependent and notoriously difficult to measure; it is also 1–2 orders of magnitude smaller than the plasma velocity in the E-region, and so it is typically set to zero.”

In the example presented by Amm (2001) the Spherical Elementary Currents Systems (SECS) (Amm, 1997; Amm and Viljanen, 1999) method is used to obtain the divergence-free currents from “the upward continuation technique for magnetic

115 disturbance fields from the ground to the ionosphere” (Amm and Viljanen, 1999). Magnetometer measurements obtained from sites in Norway, Sweden, and Finland were used in combination with electric field values from the Scandinavian Twin Auroral Radar Experiment (STARE) coherent scatter radar. The method was demonstrated for a small area using simulated magnetic fields above the ionosphere produced by a current vortex, that were compared with measured values from an overhead pass by the four Cluster II satellites. In another example Amm et al. (2015) use the SECS methods to solve for the electric field,  
 120 currents, and conductivity in the ionosphere using only measurements from two of the European Space Agency’s (ESA) Swarm spacecraft. Solutions were obtained within a region spanning  $7^\circ$  in longitude by  $20^\circ$  in latitude, that bounded the parallel tracks of the two satellites.

The notations used in Eq. (6) to (8) closely follow those of Green et al. (2007), who had demonstrated their use to obtain maps of the height-integrated Pedersen and Hall conductivity over the entire polar region. The horizontal electric field in the  
 125 ionosphere ( $E_\perp$ ) was obtained from the SuperDARN radar array, with assimilation of drift-meter, electric field measurements on the DMSP satellite along one orbit path. The SuperDARN statistical model (Ruohoniemi and Greenwald, 1996, 2005) was used to help constrain the fit of  $E_\perp$ . Ground-based magnetometer data were used to construct a map of the divergence-free ionospheric current,  $J_{df}$ , using a Spherical Cap Harmonic Analysis (SCHA) (Haines, 1988) and the techniques described by Chapman and Bartels (1940) and Backus (1986). The curl-free current,  $J_{cf}$ , was derived from magnetic field measurements on  
 130 the DMSP, Iridium, and Ørsted satellites. All data were gathered over a one-hour period while measurements by the Advanced Composition Explorer (ACE) satellite indicated that the IMF was relatively stable. The final results by Green et al. (2007) showed maps of the derived Pedersen and Hall conductivities in their Figures 9 and 10, with grey regions indicating where there was too much uncertainty in the time-averaged radar measurements. Superposed contours showed the conductances obtained by combining the Rasmussen et al. (1988) model for the solar EUV contribution with the Hardy et al. (1987) model  
 135 for the particle precipitation.

While Green et al. (2007) show results for only one event, Amm’s method is perhaps the most direct way to measure the height-integrated ionospheric conductances, and as such it seems reasonable to give the technique a more thorough test. In this paper we use a similar calculation to generate more detailed maps of the conductivity for a wider range of conditions, including variations in IMF clock angle and dipole tilt angle. Our input data consist of outputs from three separate empirical  
 140 models that were derived from large data sets: A new model of the electric potentials (Edwards, 2019), a model of the ground-level geomagnetic perturbations (Weimer, 2013), and a new FAC model from satellite magnetometers (Edwards et al., 2020). Due to the need for both electric fields and currents, from magnetic field measurements on both the ground and in space, we prefer to call this the “electrodynamic method.”

We emphasize that the height-integrated conductances that we obtain show the relationship between the total, horizontal  
 145 current and the magnitude of the electric field measured above the ionosphere. Within the ionosphere the horizontal current density varies as a function of altitude, as demonstrated by Mallinckrodt (1985). Due to a finite conductivity parallel to the magnetic field within the ionosphere, the horizontal electric field may also vary with altitude. One minor but important detail that should be mentioned is that traditionally the height-integrated conductance values are obtained by an actual integration of the conductances that are computed at a range of altitude values using Eq. (1) and (2), as done by Mallinckrodt (1985).

150     The coupled magnetosphere-ionosphere system is a complex, dynamic system that is always changing, so often it is questioned whether or not statistical models can provide an accurate representation of the polar electric fields and currents. Nevertheless, all investigations have found consistent and repeatable results that show how the dipole tilt angle and magnitude and orientation of the IMF control the global-scale configuration of the ionosphere. The Discussion section lists a number of studies done by several groups using different data sets and methods.

## 155     **2   Derivation of the ionospheric electric fields and currents**

### **2.1   IMF Measurements**

All statistical models used here incorporated measurements of the IMF and solar wind velocity. Complicating the use of these data is the fact that comparisons of measurements of the IMF taken simultaneous with multiple spacecraft in the solar wind revealed unexpected variations in time lags between the detection of similar features (Weimer et al., 2002). Similar variability  
160 in the time lags exist between IMF measurements, taken at the Lagrange point named L1, and the arrival of this IMF at the Earth's magnetosphere. We use the technique described by Weimer and King (2008) to reduce the uncertainty in these delays. We propagate the IMF to the Earth's bow shock and then add another fixed delay to account for propagation through the bow shock, the interaction with the magnetosphere, and the time required for the ionosphere to respond and reconfigure.

The paper by Weimer et al. (2010) looked at measuring these time delays using measurements of the ground-level magnetic  
165 field, and also discussed the wide range of values for these delays that were reported in previous publications. Correlations with a coupling function of the IMF were computed as a function of the lag time, after prior propagation to the bow shock. Different time periods were used for averaging the IMF values, in multiples of 5 min. Very broad peaks in the correlations were seen, attributed to the variability in the system, with the correlation increasing as the width of the averaging period was increased. The peak correlations were found at lag times of 30 min on the day side and 40 to 45 min on the night side, measured  
170 from the start of the 5-min segment at the center. For the purpose of constructing global patterns using spherical harmonics, Weimer et al. (2010) used 25-min averages of the IMF, starting at 45 min before each data sample and ending at 20 min before.

For model development the ionospheric conditions are only associated with solar wind conditions in the past. The initial 20-min lag accounts for the time it takes for the IMF signal to propagate between the bow shock and ionosphere. The use of  
175 an averaging window 25-min long (reduced to 20 min for later models) accounts for the time needed for the ionosphere to reconfigure to a new state after changes in the IMF magnitude and orientation.

The calculations using the algorithm by Weimer and King (2008) are time consuming, and need to be done with a database that spans decades. Therefore the IMF data were processed in advance of the model derivations. For greatest efficiency the sliding averages of the solar wind velocity and the Y and Z components of the IMF are stored at 5-min increments for later  
180 use.

## 2.2 The electric fields

We use an updated electric potential model by Edwards (2019), which supplements the database from the Dynamics Explorer-2 spacecraft that was used by Weimer (2005b) with a substantially larger number of measurements from the Swarm spacecraft (Lomidze et al., 2019). Previous models had derived the least-error fit of the model coefficients from electric potentials that were obtained by integrating the measured electric field values along the satellite's path ~~;~~ ~~the latest version fit the coefficients during each polar pass. The same, time-averaged IMF and solar wind values were assigned to all potentials along the entire pass. The latest version of the model is different in that the coefficients are fit~~ directly from the electric field measurements. ~~Another change is that the average of,~~ rather than the integrated potentials. As described in the previous section, the IMF and solar wind values ~~that are~~ associated with each measurement ~~use 20-min averages that~~ were calculated at 5-min intervals, ~~rather than using one average for the entire polar pass. 20-min averaging periods are used, as in the previous version,~~ after accounting for ~~solar-wind-all~~ propagation delays.

On all satellites only the component of the electric field in the direction of motion was usable for the model fits. Like the prior version of the model, this latest version is constructed using SCHA (Haines, 1985), with Legendre functions of real, non-integer degree. All model coefficients were calculated using the entire dataset, or parameter space, without prior sorting into bins.

The electric potential model is in a reference frame that is co-rotating with the Earth. Modified magnetic apex coordinates (VanZandt et al., 1972; Richmond, 1995a) are used, and the electric potentials are assumed to be constant along magnetic field lines. The electric fields in the ionosphere are obtained from the derivatives of the potentials that are produced by the model. In all results shown here an altitude of 110 km is used.

## 2.3 The divergence-free currents

The divergence-free currents are obtained from the empirical model of the ground-level magnetic fields by Weimer (2013). This model was constructed from magnetometer measurements at 149 locations during an 8-year time period, along with the simultaneous IMF measurements from the Advanced Composition Explorer (ACE) spacecraft. All sites are located in the Northern hemisphere, extending down to the magnetic equator. Quiet-time, baseline values were subtracted from the measured magnetic fields, as described in detail by Weimer et al. (2010). ~~These data were then~~ The processing is similar to that done by the SuperMAG project (Gjerloev, 2012), which did not have any results publicly available when that work had begun.

The magnetic field measurements were translated and rotated to the magnetic apex coordinate system for use in the construction of the model. The model produces values for the Northward, Eastward, and Vertical components of the magnetic field perturbations given a specification of the Y and Z components of the IMF in Geocentric Solar Magnetic (GSM) coordinates, the solar wind velocity, dipole tilt angle, and the  $F_{10.7}$  index of solar radiation. The three components were modeled separately, without use of a scalar potential, and implicitly included the effects of internal, image sources.

The formulas described by Chapman and Bartels (1940), Haines (1988), and Haines and Torta (1994) were used to derive the “ionospheric equivalent current function” (Kamide et al., 1981; Richmond and Kamide, 1988). A detailed description of

the process is provided by Weimer (2019), which includes the separation of the magnetic effects into their internal and external  
 215 sources. The magnetic fields are calculated from the gradient of a scalar potential. A SCHA technique is employed, but since  
 the size of the spherical cap is  $90^\circ$  the associated Legendre polynomials with integer degree are used, rather than Legendre  
 functions of real, non-integer degrees. The end result is an expression for the external currents in terms of spherical harmonics:

$$\psi_E(\theta, \lambda) = \frac{a}{\mu_o} \sum_{k=1}^{34} \sum_{m=0}^{\min(k,3)} \frac{2k+1}{k+1} \left(\frac{R_2}{a}\right)^k P_k^m(\cos\theta) (g_k^{m,e} \cos m\lambda + h_k^{m,e} \sin m\lambda) \quad (9)$$

where  $R_2$  is the radius of the spherical shell on which the external currents are assumed to flow, and  $a$  is the radius of the  
 220 Earth. This “equivalent current” function  $\psi_E$  has units of Amperes (or kA). The current density vector is obtained from the  
 negative gradient of this function, rotated by  $90^\circ$ . We use a spherical shell at an altitude of 110 km. External currents in  
 the magnetosphere are also projected to this shell, including the ring current. As shown by Weimer (2019), better results are  
 obtained if adjustments are made to compensate for such current. At low latitudes the Solar Quiet ( $Sq^o$ ) current systems also  
 appear in these results (Matsushita, 1975), along with the magnetic effects of inter-hemispheric, field-aligned currents, and  
 225 magnetospheric currents (Yamazaki and Maute, 2017).

## 2.4 The curl-free currents

The new FAC model that we use was developed using a very large database of magnetic field measurements from the Ørsted,  
 Challenging Mini-satellite Payload (CHAMP), and Swarm missions, along with IMF values from ACE (Edwards et al., 2020).  
 230 [The magnetic field instruments on these satellites have better accuracy and sampling rates than the Iridium satellites used for  
 the Active Magnetosphere and Planetary Electrodynamics Response Experiment \(AMPERE\) \(Anderson et al., 2008, 2014\).](#)

Like the previous version of the model (Weimer, 2005b), this new [FAC](#) model is constructed using SCHA and it is based on  
 the mathematical derivations by Backus (1986), along with Maxwell’s equations. The field-aligned currents are related to the  
 the magnetic field perturbations above the ionosphere by

$$\mu_o \mathbf{J} = \nabla \times \Delta \mathbf{B}_\perp \quad (10)$$

where  $\Delta \mathbf{B}_\perp$  are the magnetic perturbations in the plane perpendicular to the currents. Following Backus (1986), the radial  
 235 FAC is a poloidal current that is related to a toroidal magnetic field, such that

$$\mu_o J_\parallel \hat{\mathbf{r}} = \nabla \times (\hat{\mathbf{r}} \times \nabla_\perp \psi) \quad (11)$$

where  $\psi$  is a “toroidal scalar” that has units of length times magnetic induction (Tm, or more commonly used, cTm).  $\nabla_\perp$  is a  
 horizontal (perpendicular) surface gradient that Backus (1986) refers to as  $\nabla_S$ . This last equation reduces to

$$J_\parallel = \nabla_\perp^2 \psi / \mu_o \quad (12)$$

As (12) can also be written as

$$J_\parallel = \nabla_\perp \cdot (\nabla_\perp \psi / \mu_o) = \nabla_\perp \cdot \mathbf{J}_{cf} \quad (13)$$

it is seen the FAC density is related to the divergence of the curl-free “potential current,” where

$$\mathbf{J}_{cf} = \nabla_{\perp} \psi / \mu_o = -\hat{\mathbf{r}} \times \Delta \mathbf{B}_{\perp} / \mu_o \quad (14)$$

240 and  $\hat{\mathbf{r}}$  is downward in the direction of the local magnetic field. A positive field-aligned current is also downward. This result indicates that the curl-free current is in the direction of the gradient of the toroidal scalar. Additionally, this gradient is rotated 90° from the direction of the toroidal component of the magnetic field, and vice versa.

The newest model by Edwards et al. (2020) differs from the predecessors in that, rather than first integrating the magnetic field measurements to obtain a magnetic potential, the [magnetic field](#) values measured on the spacecraft are used directly in the  
245 least-error fits. Preprocessing of the data involves subtraction of the Earth’s internal field and translation into magnetic apex coordinates. As in the case of the electric field model, the IMF and solar wind values were averaged over a 20-min window, at 5-min increments. The FAC is calculated directly from Eq. (10), rather than Eq. (12), and the curl-free currents are calculated from the right side of Eq. (14), rather than the middle part. In other words, rotating the modeled magnetic field by 90° and dividing by  $\mu_o$  provides the curl-free component of  $\mathbf{J}_{\perp}$  needed to solve for the conductivity with Eq. (7) and (8).

250 [An earlier work by Edwards et al. \(2017\) had compared the field-aligned currents with four different indices of solar radiation defined by Tobiska et al. \(2008\). The expected correlations between these indices and the FAC were verified, so the latest FAC model was constructed so that it could use any one of these indices. Since the ground-level magnetic field model had used only the  \$F\_{10.7}\$  index, this same index is the one that is used for the FAC model results that are presented here.](#)

### 3 Poynting flux and Joule heating

255 In the results that follow we also include comparative maps of the distribution of the perturbation Poynting flux and Joule heating. The perturbation Poynting vector is calculated from the electric field and perturbation magnetic field that is perpendicular to the field-lines carrying the current:

$$\mathbf{S}_p = \mathbf{E}_{\perp} \times \Delta \mathbf{B}_{\perp} / \mu_o \quad (15)$$

where  $\mu_o$  is the permeability of free space. As this perturbation Poynting vector has just one component that is parallel to the current flow, the magnitude of the vector (the rate of energy flow through a spherical surface) is frequently referred to as simply  
260 Poynting flux. While it is possible to calculate a Poynting flux using the full geomagnetic field, in the absence of currents the curl of this field is zero. As indicated by Kelley et al. (1991), a Poynting flux calculated with a curl-free magnetic field has no divergence within a closed surface, and it is essentially useless; currents need to be present for energy to be dissipated within a closed region. Another important point mentioned by Kelley et al. (1991) is that “the Poynting flux yields the correct energy input even if a neutral wind is present in the ionosphere, which is almost always the case.”

265 As we have available the two-component horizontal current, it is also useful calculate the distribution of the Joule heating:

$$H_J = (\mathbf{J}_{df} + \mathbf{J}_{cf}) \cdot \mathbf{E}_{\perp} \quad (16)$$

Comparing these two quantities is useful for the simple reason that the perturbation Poynting flux can be obtained from spacecraft measurements, while the distribution of the ionospheric Joule heating cannot be easily measured or calculated, even though the later is the quantity that is more desired. As pointed out by Richmond (2010), these two quantities are not necessarily the same, and it was postulated that “the associated perturbation Poynting flux can possibly be very different from the integrated energy dissipation below.”

## 4 Results

In the figures that follow are shown results of the ionospheric conductivity calculations using Eq. (7) and (8), for different combinations of dipole tilt angle (Laundal and Richmond, 2017) and the IMF clock angle (the arc tangent of the Y and Z components of the IMF in GCM coordinates). Also shown are maps of the quantities used to obtain these results, along with the associated mappings of the perturbation Poynting flux and Joule heating. These additional maps provide useful and interesting information. All examples are for idealized conditions (not for specific events) that assume steady state solar wind and IMF values.

Figure 1 shows results with an IMF magnitude of 10 nT in the GSM Y-Z plane input to the models, at a clock angle orientation of  $180^\circ$  (entirely southward, or  $B_Z = -10$ ), and a solar wind velocity of  $450 \text{ km s}^{-1}$ . The dipole tilt angle is  $0^\circ$ , and the  $F_{10.7}$  index 160 sfu. ~~Although the FAC model has the capability to use other solar indices, the ground-level magnetometer model included only the  $F_{10.7}$  index, so that is what we need to use.~~

In the top row of the figure, parts (a)-(c) shows the electric potential and the two horizontal components of the electric field. The longitudes are marked in Magnetic Local Time (MLT), in magnetic apex coordinates, with the sun at 12 noon. The gray area on the maps show the region that is outside of the spherical cap that is used in the SCHA functions in the model. The size of this cap varies with IMF conditions. While the derivatives of the potential are originally calculated in northward and eastward polar coordinates, it is more useful to convert these to duskward and sunward components for display and use in the calculations. For example, the typical electric potential pattern has a strong electric field in the duskward direction, directed from the positive peak on the dawn side toward the negative valley on the dusk side. If the northward and eastward components are shown then this pattern is not at all obvious. Minimum and maximum values of the potential and electric fields are indicated in the lower left and right corners of all contour maps, and the locations where these values are found are marked on the map with the diamond and plus symbols respectively. For clarity the levels chosen for the counter lines avoid values at exactly zero, as the contouring algorithm tends to entirely miss the zero contour around one of the two convection cells. As mentioned before, these potentials are in a co-rotating frame of reference.

In the second row of Fig. 1, parts (d)-(f) show the equivalent current function and the duskward and sunward components of the divergence-free currents that are calculated from the gradients of this function. Since the current flows along the direction of the contour lines, clockwise around the positive peak, the sunward component of the current flow has some resemblance to the duskward electric field. As these maps are derived from the magnetic perturbation model that covers the entire hemisphere (in magnetic apex coordinates), there is no gray boundary. The color bar scale for all horizontal currents is adjusted to approx-



imately match the largest magnitude of the sunward current. Currents outside of the the electric field convection pattern appear  
300 at lower latitudes, having opposite signs. As will be seen in other examples, the patterns that are found at lower latitudes have a  
strong dependence on the magnitude and orientation of the IMF. This behavior leads us to assume that these reversed currents  
at lower latitudes are due to the magnetic effects of magnetopause and field-aligned currents, that produce a false signature of  
ionospheric flow in the equivalent current function.

In the third row of Fig. 1, parts (g)-(i) show the field-aligned current and the duskward and sunward components of the  
305 curl-free current. Via Eq. (14), these currents are just the duskward and sunward magnetic fields, transposed with one sign  
change, and divided by  $\mu_o$ . A predominately sunward magnetic field in the polar cap translates to a duskward current. These  
currents closely resemble the electric fields, as expected. These two components of the magnetic fields were produced directly  
from the SCHA functions in the FAC model, and then the FAC is calculated from their curl, Eq. (10). This model version  
(Edwards et al., 2020) had fit the spacecraft magnetic field measurements to the duskward and sunward components in order  
310 to reduce the spurious, circular harmonics in the FAC that tend to result when using polar coordinates. The total sums of the  
upward (negative) and downward (positive) FAC, integrated over the spherical cap, are indicated in the upper left and right  
corners of the contour map in units of millions of Amperes (MA). As the density of contour lines in the FAC maps tends to  
get too crowded around the largest values, lines are drawn only for every third interval marked on the color bar. As before, the  
gray area on the maps show the region outside of the SCHA cap defined by the FAC model.

315 The fourth row starts with a map of the perturbation Poynting flux in the downward direction, Fig. 1(j), calculated with Eq.  
(15). The total energy flow into the ionosphere is in the upper-right corner, in Giga-watts (GW). We note that the new electric  
potential and FAC models produce Poynting flux maps that we consider to be more realistic than the results from the prior  
models (Weimer, 2005a), with levels that are higher within the polar cap and near the cusp. Sometimes there may be a slight  
mismatch between the electric potential model and the FAC model (derived from independent data sets), that results in the  
320 electric and magnetic fields reversing directions at not exactly the same locations; such misalignment manifests as a negative  
value of the Poynting flux. These negative fluxes are simply artifacts, and colored in lighter shades of gray. In general the two  
models match up very well at the electric field reversals, and these areas are rather small in size and magnitude. As with the  
FAC map, some contour lines are omitted for clarity.

The second map in the fourth row, Fig. 1(k), shows the Pedersen conductivity that is calculated with Eq. (7), but without  
325 including the divergence-free current, from the equivalent current function. While this result is not physically meaningful, it is  
useful to include the map for diagnostic purposes and to show the wrong answer that results if the total current is not used. As  
the calculation in Eq. (7) doesn't work where the magnitude of the electric field is very low, locations where this magnitude  
is small are flagged as invalid and colored in gray on the map, in addition to latitudes that are below the spherical cap of the  
electric field or FAC models. The limiting electric field magnitude is 3 mV/m or 7% of the peak magnitude, whichever is  
330 greater. ~~While it It would be desirable to increase the limiting electric field strength, to around 10 mV/m for example, if larger  
values are used the result is that the gray areas extend too far into the auroral ovals and potentially useful conductivity values  
are lost. The outcome with increasing this limit is that a greater number of results are marked as invalid within the auroral oval.~~

At the electric field reversals there are small areas where conductivity may appear to be negative or have abnormally high values. Negative values are indicated with a blue coloring on the map, but these values are not considered to be realistic or meaningful. Likewise, large, positive values near the convections reversals should be ignored. In all maps of conductivity the maximum values that are indicated in the lower-right corner of the maps excluded results at latitudes greater than  $68^\circ$ , in order to avoid the areas around the convection reversals. The color bars on all conductivity maps have a fixed range, unlike the others that are adjusted to accommodate the largest values. A green color shows where the conductivity is greater than zero but less than 3 mho.

The next map, Fig. 1(l), shows the Hall conductivity that is calculated with Eq. (8), but without using the curl-free current from the FAC model; ~~a wrong result again, but useful to include~~. The format is the same as the Pedersen conductivity. While this result is incorrect (as was Fig. 1(k)), it is useful to include as it shows why the divergence-free component alone is not sufficient for calculating the Hall conductivity, as might be assumed. Including these maps is useful for understanding the contributions that both components have on the final numerical results. In this map there are regions where the derived conductivity is negative, which is unrealistic. These areas are marked in shades of blue that darken as the value becomes more negative. While the alignment between the electric potential and equivalent current functions in Fig. 1(a) and 1(d) is generally good, on the dawn side these negative values appear where the current function reverses direction from the clockwise flow around the positive convection cell, or counter-clockwise around the negative convection cell. As we had mentioned earlier, it is thought that the reversed flows, and the unrealistic, negative conductivity values, result from interference from magnetospheric currents. All locations with negative values are considered to have invalid results.

The bottom row in Fig. 1 shows the results using the total currents, with the two components of the current combined together. At the left, Fig. 1(m) shows the Joule heating that is calculated with from the dot product of the electric field and this total current with Eq. (16). The differences between the Joule heating and the perturbation Poynting flux maps will be discussed in more detail in Section 6. Finally, Fig. 1(n) and 1(o) show the derived values of the Pedersen and Hall conductivities, using the total currents. The auroral oval is easily seen in these results, where the conductivity changes to values greater than 3 mho. The Hall conductivity has enhanced values near 6 MLT, that peak at 32 mho, while the largest Pedersen conductivity (45 mho) is found near midnight. The regions of higher conductivities in both maps correspond to upward field-aligned current, the blue regions in Fig. 1(g), including where this FAC passes through the gap between downward current near midnight. This is a common feature in all results. On the dawn side the upward current is the lower-latitude belt, often referred to as “Region 2”, while on the dusk side it is the inner-belt, called “Region 1.” On the other hand, the Pedersen conductivity that is calculated near midnight seems too large, and most likely not realistic. We will return to this subject later.

## 5 Results from other dipole tilt and IMF clock angles

In Fig. 2 and 3 are shown maps for dipole tilt angles of  $-23^\circ$  and  $+23^\circ$ , corresponding to winter and summer conditions, while the zero tilt in Fig. 1 corresponds to near equinox conditions. As the dipole tilt angle varies every day by about  $\pm 11^\circ$ , due to the offset of the magnetic pole from the rotation axis, there is a broad range of dates when the dipole tilt angle is at the specified

values; the reference to seasons does not refer to exact dates, but a generalized time period. The format of these Figures is the same as before. Both the Pedersen and Hall conductivities in Fig. 2(n) and (o), peak at 69 and 51 mho respectively, which are greater than for the equinox conditions. The maps for summer conditions in Fig. 3 show peak Pedersen and Hall conductivities of 56 and 35 mho, that are lower than the winter values yet greater than at equinox. The positive tilt angle in the summer produces enhanced conductivities on the dayside, as expected. The enhanced Hall conductivity seen near 6 MLT in all three graphs, Fig. 2(o) in particular, agrees with the results ~~found by Green et al. (2007)~~ presented by Green et al. (2007) with their proof of concept demonstration, in their Fig. 10.

One feature to note is that while the electric potentials have similar patterns in Fig. 1–3, the equivalent current function rotates as the dipole tilt angle changes, and exhibits a sharp twist near the pole under winter conditions (negative dipole tilt, Fig. 2). Another noticeable feature is found near midnight, where the region of enhanced conductivities passes through the region in the electric potential patterns where the negative, dusk potential cell wraps around and under the positive cell. This warp in the electric potential patterns, known as the Harang discontinuity (Gjerloev and Hoffman, 2001; Marghita et al., 2009), does not appear in the equivalent current functions.

Next we turn our attention to other IMF orientation angles. Figures 4 and 5 show graphs for IMF clock angles of  $90^\circ$  and  $270^\circ$ , corresponding to positive and negative values of the Y component, with  $B_Z = 0$ . The magnitude of the IMF is 10 nT, and the dipole tilt angle is zero, the same as in Fig. 1. In both cases the conductivities are lower than when the IMF is southward ( $180^\circ$  clock angle), with conductivity values being lowest at the  $270^\circ$  clock angle. Additionally, the electric potentials, total FAC, and total Poynting flux are also much lower than when the clock angle is  $180^\circ$ . In Fig. 5 the enhanced Pedersen conductivity previously seen near 0 MLT is noticeably absent. The westward electrojet is also reduced, the region of positive duskward current near 0 MLT in subplots (e) and (h) in all examples. Examples of the results for these two IMF clock angles with negative and positive tilt angles (winter and summer) are included in the supplementary information contained at <https://doi.org/10.5281/zenodo.3985988>. This supplement also contains a set of graphs showing the same combinations of IMF clock angle and dipole tilt angle, but with the IMF magnitude reduced from 10 to 5 nT. Similar variations in the conductivities are seen in these other examples, such as the lower values when the Y component is negative ( $270^\circ$ ).

In order to get a better comparison of ~~how the both the~~ effects that the dipole tilt and IMF clock angles ~~influences have on~~ the conductivity values on the dawn and dusk sides, Fig. 6 shows a graph of the Pedersen conductivity as a function of latitude from  $50^\circ$  to  $75^\circ$ , along a meridional slice at 4 hours MLT. Results were calculated for IMF magnitudes of 1 to 29 nT, at 1 nT intervals, and stacked vertically along the ordinate. The graphs are repeated on a three-by-three grid with dipole tilt angles of  $-23^\circ$  (left column),  $0^\circ$  (middle), and  $+23^\circ$  (right column) and IMF clock angles of  $90^\circ$  (top row),  $180^\circ$  (middle), and  $270^\circ$  (bottom row). Conductivities are indicated with the colors with the scale shown at the bottom of the figure. Gray areas on the graph show where there are no valid results because the location was outside of one of the model's lower-latitude boundary, or the electric field magnitude was too low. The patterns shift to lower latitudes as the magnitude of the IMF increases due to the expansion of the auroral ovals. Blue areas also show invalid results, where the divergence-free current function has a sign opposing the electric field. Fig. 7 shows the Hall conductivities in an identical format. Figures 8 and 9 repeat the graphs for

400 the dusk side at 20 MLT. The 4 and 20 MLT slices mostly avoid the artifacts found at both high and low latitudes at all clock angles.

These graphs show that the conductivities have an asymmetrical response to the clock angle variations. On the dawn side at 4 MLT with a 90° clock angle (top rows in Fig. 6 and 7) the conductivity values are generally larger than at 270° (bottom rows), while both are exceeded when the IMF is at 180° (middle rows). The tilt angle that corresponds to winter conditions (left columns) often produce the largest values. On the dusk side (Fig. 8 and 9) the southward IMF (180°, middle rows) again produces the larger conductivity values, but the seasonal (tilt angle) differences are not always as significant. Enhancements near 70° latitude (at 10 nT IMF) are produced by the upward, Region 1 currents.

## 6 Comparing the maps of Joule heating and Poynting Flux

As mentioned earlier, Richmond (2010) shows how the distributions of the perturbation Poynting flux and the Joule heat should ~~need~~ not be identical. The Poynting vector that enters the ionosphere is not necessarily the same as the height-integrated heating rate below. The paper by Vanhamäki et al. (2012) addresses this topic as well; ~~they show that when the divergence-free and curl-free parts of the ionospheric current are combined, their gradients act to transport energy flux from regions of low Pedersen conductance toward regions of higher conductance. If,~~ showing that if only the curl-free component of the current is used to compute the Joule heating, then the mathematical result is identical to the Poynting flux distribution ~~as (also~~ shown by Weimer (2005a) ~~and Vanhamäki et al. (2012)).~~ This result is simply due to the way the curl-free component is computed from the perturbation magnetic field and mathematical identities. ~~The But when the~~ divergence-free component is ~~what causes the added to the Joule heating calculation then there are~~ differences in the distribution of energy dissipation, ~~but it does not influence the total. As shown by Vanhamäki et al. (2012) in,~~ Vanhamäki et al. (2012) explain that “between these areas the Poynting flux is transported horizontally near the ionospheric plane. ... Conductance gradients give raise to divergence-free Pedersen currents that modify the spatial distribution (but not the total amount) of Joule heating and to curl-free Hall currents that modify the spatial distribution (but not the total amount) of vertical Poynting flux.” In their section 3.1 ,Vanhamäki et al. (2012) show that the dot product of the horizontal, curl-free electric field with  $J_{df}$  should integrate to zero within the boundary where the electric field goes to zero, the result of the mathematical identities, which is why the totals are the same.

Our results in Fig. 1 through 5 do show differences between the maps of the Joule heating (m) and the perturbation Poynting flux (j). The Joule heating in the high-latitude polar cap ~~tending-tends~~ to be lower than the Poynting flux, and greater in the auroral oval, where the conductances are greater, while the integrated sums are nearly ~~the same equal~~. In order to more clearly demonstrate the differences between regions, Fig. 10 shows a map of the dot product of the electric field with the  $J_{df}$  component alone, for the same case shown in Fig. 1. The ~~energy-transfer~~ transfer of energy from the polar regions to auroral oval is clear. The numbers in the upper left and right corners show the totals of just the areas with negative and positive values, respectively. The 2 GW difference between the totals is only 0.7% of the 303 GW total Poynting flux, and such differences are always less than 1% of the total Poynting flux. As the sums should be equal, the differences are the result of numerical error. The totals in the graphs were derived from 19531 data points at and above 50° latitude, forming 38597 triangles on the

spherical cap. Each side of the spherical triangles spans an arc length of roughly  $0.5^\circ$ . The values of each quantity are evaluated at the vertices, and the mean value within each triangle are multiplied with the triangle's area, and summed.

## 435 7 Discussion

We have applied three, separate empirical models to the formulas proposed by Amm (2001) to calculate the ionospheric Pedersen and Hall conductivities, producing new maps of these conductivities for various conditions. For the most part, the values of the conductivities that are produced seem reasonable. Enhanced conductivities in the auroral ovals are seen, as expected, with values in the range of 3 to 9 mhos under moderate conditions, with some regions having substantial enhancements on top of that. Another result that was entirely expected is that conductivity values increase as the Z component of the IMF becomes more negative. In general, the range of values that we obtain are within reason, in comparison with the results found by Fuller-Rowell and Evans (1987), Hardy et al. (1987) Ahn et al. (1998), and very recently, Carter et al. (2020). At large IMF magnitudes the conductivity results that are greater than 40 mho seem too extreme, but such values are also found in the model results reported by Mukhopadhyay et al. (27 August 2020).

At positive and negative values of the Y component of the IMF the conductivity results are very different. This is significant, since the existing statistical models of conductivity (see Introduction) do not account for the orientation of the IMF. Some of the conductivity values that we found seem to be greater than what are produced with existing models. Both the Hall and Pedersen conductivities are higher for winter, or negative dipole tilt, conditions, particularly on the dawn side and toward midnight. There are some gaps and artifacts produced where the various models don't exactly line up. The results shown in Fig. 7 are in agreement with the findings by Ohtani et al. (2019) that "the nightside westward electrojet (WEJ) is more intense when the ionosphere is dark."

The equivalent currents often have patterns at lower latitudes that cause the ~~derived conductivity to seem negative, values that~~ conductivity that is calculated to have negative values; such results are not realistic ~~that and~~ therefore rejected. Magnetospheric and low-latitude, field-aligned currents are the likely source such results. The correction that was employed to compensate for the ring current actually had little effect on these results, with differences in the maximum values on the order of 2 mho if the correction was either removed or doubled, so that adjustment doesn't seem to be an issue.

One persistent feature in the results is the presence of the extraordinarily large Pedersen conductivities in a narrow band near midnight. A close examination of the map data shows that the peak value occurs in a region of low electric field strength, exactly where the sunward electric field at midnight passes through zero while changing sign from negative to positive, as latitude decreases. The curl-free, sunward current changes sign also, about one-half of a degree to the north. The duskward electric field is weak, around  $5 \text{ mV m}^{-1}$ . There is a rather strong ( $> 200 \text{ mA m}^{-1}$ ) divergence-free (equivalent) current here in the duskward direction, part of the westward, electrojet. This Hall current does not change sign with the electric field, which results in the derived Hall conductivity changing signs from positive to negative. This same pattern is found often, and in association with the Harang region where the negative, dusk electric potential cell wraps around the positive cell near midnight, on the lower-latitude side (Gjerloev and Hoffman, 2001; Marghitu et al., 2009). When the IMF is in the -Y direction

(Fig. 5, 270° orientation) the extra large Pedersen conductivity is not present, the signature of the Harang discontinuity is weak, and the westward (duskward) currents at midnight are substantially lower.

Shue and Weimer (1994) had proposed that polarization electric fields around areas of enhanced conductivity are responsible for forming the Harang discontinuity; the effect is to block the divergence of Hall currents where there are gradients in the Hall conductivity. Further evidence of this concept has been provided by Nakamizo and Yoshikawa (2019), and our results are consistent with this hypothesis. Equations (7) and (8) are not accurate where there are conductivity gradients, so the derived conductivity values near midnight, from 22 to 2 MLT, should not be trusted. Likewise, the results near noon are also suspect.

The effects of the neutral winds, which are difficult to measure, are not included in these results. One result of the neutral winds is that the heating is not entirely ohmic, but includes frictional heating from the relative motion of plasma and neutrals (Vasyliūnas and Song, 2005), and acceleration of the neutrals.

Thayer (1998) indicates that a neutral wind in the direction of the  $\mathbf{E} \times \mathbf{B}$  vector will decrease the Joule heating rate, while a component of the neutral wind in the opposite direction will act to increase it. His observations in one particular time period indicated that the neutral winds could reduce the local Joule heating rate by over 75% in the upper E region while enhancing the local heating rate by nearly 50% in the lower E region, with an overall decrease of 40% in the height-integrated values. If the electric field is in a steady direction then the neutral winds act to reduce the Joule heating rate, but if the electric fields change directions suddenly then the effect is reversed (Thayer, 1998).

Billett et al. (2018) report that the reduction in the Joule heating due to the neutral winds primarily happens at high magnetic latitudes and in the dusk sector. They report on observations showing a persistent absence of a neutral winds in the dawn circulation cell, and hence a lower reduction. They report that “the percentage contribution of the wind correction to the area-integrated Joule heating rate can vary by  $\pm 14\%$  depending on season and geomagnetic activity level,” with a greater influence occurring in the hemispheric winter months.

While our results do not include the effects of the neutral winds, they do show how the currents and electric fields are related to each other under typical conditions, which implicitly includes whatever influence the neutral winds may have. The conductivity values obtained from the electrodynamic models could be what numerical modelers actually need in order to compute electric potentials from the field-aligned currents, or vice versa, if the neutral winds are not available. If the neutral winds are significant, then these results are not indicative of the true conductivity in the usual sense. However, the results do establish a relationship between the electric field above the ionosphere, obtained from measurements, and the currents within the ionosphere, also from measurements in space and on the ground. The current-voltage relationship is really what matters.

Both Richmond (2010) and Vanhamäki et al. (2012) had demonstrated differences between maps of the perturbation Poynting flux and the energy dissipation, but with contrived distributions of energy flow and conductivity. The results shown here are the first demonstration of the such differences using more realistic conditions. ~~Theoreticians may argue that the use of  $\mathbf{J}_{af}$  and  $\mathbf{J}_{cf}$  are only a mathematical device that have no physical significance. As such the distributions shown in Fig. 10 should be treated with some degree of skepticism, particularly in the presence of neutral winds.~~

~~The accuracy of the final results are difficult to ascertain with reasonable certainty, mainly due to the presence of the small-scale fluctuations in the original measurements that are not present in the models. Of course, our smoothed calculations~~

of the conductivities do not include the small-scale enhancements that occur within the auroral arcs that are non-stationary in time and space.

All three statistical models provide a large-scale representation of the electric and magnetic field variations. ~~The models have considerable smoothing of the small-scale fluctuations contained in the original measurements. The data include a mixture of~~  
505 ~~turbulent features that may persist only for a brief period. These models all use measurements that were taken as a function of~~  
~~time, and moving from one location to another. Such fluctuations produce standard deviations that seem large, on the order of~~  
~~20% to 50% of the peak magnitude while the satellites were moving in space, or in the case of the ground-based magnetometers,~~  
~~fixed in position while moving through local time. The original measurements contain fluctuations at various frequencies, with~~  
~~the most rapid fluctuations, lasting a few seconds or so, often having the largest magnitudes.~~ Examples of these fluctuations  
510 can be seen in the recent comparison by Lomidze et al. (2019), showing electric field measurements from the Swarm satellites  
compared with values from the earlier potential model (Weimer, 2005b). ~~Despite the fluctuations, the underlying patterns are in~~  
~~very good agreement.~~ The ~~end-result models are global-scale, spatial representations of the fields that do not include such rapid~~  
~~fluctuations. We consider these variations to be mainly temporal, including a mixture of turbulent features that may persist only~~  
~~for a brief period of time, and perhaps moving in space as well.~~

515 With the removal of such fluctuations, the global-scale patterns are consistent and repeatable. For example, the elec-  
tric potential patterns obtained from double-probe measurements on the Dynamics Explorer-2 satellite (~~Weimer, 2005a~~)  
(Weimer, 1995, 2005a) are very similar to results obtained from ground radar (Ruohoniemi and Greenwald, 1996, 2005;  
Cousins and Shepherd, 2010), an electron drift instrument on the Cluster spacecraft (Haaland et al., 2007), and ion drift me-  
ters on the Defense Meteorological Satellite Program (DMSP) (Papitashvili and Rich, 2002). The field-aligned current maps  
520 obtained from different datasets are also similar and repeatable. For examples, compare the results from magnetometers on  
Dynamics Explorer-2 (Weimer, 2001, 2005a), DMSP (Papitashvili et al., 2002), the Iridium constellation (Anderson et al.,  
2008), CHAMP and Swarm (Laundal et al., 2018), and the Ørsted-CHAMP-Swarm combination (Edwards et al., 2020) that  
we use here. Laundal et al. (2018) also show maps of the equivalent currents that resemble the ones shown here.

~~The greatest~~ One consequence of the temporal variations is that the errors calculated for each model seem large. During  
525 the fitting process the difference between the model outputs and the input values (the total square error) is minimized, and  
also output with the results. The standard deviations that are calculated are on the order of 20% to 50% of the model outputs  
at their peak values. Due to the influence of the high-frequency fluctuations, we do not consider such errors to be a true  
representation of the accuracy of the spatial models, considering the repeatable results just mentioned. As a result, the accuracy  
of the final conductivity values are difficult to ascertain with reasonable certainty. Of course, our smoothed calculations of  
530 the conductivities do not include the small-scale and meso-scale enhancements that occur within the auroral arcs that are  
non-stationary in time and space.

A significant source of error is actually in the IMF measurements, since the values that are measured by a spacecraft such as  
ACE may differ from what actually impacts the Earth's magnetosphere (~~Borovsky, 2018~~). ~~Such errors cannot be determined~~  
~~until additional spacecraft are placed in the upstream solar wind. The~~ (Weimer et al., 2002; Borovsky, 2018). As indicated by  
535 Burkholder et al. (2020), “the solar wind measured by a single spacecraft at L1 often does not impact Earth in a homogeneous



manner.” Such uncertainties in the propagated IMF were greater in the electric potential model ~~also as it had~~ required use of some IMF measurements from the International Sun-Earth Explorer 3 (ISEE3), which was located farther ~~upstream, causing greater uncertainty in the IMF~~ off to the side in the solar wind. The errors in the propagated IMF measurements cannot be fully evaluated until additional spacecraft are placed in the upstream solar wind. The presence of multiple, random and uncontrolled

540 variables often makes space physics research more difficult than laboratory experiments under controlled conditions.

Mukhopadhyay et al. (27 August 2020) had tested the usefulness of four conductivity models in a numerical simulation by evaluating whether or not there was an improvement in various metrics and skill scores for the predictions of ground-level magnetic perturbations. A similar approach could be used for evaluating the conductivity values derived with the electrodynamic method, including use in ionospheric assimilation codes and coupled, ionosphere-thermosphere models.

## 545 8 Conclusions

The lack of definitive values for the ionospheric conductivities is a significant problem in magnetospheric physics. Existing empirical models are not entirely in agreement, and typically use indices of activity level rather than solar wind and IMF measurements. Amm (2001) had presented formulas for obtaining the conductivity from measurements of the electric fields, and currents derived indirectly from magnetic field measurements on the ground and in space. The ~~demonstration of the~~ method-proof of concept demonstration by Green et al. (2007) had shown promising results, which presented an opportunity to try the technique with a combination of empirical models to obtain estimates of the height-integrated Pedersen and Hall conductivities. Maps of the high-latitude, ionospheric conductivities were derived for varying combinations of the dipole tilt angle, IMF magnitude, and IMF orientation angle in the GSM Y-Z plane. There are places where the technique fails to produce valid conductivity values, so the results are not entirely satisfactory. Nevertheless, these findings should still be of some use to

555 the space science community.

The conductivity maps that were obtained have features that were expected on the basis of prior knowledge, such as enhancements within the auroral oval and increasing conductivity as the Z component of the IMF becomes more negative. The dawn and dusk sides do not have a symmetric response with respect to flipping the sign of  $B_Y$ . It was found that reversing the sign of the Y component of the IMF results in substantially different conductivity patterns, with values that are generally

560 lower when  $B_Y$  is negative, corresponding to clock angles near  $270^\circ$ . Changes in the dipole tilt angle also have a significant influence. These factors need to be considered in ~~all future conductivity models~~ the future development of conductivity models and their use.

*Code and data availability.* An archive of graphs and data can be found at <https://doi.org/10.5281/zenodo.3985988>. This archive includes the supplement with additional illustrations mentioned in the text, and reproductions of Fig. 1 to 5, but arranged in a horizontal, landscape

565 format for easier viewing on a computer screen, and similar figures generated for other cases. Maps of the dot product of the electric field and divergence-free current in each case are also included as well as the equivalent current functions down to a latitude of  $35^\circ$ . Digital data files containing the conductivity results are contained in the archive, along with Interactive Data Language (IDL) software code needed to



read and interpolate these data. All data needed to generate Fig. 6 to 9 are included, along with the associated graphs, plus additional cases at other IMF clock angles. Altogether, 696 sets are included. The archive contains the ground-level magnetic field model output values that are used to calculate the equivalent currents, the resulting spherical harmonic coefficients, and the SCHA coefficients from the electric potential and FAC models.

Data used in the development of the magnetic perturbation model are listed in the publications by Weimer et al. (2010) and Weimer (2013). Data used in the development of the field-aligned current model are available through Edwards et al. (2020). The electric potential model (Edwards, 2019) was developed with the Swarm cross-track ion drift data available at <http://earth.esa.int/swarm/> and Dynamics Explorer-2 Vector Electric Field measurements at <https://cdaweb.gsfc.nasa.gov/>. All models used solar wind and IMF measurements from the IMP8, ISEE3, and ACE satellites, at <https://cdaweb.gsfc.nasa.gov/>.

*Author contributions.* The writing of this article was led by DW with contributions from TE. DW created all figures. TE created the empirical FAC and electric potential models, and DW created the magnetic perturbation model and the equivalent currents.

*Competing interests.* The authors declare that they have no competing interests.

*Acknowledgements.* Daniel Weimer and Thom Edwards were supported by NSF grants AGS-1638270 and PLR-1543364 to Virginia Tech, with additional support from a subcontract to Hampton University, on NASA grant NNX15AE05G. The authors thank Art Richmond for helpful comments on an earlier version of the manuscript.

## References


- Ahn, B.-H., Richmond, A. D., Kamide, Y., Kroehl, H. W., Emery, B. A., de la Beaujardiere, O., and Akasofu, S.-I.: An ionospheric conductance model based on ground magnetic disturbance data, *J. Geophys. Res.*, 103, 14 769–14 780, <https://doi.org/10.1029/97JA03088>, 1998.
- Amm, O.: Direct determination of the local ionospheric Hall conductance distribution from two-dimensional electric and magnetic field data: Application of the method using models of typical ionospheric electrodynamic situations, *J. Geophys. Res.*, 100, 21 473–21 488, <https://doi.org/10.1029/95JA02213>, 1995.
- Amm, O.: Ionospheric Elementary Current Systems in Spherical Coordinates and Their Application, *J. Geomag. Geoelectr.*, 49, 947–955, <https://doi.org/10.5636/jgg.49.947>, 1997.
- Amm, O.: Method of characteristics in spherical geometry applied to a Harang-discontinuity situation, *Annales Geophysicae*, 16, 413–424, <https://doi.org/10.1007/s00585-998-0413-2>, 1998.
- Amm, O.: The elementary current method for calculating ionospheric current systems from multisatellite and ground magnetometer data, *J. Geophys. Res.*, 106, 24 843–24 856, <https://doi.org/10.1029/2001JA900021>, 2001.
- Amm, O. and Viljanen, A.: Ionospheric disturbance magnetic field continuation from the ground to the ionosphere using spherical elementary currents systems, *Earth Planets Space*, 51, 431–440, <https://doi.org/10.1186/BF03352247>, 1999.
- Amm, O., Aruliah, A., Buchert, S. C., Fujii, R., Gjerloev, J. W., Ieda, A., Matsuo, T., Stolle, C., Vanhamäki, H., and Yoshikawa, A.: Towards understanding the electrodynamics of the 3-dimensional high-latitude ionosphere: present and future, *Ann. Geophys.*, 26, 3913–3932, <https://doi.org/10.5194/angeo-26-3913-2008>, 2008.
- Amm, O., Vanhamäki, H., Kauristie, K., Stolle, C., Christiansen, F., Haagmans, R., Masson, A., Taylor, M. G. G. T., Floberghagen, R., and Escoubet, C. P.: A method to derive maps of ionospheric conductances, currents, and convection from the Swarm multisatellite mission, *Journal of Geophysical Research: Space Physics*, 120, 3263–3282, <https://doi.org/10.1002/2014JA020154>, 2015.
- Anderson, B. J., Korth, H., Waters, C. L., Green, D. L., and Stauning, P.: Statistical Birkeland current distributions from magnetic field observations by the Iridium constellation, *Ann. Geophys.*, 26, 671–687, <https://doi.org/10.5194/angeo-26-671-2008>, 2008.
- Anderson, B. J., Korth, H., Waters, C. L., Green, D. L., Merkin, V. G., Barnes, R. J., and Dyrud, L. P.: Development of large-scale Birkeland currents determined from the Active Magnetosphere and Planetary Electrodynamics Response Experiment, *Geophys. Res. Lett.*, 41, 3017–3025, <https://doi.org/10.1002/2014GL059941>, 2014.
- Backus, G.: Poloidal and toroidal fields in geomagnetic field modeling, *Rev. Geophys.*, 24, 75, 1986.
- Bilitza, D.: International reference ionosphere 2000, *Radio Science*, 36, 261–275, 2001.
- Billett, D. D., Grocott, A., Wild, J. A., Walach, M.-T., and Kosch, M. J.: Diurnal Variations in Global Joule Heating Morphology and Magnitude Due To Neutral Winds, *Journal of Geophysical Research: Space Physics*, 123, 2398–2411, <https://doi.org/10.1002/2017JA025141>, 2018.
- Borovsky, J. E.: The spatial structure of the oncoming solar wind at Earth and the shortcomings of a solar-wind monitor at L1, *Journal of Atmospheric and Solar-Terrestrial Physics*, 177, 2 – 11, <https://doi.org/10.1016/j.jastp.2017.03.014>, 2018.
- Brekke, A.: *Physics of the Upper Polar Atmosphere*, Berlin, Heidelberg: Springer Berlin Heidelberg, second edn., <https://doi.org/10.1007/978-3-642-27401-5>, 2013.
- Brekke, A. and Moen, J.: Observations of high latitude ionospheric conductances, *J. Atmos. Sol. Terr. Phys.*, 55, 1493–1512, 1993.

- Burkholder, B. L., Nykyri, K., and Ma, X.: Use of the L1 Constellation as a Multispacecraft Solar Wind Monitor, *Journal of Geophysical Research: Space Physics*, 125, e2020JA027978, <https://doi.org/10.1029/2020JA027978>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2020JA027978>, 2020.
- Carter, J. A., Milan, S. E., Paxton, L. J., Anderson, B. J., and Gjerloev, J.: Height-Integrated Ionospheric Conductances Parameterized By Interplanetary Magnetic Field and Substorm Phase, *Journal of Geophysical Research: Space Physics*, 125, e2020JA028121, <https://doi.org/10.1029/2020JA028121>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2020JA028121>, 2020.
- Chapman, S. and Bartels, J.: *Geomagnetism*, vol. Vol. II, Analysis of the data, and physical theories, Oxford University Press, London, 1940.
- Cousins, E. D. P. and Shepherd, S. G.: A dynamical model of high-latitude convection derived from SuperDARN plasma drift measurements, *Journal of Geophysical Research: Space Physics*, 115, <https://doi.org/10.1029/2010JA016017>, <http://dx.doi.org/10.1029/2010JA016017>, a12329, 2010.
- Edwards, T. R.: A New Suite of Ionospheric Electrodynamics Models, Phd thesis, Virginia Polytechnic Institute and State University, 2019.
- Edwards, T. R., Weimer, D. R., Tobiska, W. K., and Olsen, N.: Field-Aligned Current Response to Solar Indices, *J. Geophys. Res. Space Physics*, 122, 5798–5815, <https://doi.org/10.1002/2016JA023563>, 2017.
- Edwards, T. R., Weimer, D. R., Olsen, N., Lühr, H., Tobiska, W. K., and Anderson, B. J.: A Third Generation Field-Aligned Current Model, *J. Geophys. Res. Space Physics*, 125, <https://doi.org/10.1029/2019JA027249>, 2020.
- Emmert, J. T., Drob, D. P., Picone, J. M., Siskind, D. E., Jones Jr, M., Mlynczak, M. G., Bernath, P. F., Chu, X., Doornbos, E., Funke, B., Goncharenko, L. P., Hervig, M. E., Schwartz, M. J., Sheese, P. E., Vargas, F., Williams, B. P., and Yuan, T.: NRLMSIS 2.0: A whole-atmosphere empirical model of temperature and neutral species densities, *Earth and Space Science*, p. e2020EA001321, <https://doi.org/10.1029/2020EA001321>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2020EA001321>, 2020.
- Fuller-Rowell, T. J. and Evans, D. S.: Height-Integrated Pedersen and Hall Conductivity Patterns Inferred From the TIROS-NOAA Satellite Data, *J. Geophys. Res.*, 92, 7606–7618, 1987.
- Galand, M. and Richmond, A.: Ionospheric electrical conductances produced by auroral proton precipitation, *J. Geophys. Res.*, 106, 117–125, 2001.
- Gjerloev, J. W.: The SuperMAG data processing technique, *J. Geophys. Res. Space Physics*, 117, <https://doi.org/10.1029/2012JA017683>, 2012.
- Gjerloev, J. W. and Hoffman, R. A.: The convection electric field in auroral substorms, *Journal of Geophysical Research: Space Physics*, 106, 12 919–12 931, <https://doi.org/10.1029/1999JA000240>, 2001.
- Green, D. L., Waters, C. L., Korth, H., Anderson, B. J., Ridley, A. J., and Barnes, R. J.: Technique: Large-scale ionospheric conductance estimated from combined satellite and ground-based electromagnetic data, *J. Geophys. Res.*, 112, <https://doi.org/10.1029/2006JA012069>, 2007.
- Haaland, S. E., Paschmann, G., Forster, M., Quinn, J. M., McIlwain, R. B. T. C. E., Vaith, H., Puhl-Quinn, P., and Kletzing, C. A.: High-latitude plasma convection from Cluster EDI measurements: method and IMF-dependence, *Ann. Geophys.*, 25, 239–253, 2007.
- Haines, G.: Computer programs for spherical cap harmonic analysis of potential and general fields, *Computers and Geosciences*, 14, 413–447, [https://doi.org/10.1016/0098-3004\(88\)90027-1](https://doi.org/10.1016/0098-3004(88)90027-1), 1988.
- Haines, G. V.: Spherical cap harmonic analysis, *J. Geophys. Res.*, 90, 2583, <https://doi.org/10.1029/JB090iB03p02583>, 1985.
- Haines, G. V. and Torta, J. M.: Determination of equivalent current sources from spherical cap harmonic models of geomagnetic field variations, *Geophys. Res. Int.*, 118, 499–514, <https://doi.org/10.1111/j.1365-246X.1994.tb03981.x>, 1994.

- Hardy, D. A., Gussenhoven, M. S., Raistrick, R., and McNeil, W. J.: Statistical and functional representations of the patterns of auroral energy flux, number flux, and conductivity, *J. Geophys. Res.*, 92, 12 275, 1987.
- Hedin, A. E.: Extension of the MSIS Thermosphere Model into the Middle and Lower Atmosphere, *J. Geophys. Res.*, 96, 1159, 1991.
- Kamide, Y., Richmond, A. D., and Matsushita, S.: Estimation of ionospheric electric fields, ionospheric currents, and field-aligned currents from ground magnetic records, *Journal of Geophysical Research*, 86, 801–813, <https://doi.org/10.1029/JA086iA02p00801>, 1981.
- 660 Kelley, M. C., Knudsen, D. J., and Vickrey, J. F.: Poynting flux measurements on a satellite: A diagnostic tool for space research, *Journal of Geophysical Research: Space Physics*, 96, 201–207, <https://doi.org/10.1029/90JA01837>, 1991.
- Laundal, K. M. and Richmond, A. D.: Magnetic coordinate systems, *Space Science Reviews*, 206, 27–59, <https://doi.org/10.1007/s11214-016-0275-y>, 2017.
- 665 Laundal, K. M., Finlay, C. C., Olsen, N., and Reistad, J. P.: Solar wind and seasonal influence on ionospheric currents from Swarm and CHAMP measurements, *J. Geophys. Res. Space Physics*, 123, 4402–429, <https://doi.org/10.1029/2018JA025387>, 2018.
- Liemohn, M. W.: The Case for Improving the Robinson Formulas, *Journal of Geophysical Research: Space Physics*, 125, e2020JA028 332, <https://doi.org/10.1029/2020JA028332>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2020JA028332>, 2020.
- Lomidze, L., Burchill, J. K., Knudsen, D. J., Kouznetsov, A., and Weimer, D. R.: Validity Study of the Swarm Horizontal Cross-Track Ion Drift Velocities in the High-Latitude Ionosphere, *Earth and Space Science*, 6, 411–432, <https://doi.org/10.1029/2018EA000546>, 2019.
- 670 Mallinckrodt, A. J.: A numerical simulation of auroral ionospheric electrodynamics, *J. Geophys. Res.*, 90, 409–417, <https://doi.org/10.1029/JA090iA01p00409>, 1985.
- Marghitu, O., Karlsson, T., Klecker, B., Haerendel, G., and McFadden, J.: Auroral arc and oval electrodynamics in the Harang region, *Journal of Geophysical Research: Space Physics*, 114, <https://doi.org/10.1029/2008JA013630>, 2009.
- 675 Matsushita, S.: Morphology of slowly-varying geomagnetic external fields – a review, *Physics of the Earth and Planetary Interiors*, 10, 299–312, [https://doi.org/10.1016/0031-9201\(75\)90056-4](https://doi.org/10.1016/0031-9201(75)90056-4), 1975.
- Mukhopadhyay, A., Welling, D. T., Liemohn, M. W., Ridley, A. J., Chakraborty, S., and Anderson, B. J.: Conductance Model for Extreme Events: Impact of Auroral Conductance on Space Weather Forecasts, submitted to *Space Weather*, pp. arXiv [preprint], [arXiv:2008.12 276v1](https://arxiv.org/pdf/2008.12276v1), <https://arxiv.org/pdf/2008.12276.pdf>, 27 August 2020.
- 680 Nakamizo, A. and Yoshikawa, A.: Deformation of Ionospheric Potential Pattern by Ionospheric Hall Polarization, *Journal of Geophysical Research: Space Physics*, 124, 7553–7580, <https://doi.org/10.1029/2018JA026013>, 2019.
- Ohtani, S., Gjerloev, J. W., Johnsen, M. G., Yamauchi, M., Brändström, U., and Lewis, A. M.: Solar Illumination Dependence of the Auroral Electrojet Intensity: Interplay Between the Solar Zenith Angle and Dipole Tilt, *Journal of Geophysical Research: Space Physics*, 124, 6636–6653, <https://doi.org/10.1029/2019JA026707>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019JA026707>, 2019.
- 685 Papitashvili, V. O. and Rich, F. J.: High-latitude ionospheric convection models derived from Defense Meteorological Satellite Program ion drift observations and parameterized by the interplanetary magnetic field strength and direction, *Journal of Geophysical Research: Space Physics*, 107, <https://doi.org/10.1029/2001JA000264>, <http://dx.doi.org/10.1029/2001JA000264>, 2002.
- Papitashvili, V. O., Christiansen, F., and Neubert, T.: A new model of field-aligned currents derived from high-precision satellite magnetic field data, *Geophys. Res. Lett.*, 29, <https://doi.org/10.1029/2001GL014207>, 2002.
- 690 Picone, J., Hedin, A., Drob, D., and Aikin, A.: NRLMSISE-00 empirical model of the atmosphere: Statistical comparisons and scientific issues, *J. Geophys. Res.*, 107, <https://doi.org/10.1029/2002JA009430>, 2002.
- Prölss, G. W. and Bird, M. K.: *Physics of the Earth's Space Environment: an introduction*, Springer-Verlag Berlin Heidelberg, ISBN 3-540-21426-7, 2004.


- 695 Rasmussen, C. E., Schunk, R. W., and Wickwar, V. B.: A photochemical equilibrium model for ionospheric conductivity, *Jgr*, 93, 9831–9840, 1988.
- Rees, M. H.: *Physics and Chemistry of the Upper Atmosphere*, Cambridge Univ. Press, New York, 1989.
- Richmond, A. D.: Ionospheric electrodynamics using magnetic apex coordinates, *J. Geomag. Geoelectr.*, 47, 191, <https://doi.org/10.5636/jgg.47.191>, 1995a.
- 700 Richmond, A. D.: Ionospheric electrodynamics, in: *Handbook of Atmospheric Electrodynamics*, vol. 2, edited by Volland, H., pp. 249–290, CRC Press, Boca Raton, Fla, 1995b.
- Richmond, A. D.: On the ionospheric application of Poynting’s theorem, *Journal of Geophysical Research: Space Physics*, 115, <https://doi.org/10.1029/2010JA015768>, 2010.
- Richmond, A. D. and Kamide, Y.: Mapping electrodynamic features of the high-latitude ionosphere from localized observations: technique, *J. Geophys. Res.*, 93, 5741–5759, <https://doi.org/10.1029/JA093iA06p05741>, 1988.
- 705 Ridley, A. J., Gombosi, T. I., and DeZeeuw, D. L.: Ionospheric control of the magnetosphere: Conductance, *Annales Geophysicae*, 22, 567–584, 2004.
- Robinson, R. M., Vondrak, R. R., Miller, K., Babbs, T., and Hardy, D.: On calculating ionospheric conductances from the flux and energy of precipitating electrons, *J. Geophys. Res.*, 92, 2565, 1987.
- Ruohoniemi, J. M. and Greenwald, R. A.: Statistical patterns of high-latitude convection obtained from Goose Bay HF radar observations, *J. Geophys. Res.*, 101, 21 743–21 763, <https://doi.org/10.1029/96JA01584>, 1996.
- 710 Ruohoniemi, J. M. and Greenwald, R. A.: Dependencies of highlatitude plasma convection: Consideration of interplanetary magnetic field, seasonal, and universal time factors in statistical patterns, *J. Geophys. Res.*, 110, <https://doi.org/10.1029/2004JA010815>, 2005.
- Shue, J.-H. and Weimer, D. R.: The relationship between ionospheric convection and magnetic activity, *J. Geophys. Res. Space Physics*, 99, 401–415, <https://doi.org/10.1029/93JA01946>, 1994.
- 715 Thayer, J. P.: Height-resolved Joule heating rates in the high-latitude E region and the influence of neutral winds, *Journal of Geophysical Research: Space Physics*, 103, 471–487, <https://doi.org/10.1029/97JA02536>, 1998.
- Tobiska, W. K., Bouwer, S. D., and Bowman, B. R.: The development of new solar indices for use in thermospheric density modeling, *J. Atmos. Sol. Terr. Phys.*, 70, 803, 2008.
- Vanhamäki, H., Yoshikawa, A., Amm, O., and Fujii, R.: Ionospheric Joule heating and Poynting flux in quasi-static approximation, *Journal of Geophysical Research: Space Physics*, 117, <https://doi.org/10.1029/2012JA017841>, 2012.
- 720 VanZandt, T. E., Clark, W. L., and Warnock, J. M.: Magnetic Apex Coordinates: A magnetic coordinate system for the ionospheric  $F_2$  layer, *J. Geophys. Res.*, 77, 2406, <https://doi.org/10.1029/JA077i013p02406>, 1972.
- Vasyliūnas, V. M. and Song, P.: Meaning of ionospheric Joule heating, *J. Geophys. Res.*, 110, A02301, <https://doi.org/10.1029/2004JA010615>, 2005.
- 725 Weimer, D.: Models of high-latitude electric potentials derived with a least error fit of spherical harmonic coefficients, *J. Geophys. Res.*, 100, 19,595, <https://doi.org/10.1029/95JA01755>, 1995.
- Weimer, D.: Maps of field-aligned currents as a function of the interplanetary magnetic field derived from Dynamic Explorer 2 data, *J. Geophys. Res.*, 106, 12,889, <https://doi.org/10.1029/2000JA000295>, 2001.
- Weimer, D. R.: Improved ionospheric electrodynamic models and application to calculating Joule heating rates, *J. Geophys. Res.*, 110, A05306, <https://doi.org/10.1029/2004JA010884>, 2005a.
- 730

- Weimer, D. R.: Predicting Surface Geomagnetic Variations Using Ionospheric Electrodynamic Models, *J. Geophys. Res.*, 110, A12307, <https://doi.org/10.1029/2005JA011270>, 2005b.
- Weimer, D. R.: An empirical model of ground-level geomagnetic perturbations, *Space Weather*, 11, 107–120, <https://doi.org/10.1002/swe.20030>, 2013.
- 735 Weimer, D. R.: Derivation of hemispheric ionospheric current functions from ground-level magnetic fields, *J. Geophys. Res. Space Physics*, 124, <https://doi.org/10.1029/2018JA026191>, 2019.
- Weimer, D. R. and King, J. H.: Improved calculations of interplanetary magnetic field phase front angles and propagation time delays, *J. Geophys. Res.*, 113, A01105, <https://doi.org/10.1029/2007JA012452>, 2008.
- Weimer, D. R., Ober, D. M., Maynard, N. C., Burke, W. J., Collier, M. R., McComas, D. J., Nagai, T., and Smith, C. W.: Variable time delays  
 740 in the propagation of the interplanetary magnetic field, *J. Geophys. Res.*, 107, <https://doi.org/10.1029/2001JA009102>, 2002.
- Weimer, D. R., Clauer, C. R., Engebretson, M. J., Hansen, T. L., Gleisner, H., Mann, I., and Yumoto, K.: Statistical Maps of Geomagnetic Perturbations as a Function of the Interplanetary Magnetic Field, *J. Geophys. Res.*, 115, A10320, <https://doi.org/10.1029/2010JA015540>, 2010.
- Wiltberger, M., Wang, W., Burns, A. G., Solomon, S. C., Lyon, J. G., and Goodrich, C. C.: Initial results from the coupled magnetosphere  
 745 ionosphere thermosphere model: Magnetospheric and ionospheric responses, *J. Atmos. Sol. Terr. Phys.*, 66, 1411, 2004.
- Yamazaki, Y. and Maute, A.: Sq and EEJ—A Review on the Daily Variation of the Geomagnetic Field Caused by Ionospheric Dynamo Currents, *Space Science Reviews*, 206, <https://doi.org/10.1007/s11214-016-0282-z>, 2017.




**Figure 1.** Conductivity input data and results, for IMF  $B_T$  magnitude 10 nT at  $180^\circ$  clock angle, the solar wind velocity is 450 km s $^{-1}$ , the  $F_{10.7}$  index is 160 sfu, and the dipole tilt angle is  $0^\circ$  corresponding to near equinox. Details are explained in the text.






**Figure 2.** Conductivity input data and results for the same conditions as in Fig. 1, except that the dipole tilt angle is for winter conditions. The IMF  $B_T$  magnitude is 10 nT at  $180^\circ$ , the solar wind velocity is 450 km s $^{-1}$ , the  $F_{10.7}$  index 160 sfu, and the dipole tilt angle is  $-23^\circ$ .







**Figure 3.** Conductivity input data and results for the same conditions as in Fig. 1, except that the dipole tilt angle is for summer conditions. The IMF  $B_T$  magnitude is 10 nT at  $180^\circ$ , the solar wind velocity is 450 km s $^{-1}$ , the  $F_{10.7}$  index 160 sfu, and the dipole tilt angle is  $+23^\circ$ .




**Figure 4.** Conductivity input data and results for the same conditions as in Fig. 1, except that IMF clock angle is changed  $90^\circ$ . The IMF  $B_T$  magnitude is 10 nT, the solar wind velocity 450 km s $^{-1}$ , the  $F_{10.7}$  index 160 sfu, and the dipole tilt angle is  $0^\circ$ .




**Figure 5.** Conductivity input data and results for the same conditions as in Fig. 1, except that IMF clock angle is changed  $270^\circ$ . The IMF  $B_T$  magnitude is 10 nT, the solar wind velocity 450 km s $^{-1}$ , the  $F_{10.7}$  index 160 sfu, and the dipole tilt angle is  $0^\circ$ .



**Figure 6.** Pedersen conductivity as a function of latitude from  $50^\circ$  to  $75^\circ$ , at 4 hours MLT. Values were calculated for IMF magnitudes of 1 to 29 nT, at 1 nT intervals, indicated on the ordinate. Multiple graphs are shown for dipole tilt angles of  $-23^\circ$  (left column),  $0^\circ$  (middle), and  $+23^\circ$  (right column) and IMF clock angles of  $90^\circ$  (top row),  $180^\circ$  (middle), and  $270^\circ$  (bottom row). Conductivity values are colored according to the scale at the bottom. Gray and blue areas indicate invalid results, as noted in the text.




**Figure 7.** Hall conductivity as a function of latitude from 50° to 75°, at 4 hours MLT. The format of the graph is the same as in Fig. 6.




**Figure 8.** Pedersen conductivity as a function of latitude from  $50^\circ$  to  $75^\circ$ , at 20 hours MLT. The format of the graph is the same as in Fig. 6, with only a change in the MLT.





**Figure 9.** Hall conductivity as a function of latitude from 50° to 75°, at 20 hours MLT. The format of the graph is the same as in Fig. 6 to 8.



**Figure 10.** Dot product of electric field and divergence-free current, for the same conditions as in Fig. 1. The IMF  $B_T$  magnitude is 10 nT at  $180^\circ$ , the solar wind velocity is  $450 \text{ km s}^{-1}$ , the  $F_{10.7}$  index 160 sfu, and the dipole tilt angle is  $0^\circ$ . The integrated sum of all negative values is indicated in the upper left corner, and this total for all positive values in indicated in the upper right corner, in units of GigaWatts.