

Intermediate Layers responses to Geomagnetic Activity During the 2009 Deep Solar Minimum Over the Brazilian Low Latitude Sector

Ângela M. Santos¹, Christiano G. M. Brum², Inez S. Batista¹, José H. A. Sobral¹, Mangalathayil A. Abdu¹,
and Jonas R. Souza¹

¹National Institute for Space Research, São José dos Campos, Brazil.

²Arecibo Observatory, University of Central Florida, Arecibo, Puerto Rico.

Correspondence to: Ângela Santos (angelamacsantos@gmail.com; angelasantos_1@yahoo.com.br)

Abstract. This work presents the daytime behaviour of the Intermediate Layers (*ILs*) parameters (the virtual height - $h'IL$, and the top frequency - $ftIL$) over the low latitude region of Cachoeira Paulista (CP, 22.42° S; 45° W, I: -34.4°) for the deepest solar minimum of the last 100 years. In such a unique condition, this research reveals the *ILs*' quiet state seasonal behaviour as well as its responses to moderate changes in the geomagnetic activity. The main results show that even small variations of geomagnetic activity (quantified by the planetary Kp index) are able to modify the dynamics of the *ILs* parameters. For the first time, it was observed that during the summer, the $h'IL$ decreases rapidly with the increase of geomagnetic activity, mainly in the early morning hours, while in the following hours, a smoothed rise of the *IL* was found in all seasons analysed. Regarding to the *IL* frequency, it was observed that after 12:00 LT, there is a tendency to decrease with the increase of the magnetic disturbances, being this characteristic more intense after 16:00 LT for summer and winter. For the equinox, such variation was detected, however with half of the amplitude of the other seasons. In addition, it stands out the domain of the annual periodicity of the $ftIL$ while the $h'IL$ presents a semiannual component under the geomagnetic quiet condition.

1 Introduction

The deep solar minimum of the solar cycle 23/24 provides an unprecedented opportunity to understand the variability of Earth's ambient ionosphere. During this period, an unusually inactive state

of the Sun with only relatively small sunspot-carrying active regions was observed. The solar fluxes (UV, EUV, and X-rays) responsible for the heating of the upper atmosphere and production of the ionosphere and the well-known 10.7 cm solar radio flux (F10.7cm) presented very low values when compared to the previous solar cycle of the modern era (see for example Balan et al., 2012 and Kutiev et al., 2013). During this period, reduction of ionospheric temperatures and densities were detected over several latitudes (Coley et al., 2010; Heelis et al., 2009; Yue et al., 2010; Klenzing et al., 2011; Aponte et al., 2013). The thermospheric total mass density from the prolonged minimum in solar activity between cycles 23 and 24 with that of the previous solar minima, presented a reduction of about 10–30% compared with the climatologically expected levels (Emmert et al. 2010). Likewise, Heelis et al. (2009) and Aponte et al. (2013) reported an unprecedented contraction of the topside ionosphere to altitudes never reported before.

Great efforts have been made to better understand the behaviour of the different ionospheric layers over the equatorial and low latitudes sectors during this period. Under such condition, it is expected that the effects caused by geomagnetic activities are highlighted, since the variability of radiation coming from the Sun in this case can be neglected. Liu et al. (2012), for example, discussed the impacts of the high-speed stream in the equatorial ionization anomaly (EIA) development. They showed that the inhibition in the EIA formation was probably due to a westward disturbance dynamo electric field. Santos et al. (2016) investigated the behaviour of the equatorial F region zonal and vertical plasma drifts over Jicamarca during the weak geomagnetic storm of June 2008. Based on a realistic low-latitude ionospheric model (SUPIM - Sheffield University Plasmasphere-Ionosphere Model), they showed that the perfect anti-correlation between the vertical and the zonal drifts close to the evening prereversal enhancement of the zonal electric field was driven mainly by a vertical Hall electric field induced by the primary zonal electric field in the presence of an enhanced nighttime E region ionization (see Abdu et al., 1998). Sreeja et al. (2011), in their turn, showed that the daytime E-region westward drift over Trivandrum (8.5°N , 77°E ; dip latitude $\sim 0.5^{\circ}\text{N}$) presented a reduction that was simultaneous with the disappearance of the equatorial sporadic E layer (E_{sq}) echoes in the ionograms. In this case, it was suggested that an additional overshielding electric field (westward/eastward during the day/night), superposed on the ionosphere during the storm main phase, contributed to the observed reduction in the drift.

While the effects of the geomagnetic storms on the E , F , and sporadic- E (Es) layers are widely
 55 investigated, little information can be found about such effects on those layers located in the ionospheric
 valley, especially during the deep solar cycle minimum of 2008-2009. These layers, which are known as
 “intermediate layers (or just ILs)” are defined as a region of enhanced electron density located in the
 ionospheric valley that extends from the peak altitude of the daytime E -region to the bottom side of the
 F -region. As the behaviour of the ILs is little discussed in the literature, and in order for the reader to
 60 become familiar with the subject that this paper is going to deal with, we will first present [a brief summary
 of the main characteristics of the \$ILs\$ over the equatorial and low latitudes sectors over Brazil, and then
 we will be able to understand how the magnetic activity can affect these layers.](#)

Recently, dos Santos et al. (2019) and Santos et al. (2020; 2021) have studied the essential
 characteristics of the ILs over the Brazilian sector during epochs of minimum and maximum solar activity.
 65 It was observed that these layers are predominantly diurnal and present a typical downward movement
 that can last from minutes to hours. Depending on the height at which the ILs are formed, they can descend
 and merge with the normal ongoing sporadic - E (Es) layers. The ILs ’ occurrence over Brazil is high and
 seems to be dependent on the magnetic inclination angle and independent (or weakly dependent) on the
 solar activity. [Nocturnal \$ILs\$ also were observed over Brazil but they are very unusual.](#) Regarding the
 70 shape in which the ILs are seen in the ionograms, it was verified that they presented a curved format
 similar to the “h” type Es layer, however the ILs with a straight format and spreading base appearance
 also were observed.

The studies conducted so far on the ILs over Brazil give us some indications that the dynamics of
 these layers can be influenced by the atmospheric tides, gravity waves, and electric fields (Nygrén et al.,
 75 1990; Wilkinson et al., 1992). The day-to-day variability in the average ILs ’ descent velocity also
 suggests the influence of a periodic perturbation with a periodicity of some days. The velocity values
 found are compatible with those of the semidiurnal and quarter-diurnal tides. However, the larger
 descending rate (> 10 km/h) observed over the equatorial region may reveal the additional influence of
 the gravity waves in IL ’s dynamics. Additionally, Santos et al. (2021) reported interesting events in which
 80 the ILs presented an upward movement at the same time in which the F layer rises due to the evening

prereversal enhancement of the zonal electric field. Such characteristic was observed in most of the cases during a period of high solar activity, between October and April months, however a single case also was observed in 2009. Another interesting characteristic observed is that the *ILs* could suffer in some way the influence of the prompt penetration electric fields. Dos Santos et al. (2019), for example, showed a case in which a daytime *IL* over the equatorial region of São Luis (2°S; 44°W) on October 9, 2009, presented a strong upward movement that carried the *IL* to the base of the *F2* layer in ~ 1.5 hours. This anomalous rise was probably caused by the joint action of the atmospheric gravity wave propagation and the dawn to dusk PPEF. Santos et al. (2021) also reported the ascending *ILs*, however, during sunset times. As mentioned by the authors, it is possible that the *ILs* in these cases had been caused by the action of the PRE and in some events by the additional contribution from the prompt penetration electric fields. In all the studied events, the *ILs* were located at altitudes higher than or equal to 175 km, except the event of November 10, 2003, when an *Es* layer located at about 120 km of altitude presented an abrupt rise reaching 290 km of altitude in a time interval of ~ 1.25 hours. This rapid rise of the *Es/IL* layers probably was caused by an eastward electric field of ~ 0.6 mV/m arising from the PRE and the PPEF (for more details, see Santos et al., 2021).


The focus of this paper is to investigate the geomagnetic activity effects on the intermediate layers over the Brazilian low latitude sector during the deep solar minimum of 2009, regardless of the reasons why such storms were generated. As mentioned previously, this epoch is especially suited to develop studies like the one proposed here due to the very low values and little variation of the solar decimetric flux (10.7cm). In this case, the effects caused in the *ILs* by the variability of radiation coming from the Sun can be neglected, and consider only those caused by geomagnetic variations. The data and methodology used to investigate the possible influence of the geomagnetic storms in the intermediate layers is given in Section 2. The results are presented in Section 3 and finally, in Section 4, the discussion and conclusions.

2. Data set

In this paper, the ionospheric sounding data collected by the Digisonde operated over the low latitude site, Cachoeira Paulista (CP, 22.42° S; 45° W, I: -34.4°), during the deep solar minimum of 2009 are used to verify the possible dependence of the *ILs* on the geomagnetic activity. The ionospheric survey made by the Digisonde is based on the reflection of the electromagnetic signal transmitted vertically to the ionosphere with a peak power of the order of 10kW (for the case of Digisonde DGS256, that is the model used to collect data for 2009 over CP) at frequencies ranging from 0.5 to 30 MHz. The vertical radio sounding makes use of the fact that radio waves are reflected in the ionosphere at the height where the local cut-off frequency equals the frequency of the radio wave. The ionospheric information is recorded in the form of ionograms that display the virtual height of the returned echoes versus their frequency, generally registered at 10 and/or 15-min intervals. The Digisonde data used in this work were pre-processed through the ARTIST software (Automatic Real Time Ionogram Scaler with True Height) and also manually post-processed using the SAO-explorer software following the same criteria described by Dos Santos et al. (2019). For more details about Digisonde, see for example Reishish (1986) and Reishish et al. (2009). The *ILs*' virtual height ($h'IL$) and top frequency (f_{tIL}) are analysed as a function of the Kp index. All the observed *ILs* were included in the analysis, regardless if they present a descending or an ascending movement.

Before going into detail on the topic that this work proposes, we will first give an overview of the behaviour of *ILs* on the sector of CP. Figure 1 shows the variability of the parameter of frequency and height of the *ILs* (panels **a** and **b**), their distribution with the local time (panel **c**), as well as their rate of occurrence for different seasons of the year. The red, blue and grey colors are used to represent the summer (December solstice), winter (June solstice) and equinoxes, respectively. It can be observed that in general the *ILs* attain higher frequencies (> 6 MHz) after 11:00 LT (panel **a**) (except in some cases), and present a high variability in height during all the analysed period (panel **b**). The downward movement of the *ILs* is one of the important characteristic that also can be observed in panel **b**. As indicated in panel **c**, the occurrence of the *ILs* along the day is not continuous, which means that the *ILs* can appear and disappear many times during the day or simply not occur. Additionally, panel **c** also shows that in some periods (especially in winter, with some exceptions), there is a tendency that the *ILs* be formed a little later (after 08-09 LT). Regarding panel **d**, it can be seen that the *ILs* occurrence in the low latitude sector

of CP increases significantly in the first hours of the day attaining its maximum at ~14:00 LT in summer,
 135 ~12:00 LT in winter, ~10:30 LT in the equinoxes. In general, the probability of occurrence decays
 drastically as the night-time period approaches (Note: the seasons were equally divided, i.e., 121 days
 around the solstices and 61 days around the equinoxes).



140

Figure 1: Behaviour of the frequency (f_iL) and height (h_iL) parameters of the intermediate layers over Cachoeira Paulista during 2009 (panels a and b, respectively) in function of local time; distribution of the ILs occurrence with the local time in function of day of the year for 2009 (panel c), and; the seasonally occurrence probability of the ILs . In each panel, red, blue and grey are used to represent Summer, Winter and Equinoxes, respectively (panel d).


145

Figure 2 summarizes the geophysical condition of the data distribution according to the solar and geomagnetic activities based on the F10.7P index and Kp_{av} index, respectively. The F10.7P (grey line in top left panel) is a combination of the daily decimetric solar flux index (F10.7) and one more term (F10.7A), which corresponds to the average of the 81 previous days, thus $F10.7P = (F10.7A + F10.7)/2$

(given in Solar Flux Units (SFU); $1 \text{ SFU} = 10^{-22} \text{ W}/(\text{m}^2 \text{ Hz})$). F10.7P was chosen because several authors
 150 have shown that the ionospheric parameters are better described by this index (Brum et al., 2011 and
 2012; Goncharenko et al., 2013 and references therein). In fact, Brum et al. (2011) and Brum et al. (2012)
 have shown that the best description of the UV-EUV (based on UV-EUV irradiance data from Pioneer
 Venus Orbiter (10–150 nm) and by the Solar EUV Monitor on board the Solar Heliospheric Observatory
 (26–34 nm and 0.1–50 nm bands)) is given by F10.7P when compared with F10.7. In addition, their works
 155 have shown that the UV-EUV emissions tend to increase with F10.7P until a certain threshold (around
 175 SFU). However, for low solar activity, the UV-EUV variations with the F10.7P can be well
 represented by a linear function and this feature is very important for the methodology employed in this
 work, as seen below. [For more details about F10.7 index, see Tapping \(2013\) and Tapping and Morgan](#)
[\(2017\)](#). The K_{pav} (grey line in left bottom panel) is the average of the 3 hours data current K_p value
 160 ($K_{p(ref)}$) and the previous 3 and 6 hours, that is, $K_{pav} = (K_{p(ref)} + K_{p(ref-3)} + K_{p(ref-6)})/3$, which gives the
 standard behaviour of the geomagnetic activity and avoid sharp gradients in the temporal edges of this
 index (every 3 hours). Then, in the case of K_{pav} , different values can be defined per day, since the ILs can
 occur in different intervals of the day.

The occurrence number in hours of the K_{pav} level during 2009 is presented on the right bottom
 165 panel of Figure 2 (red bars). It is observed that all of the data were acquired during very low to normal
 geomagnetic activity ($K_{pav} \leq 3^+$ or 3.3) according to the Wrenn et al. (1987) classification. Such
 distribution is very similar to that found by Terra et al. (2020) when the authors analyzed the MSTID
 events for the period starting in the middle of 2018 to the end of 2019 (also low solar activity). Note that
 the occurrence of various levels of magnetic activity is well distributed throughout the year (left bottom
 170 panel of Figure 2) and this behaviour is the optimum condition for the kind of analysis of this work, as
 will be seen in this report. [Tsurutani et al. \(2011\) have studied part of the period in analysis and showed](#)
[that the causes of the low geomagnetic activity during the end of cycle #23 can be related to the solar](#)
[midlatitude small coronal holes, low IMF Bz variances, low solar wind speeds, and low solar magnetic](#)
[fields](#). Regarding the solar activity, the period that encompasses our dataset is the end of solar cycle #23
 175 and the beginning of solar cycle #24. A growth of activity and fluctuations of F10.7P along the year is

observed, varying from 66.5 SFU to 78.1 SFU (average of 70.1 SFU, top right panel) and an uneven distribution of F10.7P (left upper panel) may be noted. Schrijver et al. (2011) showed that in agreement with the yearly-averaged sunspot number, only 5 of 28 cycles since 1700 had a minimum lower than in early 2009. From mid-2008 until 2009/09, the fraction of spot-free days fluctuated around 82%,
 180 unprecedented in the age of modern instrumentation. Using Johann Heinrich Müller's sunspot observations from 1709, Carrasco et al. (2021) concluded that one of the most active years in the Maunder Minimum (1709), was still less active than most years in the Dalton Minimum and also less active than those of the most recent solar cycles. As commented by the authors, the solar activity level in 2009 was similar to that in 1709 according to its most probable value. All the characteristics mentioned above
 185 reinforce the importance of the period chosen here to analyse the possible dependence of ILs on geomagnetic activity since 2009 can be, in some way, comparable to the Maunder Minimum epoch of greatly suppressed solar activity and considered as the weakest cycle of the past 100+ years (Zharkova, 2020).



190

Figure 2: Variability of the solar and geomagnetic activity quantified by the F10.7P and the Kp_{av} indices (upper and bottom block of panels, respectively) for the studied period. Left panels show the geophysical conditions as a function

of the day of the year while the right panels their corresponding number of occurrences (in hours) under different geophysical conditions. The dots of the left upper panel represent the 41 days' averages and the standard deviation of F10.7P, while dots of the left bottom panel represent the same range of days of the upper panel and its respective standard deviation, but for $Kp_{av} \leq 2.3$ (geomagnetic condition used to construct the quiet time condition of $h'IL$ and $ftIL$) (Note: Left panels show the geophysical conditions as a function of the day of the year (grey line)). The red continuous lines are the reconstruction of these variabilities using Fast Fourier Transform (FFT). The given occurrence in the right column of panels is the number of hours for a given interval of Kp_{av} (0.125) and F10.7P (0.5 SFU).

200

From the $h'IL$ and $ftIL$ data, an empirical climatological model was developed that accounted for the dependences of these parameters on time and season, under low solar and geomagnetic activities. Determining the variability of ILs ' parameters in function of time and season make it possible the isolation of any changes related to geomagnetic activity. The first step in our methodology was to extract the seasonal quiet time behaviour of the $h'IL$ and $ftIL$ parameters. For this, it was employed the weighted arithmetic mean defined as $\bar{x}(t_{ref}, d)$ represented as (equation 1):

$$\bar{x}(t_{ref}, d_{ref}) = \frac{\sum_{d_{ref}-20}^{d_{ref}+20} [x(t_{ref}, d) (|d_{ref}-d|)]}{\sum_{d_{ref}-20}^{d_{ref}+20} (|d_{ref}-d|)} \quad (1)$$

where x denotes $h'IL$ or $ftIL$ values under the geomagnetic activity condition below $Kp_{av} \leq 2.3$ for the time reference t_{ref} and the selected day of the year ($d=DOY$). The average value of height and frequency of the ILs was calculated considering 20 days adjacent to the d_{ref} and 30 minutes around the t_{ref} .

210

From the quiet time variability of the $h'IL$ and $ftIL$ obtained by weighted arithmetic mean process described above, a simple model was built using finite Fourier series reconstruction following the procedure by Souza et al. (2010) and Brum et al. (2011), given by

$$xV_{(t,d)} = A0_{(t)} + 2 \sum_{m=1}^4 [Am_{(t)} \cos(2\pi m f_1 d) + Bm_{(t)} \sin(2\pi m f_1 d)] \quad (2)$$

215 where $xV_{(t,d)}$ is the reconstructed variable as a function of time in LT (t) and DOY (d) (xV stands for $h'IL$ or $ftIL$), f_1 is the fundamental frequency of the parameter to be reconstructed ($1/365$), $A0_{(t)}$ is the annual average of the such parameter for a given (t), and finally, $Am_{(t)}$ and $Bm_{(t)}$ are the m^{th} Fourier coefficients also as a function of time. The terms $A0_{(t)}$, $Am_{(t)}$ and $Bm_{(t)}$ were incorporated to the model using polynomial fittings in function of time (LT), as shown in Figure 3, for the harmonics $m=1$ (one year),
 220 $m=2$ (~6 months), $m=3$ (~4 months) and $m=4$ (~3 months). The upper left ($h'IL$) and right ($ftIL$) panels show the time dependence of $A0_{(t)}$ open circles for Fourier coefficients. Similarly, the values of $Am_{(t)}$ and $Bm_{(t)}$ are presented in the lower panels by the blue and red circles, respectively. In all the panels of this figure, the best polynomial fitting is represented by the continuous lines following the same colour scale described above.

225




Figure 3: Dependence of the $h'IL/fiIL$'s FFT coefficients as a function of LT (left/right, respectively). The circles are the values obtained by the FFT decomposition, while the continuous lines are the best polynomial approximation (more information in the manuscript body). The colours blue and red are used to represent the coefficients values of $Am(t)$ and $Bm(t)$, respectively.

230

Based on the model output above described, Figure 4 shows the behaviour of the $h'IL$ and $fiIL$ during the year from 06:00 to 18:00 LT (top and bottom panel, respectively). The right panels show the dispersion diagram between the model and its respective weighted arithmetic mean (under $Kp_{av} \leq 2.3$)

obtained by equation 1, wherein it is possible to see the good correlation between the observation and the modeled data. The left panels show the dominance of the semi-annual and annual variation of the ILs' virtual height and top frequency, respectively. It is interesting to observe that the upper intermediate layers (>160 km) are formed as winter approaches in southern hemisphere between $\sim 06:00$ LT and $11:00$ LT, with a maximum in April/May (DOY 92-153) before the local noon. A second maximum is observed from the beginning of November to middle of January (DOY 304-15), however, in a more restricted range of time (prior $\sim 09:00$ LT). After $12:00$ LT, the ILs are generally located at altitudes at or below to 150 km. In addition, it is observed that the evolution of the ILs to altitudes below 120 km was more evident between the months of April and May (DOY 92-153) at the end of the day. The bottom left panel shows an annual variation of the top frequencies, with a maximum at about $12:00$ - $13:00$ LT from November to February (DOY 304-62). It can be observed that the upper ILs present lower frequencies when compared to the layers located near 150 km. As the ILs descend, they can reach the E region and merge with the existing sporadic- Es layer increasing, in this way, the top frequency of the layer due to the presence of the metallic ions.




Figure 4: Contour plot of the annual variation of the modeled virtual height ($h'IL$, left top panel) and top frequency ($f'IL$, left bottom panel) of the intermediate layers over Cachoeira Paulista. The dispersion diagrams on the right hand-side show the correlation between the weighted arithmetic mean and model results. In each panel on the right, the Correlation Coefficient (R) and Root Mean Square Error ($RSME$) values are also provided.

Figure 5 exemplify how the dependence of the different parameters of the intermediate layers in respect to geomagnetic activity was investigated in this work using ΔKp_{av} . The ΔKp_{av} is the mean of the respective Kp_{av} (grey line in the left bottom panel of Figure 2, $Kp_{av} = (Kp_{(ref)} + Kp_{(ref-3)} + Kp_{(ref-6)})/3$) minus the average of any value below $Kp_{av} \leq 2.3$ in a range of ± 20 days (this is the geomagnetic condition that the model was developed, the red line shown in the left bottom panel of Figure 2). Noticed: The usage of the residuals minimizes the background quiet time behaviour variation along the time (LT and season), enhancing this way the detection of the real contribution or not of the geomagnetic activity on the ILs parameters. The upper panel of Figure 5 shows the whole dataset sorted from the lowest to the highest ΔKp_{av} values and divided into eight sections with the same percentage of samples for each range of ΔKp_{av} (12.5%, represented by the black vertical lines) for the summer (December solstice) at 17:30 LT ± 30 minutes. Specifically, for this example, the selected range represents 178 data points, i.e., each 12.5% displays the behaviour of ~ 22 individual data samples. This panel also displays the respective F10.7P values (red line) and its respective average and standard deviation (blue open circles) for the same sorted 12.5% occurrence range of ΔKp_{av} . Note that the F10.7P mean variation for each range does not vary much which leads us to emphasize that the following variations of ILs are due to geomagnetic activity. The bottom panels show the $h'IL$ and $ftIL$ responses to the geomagnetic activity by the residual average obtained by the difference of the data and the model output presented in Figure 4 in function of ΔKp_{av} . The open circles represent the average values of the height/frequency residuals ($\Delta h'IL$ and $\Delta ftIL$, respectively) for the eight different levels of ΔKp_{av} and their respective standard deviations (vertical and horizontal lines). The linear fitting is indicated by the blue lines. The slope (SLP) of the dependence of $h'IL$ and $ftIL$ with respect to the geomagnetic activity variation (km. ΔKp^{-1} and MHz. ΔKp^{-1}) and the correlation factor (R) are also shown. In this example, it can be clearly observed that as the geomagnetic activity increases, the height of the intermediate layers also increases. The opposite occurs with the frequency when an increase of the ΔKp_{av} causes a decrease in this parameter.




Figure 5: Responses of the intermediate layer to the geomagnetic activity for summer at 17:30 LT \pm 30 minutes. The upper panel shows the ΔKp_{av} (see more information in the text) data organized from the lowest to the highest values and divided into eight sections with the same percentage of samples. In addition, the values of F10.7P with respect to ΔKp_{av} and the average of the F10.7P (blue open circles) for each section are also presented. The bottom panels show the linear regression fitting over the height and frequency residual variability relative to the average ΔKp_{av} values.

The same methodology explained in the case of Figure 5 was applied to all the data between 06:00 and 18:00 LT for each season. Figure 6 shows the *ILs*' dependence on the geomagnetic activity in terms of height (first two columns from left to right) and frequency (two columns of the left) for the different seasons of the year (the data were grouped in seasons as shown in panel c of Figure 1). The correlation coefficient (R) of both parameters is also shown at the right column of each block of panels. The variations of the geomagnetic activity presented in this figure were 2.13 ± 0.28 (summer), 2.18 ± 0.24 (equinoxes) and 1.72 ± 0.21 (winter) in Kp index. It is observed that the higher variability in *ILs*' height with geomagnetic activity occurs during the summer period. In this case, the *IL* was located lower than the expected position with the increase of the geomagnetic activity in the beginning of the day. In the following hours, this condition decreases until a moment in which an opposite behaviour occurs, that is, a small rise of the *IL* begins to be observed with the increase of the ΔKp_{av} . Although some fluctuation in the R-value can be observed (mainly during the equinox and winter), there is a tendency that the *ILs*' height increase with the ΔKp_{av} variation in all seasons after $\sim 12:00$ LT, as can be seen by the positive values of $\text{km} \cdot \Delta Kp_{av}^{-1}$. Regarding the behaviour of the frequency, it is noticed that in the summer, the *fIL* parameter presented a tendency of increase with the geomagnetic activity in the beginning of the day ($\sim 0.1 \text{ MHz} \cdot \Delta Kp_{av}^{-1}$), however from $\sim 12:00$ LT on there is a significant decrease in the *fIL* with ΔKp_{av} (mainly after 16:00 LT) as can be confirmed by the negative values of the coefficient correlation (R). During the equinox, the general tendency is that the *fIL* decrease along the day, and during the winter, little or no response of the top frequency to ΔKp_{av} variability can be observed prior 16:00 LT and sharp decrease after the referred period similar to what is observed during the summer.




Figure 6: Geomagnetic activity effects on $h'IL$ and fIL parameters for different seasons. The first two panels (from left to right) show the linear regression of the $h'IL$ as a function of the ΔKp_{av} index over the different times of the day and the correlation coefficient R . The two right panels indicate the same but for the fIL parameter.

4. Discussion and Conclusions.

It is well known that geomagnetic activity can drastically modify the low-latitude ionospheric dynamics. During the last solar minimum, a unique opportunity was available to investigate such dynamics, since the effects of the solar activity, that dominates the temporal variability of ionospheric properties, could be practically disregarded due to very low solar radiation variation. Using Digisonde data from a Brazilian low latitude station, Cachoeira Paulista, we studied the impacts of the geomagnetic activity in the height and top frequency of the intermediate layers during the deep solar minimum of 2009.

The results summarized in Figure 6 revealed, for the first time, that the most expressive response of the ILs over the low latitude region of Brazil to the geomagnetic activity occurred during the early morning hours ($\sim 06:00 - 08:00$ LT) of the summer when the ILs presented a significant variation of their altitude with the increase of the ΔKp_{av} (as indicated by the negative values of $km.\Delta Kp_{av}^{-1}$ in $h'IL$ panels).

One of the hypotheses to explain such variation in the $h'IL$ parameter is that this behaviour can be related to dusk-to-dawn directed PPEF (see for example Tsurutani et al., 2008). Such electric fields have westward polarity during daytime and therefore it may be one of the factors responsible for the occurrence of lower $h'IL$ at this time. As pointed out by Santos et al. (2021), depending on the height at which the IL s are located, the disturbance electric field can affect considerably the vertical displacement of the intermediate layers.

Another interesting point that needs to be considered is that the movement of the IL s can also be influenced by the regular undisturbed day-to-day variations in the zonal electric field of the ionosphere that is directed to east/west during the daytime/nighttime hours. Therefore, it is possible that in the first 2-3 hours of our analysis period, the lowering of the $h'IL$ with the increase of ΔKp_{av} (negative values of the rate $km.\Delta Kp_{av}^{-1}$) could be a result of a competition between the eastward zonal electric field created by the E-region dynamo and the disturbance westward electric field arising from the *overshielding* process. In the following hours, an opposite situation was observed, that is, a small rise of the IL s occurred in all seasonal periods analyzed, as denoted by the positive values of the rate $km.\Delta Kp_{av}^{-1}$ in the first set of panels on the left in Figure 6. As mentioned by dos Santos et al. (2019), the rise of an IL during daytime can also be a result of the joint action of the of the eastward PPEF (*undershielding*) and gravity wave propagation. In the case studied by dos Santos et al. (2019), the rise of the IL also was accompanied by a decrease in their top frequency. Wakai (1967) reported that the height of the intermediate layer over Boulder (40°N; 105°W) also can be influenced by the magnetic disturbances, however their observation was made during the night-time.

At the same time in which a decrease in the IL s height is observed, an increase in the $ftIL$ parameter occurs during the summer between 08:00 and 12:00 LT attaining a maximum of $0.2\text{ MHz}.\Delta Kp_{av}^{-1}$ at 09:00 LT. It is interesting to observe that before 12:00 LT, for example, the rate variation of the $ftIL$ was positive indicating that when they are located in lower altitudes, the IL s' top frequency increased in relation to its quiet time values. This increase in the frequency is expected since as the layer descends, it can suffer an additional increase of ionization arising from the metallic ions that contribute to the ion density in the ongoing sporadic-E (Es) layers. As the IL s presented a rise after 13:00 LT, the tendency was that the $ftIL$

345 decrease with the increase of ΔKp_{av} . Note that the after 16:00 LT, this decreased is more accentuated during the summer and winter. Analyzing the incoherent scatter data from the mid-latitude region of Arecibo, Raizada et al. (2017) showed that the integrated electron content (*E*-region total electron content - ErTEC) between 80 and 150 km altitude regions presented a maximum variability throughout the night due to geomagnetic activity for both low and high solar activity during equinox periods. Besides that, the authors also verified that the integrated electron content during geomagnetically disturbed/normal conditions and high solar flux periods displays positive changes during summer and equinox, while it is negative in winter. Wakai (1967) reported a study about the maximum electron concentration of the nighttime *E* layer, the valley above it, and the appearance of the intermediate layer from analysis of the low-frequency ionogram obtained at Boulder on three nights of quiet, moderate, and severe geomagnetic activity. They observed an increase of the ionization in the nighttime valley at times of increased magnetic activity and the appearance of an intermediate layer in ~ 150 -160 km during periods of moderated activity. Eventually, the *IL* can be impacted by the energetic particle precipitation (EPP) (see for example Santos et al., 2016a, b), mainly during the occurrence of intense geomagnetic storms. Furthermore, as the present study refers to a period in which the geomagnetic storms were considerably weaker. That said, we believe that if *ILs* were impacted in any way by the EPP, it would not be relevant to our investigation at this moment. In addition, theoretical simulation of ion-pair production by EPP over Cachoeira Paulista have shown that the peak production of electrons is comfortable below the *IL*'s minimum height used in this work (Brum et al., 2006; Brum et al., 2021).

The effects of the magnetic storms on the intermediate layer were studied also by Rodger et al. (1981) using ionosonde data from South Georgia (54°S; 34°W). They showed that the rate of the downward movement and the final height of the nocturnal intermediate layer are independent of the season or magnetic activity. Additionally, they observed that the probability of formation of an *IL* when the minimum virtual height of the *F2* layer is above 220 km is very low, but it can increase during magnetically disturbed periods. As shown by Santos et al. (2020), the *ILs*' occurrence over Cachoeira Paulista (22.42° S; 45° W) was very high both in 2009 (a low solar activity year and the same period of this report) and 2003 (a high solar activity year). These results show that in general, the *ILs* occurrence

resulted to be independent of the magnetic disturbances, since the referred two periods of geomagnetic activity are totally different from each other. However, their development/dynamics over Brazilian sector can be affected by disturbed electric fields, as shown by the results here presented and others previous publications (dos Santos et al. 2019, and Santos et al. 2021).

Although the impacts of the geomagnetic activity on different layers of the ionosphere have been extensively studied, there is a lack of information about what happens in the ionospheric valley region during such conditions, mainly over the low and equatorial latitudes. Using the low-power VHF radar data over the equatorial site of Jicamarca, Chau and Kudeki (2006) showed that the 150-km echoes were not affected by the electric field reversal caused by a magnetic disturbance ($K_p=5$). As mentioned by them, a statistical study on the *ILs* occurrence based on the magnetic activity index K_p did not identify any correlation between magnetic activity and the 150-km echoes. On the other hand, our results show that a small variation in $\Delta K_{p_{av}}$ index (~ 2.0) can affect the *ILs*, especially in the morning period of the summer and late afternoon of all season over the low latitude sector over Brazil. Although the techniques used by us are different from those used by Chau and Kudeki (2006), the contrasting result reveals that the ionospheric valley is a complex region and additional studies need to be performed to understand the physical mechanism governing the generation of the intermediate layers during the occurrence of magnetic disturbances. It is important to emphasize that for the first time it was shown that a small variation in $\Delta K_{p_{av}}$ index (by ~ 2.0) was able to impact the dynamics of the intermediate layer over the low latitude region during the period of deep solar minimum. The main results of this work are summarized as follow:

1. A small variation in the geomagnetic activity during low solar activity can affect both the parameter of height and frequency of the *ILs* over the low latitude Brazilian sector and such responses are dependent on local time and season;
2. During the summer, the height of the *ILs* tend to be lower with the increase of the magnetic activity in the first hours of the day. This characteristic was probably caused by a dusk to dawn electric field;
3. During daytime, the smoothed rise of the *h'IL* can be related to the regular day-to-day undisturbed zonal electric field of the ionosphere;

4. In respect to the top frequency dependence with geomagnetic activity, before 12:00 LT it was
 400 observed positive or null variation in all seasons. After midday, there is a tendency that the *ftIL*'s
 decrease with the magnetic disturbances, being this characteristic more intense after 16:00 LT for the
 summer and winter, and;
5. The domain of a semi-annual and an annual component variation was observed in parameters of
 height and top frequency of the *ILs*, respectively, for geomagnetic very quiet time condition.

405

Data availability. The Digisonde data can be downloaded in Zenodo (identified as CAJ2M 2009 in
<https://doi.org/10.5281/zenodo.3967542>).

Author contributions. AMS and CGMB processed the data, performed the analysis and wrote the paper.
 ISB, MAA, JHAS, JRS contributed in the interpretation of the data.

410 *Competing interests.* The authors declare that they have no conflict of interest.

Special issue statement. From the Sun to the Earth's magnetosphere–ionosphere–thermosphere

Acknowledgments: AMS thanks the financial support from FAPESP (process number: 2015/25357-4)
 and CNPq (165743/2020-4). The Kp index was obtained from the World Data Center for Geomagnetism,
 415 Kyoto (<http://wdc.kugi.kyoto-u.ac.jp/index.html>) and Solar Radio Flux (F10.7 cm) from the National
 Oceanic and Atmospheric Administration (NOOA). ISB thanks CNPq grant numbers 306844/2019-2 and
 405555/2018-0. One of us (JHAS) had Grant number 303383/2019-4 from the Conselho Nacional de
 Desenvolvimento Científico e Tecnológico (CNPq). J. R. Souza would like to thank the CNPq (grant
 307181/2018-9) for research productivity sponsorship and the INCT GNSS-NavAer supported by CNPq
 420 (465648/2014-2), FAPESP (2017/50115-0) and CAPES (88887.137186/2017-00). The Arecibo
 Observatory is operated by the University of Central Florida under a cooperative agreement with the

National Science Foundation (AST-1744119) and in alliance with Yang Enterprises and Ana G. Méndez-Universidad Metropolitana.

425 **References.**

Balan, N., C. Y. Chen, J. Y. Liu, and G. J. Bailey (2012), Behaviour of the low latitude ionosphere-thermosphere system at long deep solar minimum, *Indian J. Radio Space Phys.*, 41, 89– 97.

Brum, C.G.M.; Abdu, M.A.; Batista, I.S.; Carrasco, A.J. and Santos, P.M.T. (2006). Numerical Simulation of Nighttime Electron Precipitation in the Lower Ionosphere over a Sub-Auroral Region. *Advances in Space Research*, v.37, p.1051 – 1057. doi:10.1016/j.asr.2006.02.003.

Brum, C. G. M., F. S. Rodrigues, P. T. dos Santos, A. C. Matta, N. Aponte, S. A. Gonzalez, and E. Robles (2011), A modeling study of foF2 and hmF2 parameters measured by the Arecibo incoherent scatter radar and comparison with IRI model predictions for solar cycles 21, 22, and 23, *J. Geophys. Res.*, 116, A03324, doi:10.1029/2010JA015727.

435 Brum, C. G. M., C. A. Tepley, J. T. Fentzke, E. Robles, P. T. dos Santos, and S. A. Gonzalez (2012), Long-term changes in the thermospheric neutral winds over Arecibo: Climatology based on over three decades of Fabry-Perot observations, *J. Geophys. Res.*, 117, A00H14, doi:10.1029/2011JA016458.

Brum, C. G. M. (2021), The impacts of particle precipitation spectrum on the 30MHz cosmic noise absorption over the under the South Atlantic Anomaly Region, 2021 XXXIVth General Assembly and Scientific Symposium of the International Union of Radio Science (URSI GASS), pp. 1-4, doi: 10.23919/URSIGASS51995.2021.9560414.

Carrasco, V. M. S., Hayakawa, H., Kuroyanagi, C., Gallego, M. C., & Vaquero, J. M. (2021). Strong evidence of low levels of solar activity during the Maunder Minimum. *Monthly Notices of the Royal Astronomical Society*, 504(4), 5199–5204. doi:10.1093/mnras/stab1155.

445 Chau, J. L. and Kudeki, E.: Statistics of 150-km echoes over Jicamarca based on low-power VHF observations, *Ann. Geophys.*, 24, 1305–1310, <https://doi.org/10.5194/angeo-24-1305-2006>, 2006.

- Coley, W. R., R. A. Heelis, M. R. Hairston, G. D. Earle, M. D. Perdue, R. A. Power, L. L. H. B. J. Holt, and C. R. Lippincott (2010), Ion temperature and density relationships measured by CINDI from the C/NOFS spacecraft during solar minimum, *J. Geophys. Res.*, 115, A02313, doi:10.1029/2009JA014665.
- 450 Dos Santos, Â. M., Batista, I. S., Abdu, M. A., Sobral, J. H. A., Souza, J. R., & Brum, C. G. M. (2019). Climatology of intermediate descending layers (or 150 km echoes) over the equatorial and low-latitude regions of Brazil during the deep solar minimum of 2009. *Annales Geophysicae*, 37(6), 1005–1024. <https://doi.org/10.5194/angeo-37-1005-2019>.
- 455 Echer, E., B. T. Tsurutani, and W. D. Gonzalez (2012), Extremely low geomagnetic activity during the recent deep solar cycle minimum, *Proc. Int. Astron. Union*, 7, 200–209, doi:10.1017/S174392131200484X.
- Emmert, J. T., J. L. Lean, and J. M. Picone (2010), Record-low thermospheric density during the 2008 solar minimum, *Geophys. Res. Lett.*, 37, L12102, doi:10.1029/2010GL043671.
- 460 Goncharenko, L. P., V.W. Hsu, C. G. M. Brum, S.-R. Zhang, and J. T. Fentzke (2013), Wave signatures in the midlatitude ionosphere during a sudden stratospheric warming of January 2010, *J. Geophys. Res.: Space Physics*, 118, doi:10.1029/2012JA018251.
- Hady, A. A. (2013), Deep solar minimum and global climate changes. *Journal of Advanced Research*, V4, I3, P.209-214. doi:10.1016/j.jare.2012.11.001.
- 465 Heelis, R. A., W. R. Coley, A. G. Burrell, M. R. Hairston, G. D. Earle, M. D. Perdue, R. A. Power, L. L. H. B. J. Holt, and C. R. Lippincott (2009), Behaviour of the O⁺/H⁺ transition height during the extreme solar minimum of 2008, *Geophys. Res. Lett.*, 36, L00C03, doi:10.1029/2009GL038652.
- Klenzing, J., F. A. Simoes, S. Ivanov, R. A. Heelis, D. Bilitza, R. F. Pfaff, and D. Rowland (2011), Topside equatorial ionospheric density and composition during and after extreme solar minimum, *J. Geophys. Res.*, 116, A12330, doi:10.1029/2011JA017213.
- 470

- Kutiev I, Tsagouri I, Perrone L, Pancheva D, Mukhtarov P, Mikhailov A, Lastovicka J, Jakowski N, Buresova D, Blanch E, Andonov B, Altadill D, Magdaleno S, Parisi M & Miquel Torta J (2013), Solar activity impact on the Earth's upper atmosphere. *J.Space Weather Space Clim.*, 3, A06, doi:10.1051/swsc/2013028.
- 475 Liu, J., L. Liu, B. Zhao, Y. Wei, L. Hu, and B. Xiong (2012), High-speed stream impacts on the equatorial ionization anomaly region during the deep solar minimum year 2008, *J. Geophys. Res.*, 117, A10304,
- Nygrén, T., Lanchester, B. S., Huuskonen, A., Jalonon, L., Turunen, T., Rishbeth, H., & Van Eyken, A. P. (1990). Interference of tidal and gravity waves in the ionosphere and an associated sporadic-E layer. *Journal of Atmospheric and Terrestrial Physics*, 1526, 9, [https://doi.org/10.1016/0021-9169\(90\)90056-](https://doi.org/10.1016/0021-9169(90)90056-S)
- 480 S.
- Raizada, S.; Brum, C.G.M.; D. Mathews, J.D; Gonzalez, C; Franco, E. (2017). Characteristics of nighttime E-region over Arecibo: Dependence on solar flux and geomagnetic variations, *Advances in Space Research*, Volume 61, Issue 7, Pages 1850-1857, <https://doi.org/10.1016/j.asr.2017.07.006>.
- Reinisch, B. W.: New techniques in ground-based ionospheric sounding and studies, *Radio Sci.*, 21, 331–
- 485 341, 1986.
- [Reinisch, B. W., Galkin, I. A., Khmyrov, G. M., Kozlov, A. V., Bibl, K., Lisysyan, I. A., ... & Luo, Y. New Digisonde for research and monitoring applications. *Radio Science*, 44\(01\), 1-15, 2009.](#)
- Rodger, A. S., Fitzgerald, P. H., and Broom, S. M.: The nocturnal intermediate layer over South Georgia, *J. Atmos. Terr. Phys.*, 43,1043–1050, 1981.
- 490 Santos, A. M., M. A. Abdu, J. R. Souza, J. H. A. Sobral, and I. S. Batista (2016a), Disturbance zonal and vertical plasma drifts in the Peruvian sector during solar minimum phases, *J. Geophys. Res. Space Physics*, 121, doi:10.1002/2015JA022146.
- Santos, A. M., M. A. Abdu, J. R. Souza, J. H. A. Sobral, I. S. Batista, and C. M. Denardini (2016b), Storm time equatorial plasma bubble zonal drift reversal due to disturbance Hall electric field over the
- 495 Brazilian region, *J. Geophys. Res. Space Physics*, 121, doi:10.1002/2015JA022179.

- Santos, A. M., Batista, I. S., Sobral, J. H. A., Brum, C. G. M., Abdu, M. A., & Souza, J. R. (2020). Some differences in the dynamics of the intermediate descending layers observed during periods of maximum and minimum solar flux. *Journal of Geophysical Research: Space Physics*, 125, e2019JA027682. <https://doi.org/10.1029/2019JA027682>.
- 500 Santos, Â. M., Batista, I. S., Brum, C. G. M., Sobral, J. H. A., Abdu, M. A., Souza, J. R., &. (2021). F region electric field effects on the intermediate layer dynamics during the evening prereversal enhancement at equatorial region over Brazil. *Journal of Geophysical Research: Space Physics*.
- Scherliess, L., and B. G. Fejer (1997), Storm time dependence of equatorial disturbance dynamo zonal electric fields, *J. Geophys. Res.*, 102, 24,037–24,046, doi:10.1029/97JA02165.
- 505 [Schrijver, C. J., W. C. Livingston, T. N. Woods, and R. A. Mewaldt \(2011\), The minimal solar activity in 2008–2009 and its implications for longterm climate modeling, *Geophys. Res. Lett.*, 38, L06701, doi:10.1029/2011GL046658.](#)
- Souza, J.R., Brum, C.G.M., Abdu, M.A., Batista, I.S., Asevedo, W.D., Bailey, G.J., Bittencourt, J.A. (2010), Parameterized Regional Ionospheric Model and a comparison of its results with experimental
- 510 data and IRI representations, *Advances in Space Research*, 46, 1032-1038, <https://doi.org/10.1016/j.asr.2009.11.025>.
- Sreeja, V., T. K. Pant, L. Jose, and S. Ravindran (2011), Westward electric field penetration to the dayside equatorial ionosphere during the main phase of the geomagnetic storm on 22 July 2009, *J. Geophys. Res.*, 116, A03303, doi:10.1029/2010JA016013.
- 515 [Tapping, K. F. \(2013\). The 10.7 cm solar radio flux \(F10.7\). *Space Weather*, 11\(7\), 394–406. doi:10.1002/swe.20064.](#)
- [Tapping, K., & Morgan, C. \(2017\). Changing Relationships Between Sunspot Number, Total Sunspot Area and F_{10.7} in Cycles 23 and 24. *Solar Physics*, 292\(6\), doi:10.1007/s11207-017-1111-6.](#)
- Terra, P., Vargas, F., Brum, C. G. M., & Miller, E. S. (2020). Geomagnetic and solar dependency of
- 520 MSTIDs occurrence rate: A climatology based on airglow observations from the Arecibo Observatory

ROF. Journal of Geophysical Research: Space Physics, 125, e2019JA027770.
<https://doi.org/10.1029/2019JA027770>.

Tsurutani, B. T., et al. (2008), Prompt penetration electric fields (PPEFs) and their ionospheric effects during the great magnetic storm of 30–31 October 2003, *J. Geophys. Res.*, 113, A05311, doi:10.1029/2007JA012879.

Tsurutani, B. T., Echer, E., and Gonzalez, W. D. (2011). The solar and interplanetary causes of the recent minimum in geomagnetic activity (MGA23): a combination of midlatitude small coronal holes, low IMF BZ variances, low solar wind speeds and low solar magnetic fields, *Ann. Geophys.*, 29, 839–849, <https://doi.org/10.5194/angeo-29-839-2011>

530 Zharkova, V. (2020) Modern Grand Solar Minimum will lead to terrestrial cooling, *Temperature*, 7:3, 217-222, DOI: 10.1080/23328940.2020.1796243.

Zerbo J.L., Amory-Mazaudier C., Ouattara F., 2013. Geomagnetism during solar cycle 23: Characteristics. *J. Adv. Res*, 4(3), 265–274. doi:10.1016/j.jare.2013.08.010.

Wakai, N. (1967), Quiet and disturbed structure and variations of the nighttime E-region, *J. Geophys. Res.*, 72, 4507-4517.

Wilkinson, P. J., Szuszczewicz, E. P., & Roble, R. G. (1992). Measurements and modeling of intermediate, descending, and sporadic layers in the lower ionosphere: Results and implications for global-scale ionospheric-thermospheric studies. *Geophysical Researcher Letter*, 19(2), 95–98. <https://doi.org/10.1029/91GL02774>.

540 Wrenn, G., Rodger, A. S., & Rishbeth, H. (1987). Geomagnetic storms in the Antarctic F-region. I. Diurnal and seasonal patterns for main phase effects. *Journal of Atmospheric and Solar-Terrestrial Physics*, 49(9), 901–913. [https://doi.org/10.1016/0021-9169\(87\)90004-3](https://doi.org/10.1016/0021-9169(87)90004-3)

Yue, X., W. S. Schreiner, J. Lei, C. Rocken, Y. Kuo, and W. Wan (2010), Climatology of ionospheric upper transition height derived from COSMIC satellites during the solar minimum of 2008, *J. Atmos. Terr. Phys.*, 72, 1270–1274, doi:10.1016/j.jastp.2010.08.018.

