

Dayside magnetopause reconnection and flux transfer events: BepiColombo Earth-flyby observations

Weijie Sun¹, James A. Slavin¹, Rumi Nakamura², Daniel Heyner³, Karlheinz J. Trattner⁴, Johannes Z. D. Mieth³, Jiutong Zhao⁵, Qiu-Gang Zong⁵, Sae Aizawa^{6,7}, Nicolas Andre⁶, Yoshifumi Saito⁸

5 ¹Department of Climate and Space Sciences and Engineering, University of Michigan, Ann Arbor, MI 48109, United States

²Space Research Institute, Austrian Academy of Sciences, Schmiedlstraße 6, 8042 Graz, Austria

³Institut für Geophysik und extraterrestrische Physik, Technische Universität Braunschweig, 38106 Braunschweig, Germany

10 ⁴Laboratory for Atmospheric and Space Physics, University of Colorado, Boulder, CO 80303, USA

⁵School of Earth and Space Sciences, Peking University, Beijing 100871, China

⁶Institut de Recherche en Astrophysique et Planétologie, CNRS-UPS-CNES, Toulouse, France

⁷Department of Geophysics, Graduate School of Science, Tohoku University, Sendai, Japan

⁸Japan Aerospace Exploration Agency, Institute of Space and Astronautical Science, Kanagawa, Japan

15 *Correspondence to:* Weijie Sun (wjsun@umich.edu)

Abstract. This study analyzes the flux transfer event (FTE)-type flux ropes and magnetic reconnection around the dayside magnetopause during BepiColombo's Earth flyby. The magnetosheath has a high plasma β (~ 8) and the interplanetary magnetic field (IMF) has a significant radial component. Six flux ropes are identified around the magnetopause. The motion of flux ropes together with the maximum magnetic shear model suggest that the reconnection X-line possibly swipes BepiColombo near the magnetic equator due to an increase of the radial component of IMF. The flux rope with the highest flux content contains a clear coalescence signature, i.e., two smaller flux ropes merge, supporting theoretical predictions the flux contents of flux ropes can grow through coalescence. The coalescence of the two FTE-type flux ropes takes place through secondary reconnection at the point of contact between the two flux ropes. The BepiColombo measurements indicate a large normalized guide field and a reconnection rate comparable to that measured at the magnetopause (~ 0.1).

1. Introduction

Flux transfer events (FTEs) are frequently observed near the outer boundaries, i.e., magnetopause, of planetary magnetospheres, including on Earth (e.g., Russell and Elphic, 1978; Saunders et al., 1984; Wang et al., 2005), Mercury (Russell and Walker, 1985; Slavin et al., 2009, 2010, 2012; Imber et al., 2014; Sun et al., 2020a; Zhong et al., 2020), Saturn (Jasinski et al., 2016, 2021) and Jupiter (Walker and Russell, 1985; Lai et al., 2012). Some of the FTEs have magnetic flux ropes at their cores, which consist of helical magnetic field lines surrounding stronger magnetic fields paralleling their central axes (Paschmann et al., 1982; Lee et al., 1993). These FTE-type flux ropes are created by multiple X-line reconnections in the magnetopause during intervals of significant magnetic shear across the magnetopause current sheet (Lee and Fu, 1985; Raeder, 2006). As a result, the FTE-type flux ropes signal

35 not only the occurrence of magnetic reconnection but their directions of travel can be used to infer the relative location of the reconnection X-lines at the magnetopause.

The FTEs usually include magnetic field lines with one end connecting to the IMF and the other to magnetospheric cusp. They transport magnetic flux from the dayside to the nightside magnetosphere that drives the Dungey cycle in planetary magnetospheres with global intrinsic magnetic fields. Sun et al. (2022) has recently reviewed the

40 contributions of FTE-type flux ropes to the Dungey cycle in dipolar planetary magnetospheres. In Mercury's magnetosphere, FTE-type flux ropes transport the majority (>60%) of the circulated flux (Slavin et al., 2010; Sun et al., 2020a). In contrast, FTE-type flux ropes are estimated to transport only a small portion (<5%) of the circulated flux at Earth (Lockwood et al., 1995; Fear et al., 2017). For the giant outer planetary magnetospheres at Jupiter and Saturn, they appear to transport a negligible magnetic flux (< 1%) for the solar wind-driven portion of their internal

45 convection (Jasinski et al., 2021).

The FTEs on Earth's magnetosphere appear most frequently during periods of the southward interplanetary magnetic field (IMF) when the magnetic shear angle across the magnetopause is larger than 90° (e.g., Rijnbeek et al., 1984; Kuo et al., 1995; Wang et al., 2006). The locations of magnetopause X-lines are closely related to the orientation of the IMF. For example, during the purely southward IMF, reconnection most likely occurs on the

50 magnetopause near the subsolar point (Dungey, 1961). During the purely northward IMF, reconnections occur on the magnetopause tailward of the cusp (Dungey, 1961; Song and Russell, 1992; Shi et al., 2009, 2013; Gou et al., 2016). Magnetic reconnection is also thought to occur at the dayside magnetopause under the strong radial IMF (B_x dominate) (Belenkaya, 1998; Luhmann et al., 1984; Pi et al., 2017; Tang et al., 2013; Toledo-Redondo et al., 2021), but the strong radial IMF conditions are less well studied.

55 Coalescence, which refers to the merging of neighboring flux ropes, is thought to be an important process in space plasma physics (Biskamp and Welter, 1980; Dorelli and Bhattacharjee, 2009; Fermo et al., 2011; Hoilijoki et al., 2017). The merging of flux ropes is associated with secondary reconnection, and changes in magnetic field configuration caused by this secondary reconnection can energize particles, especially electrons (Drake et al., 2006). Furthermore, several studies have suggested that FTE-type flux ropes are initially formed between electron to ion

60 scales. They then grow through coalescence, thereby, increasing their magnetic flux contents and scales (Fermo et al., 2011; Akhavan-Tafti et al., 2018). NASA's Magnetospheric Multiscale (MMS) Mission (Burch et al., 2016) has provided several observations of secondary reconnections between neighboring flux ropes (see, Zhou et al., 2017), between flux rope and Earth's dipole magnetic field (Poh et al., 2019), and between interlinked flux tubes (Øieroset et al., 2016; Kacem et al., 2018).

65 This study investigates FTE-type flux ropes and reconnection at Earth's dayside magnetopause during BepiColombo's flyby on 10 April 2020. The paper is arranged as follows. Section 2 introduces the BepiColombo mission and the measurements during Earth's dayside magnetopause crossing. Section 3 analyzes the distribution of magnetopause reconnection with a strong radial IMF component, and the properties of the flux ropes, including a coalescence event. Section 4 provides a summary of our results.

2. BepiColombo Dayside Magnetopause Crossing

2.1. Spacecraft and Instrumentation

BepiColombo is a joint mission by European Space Agency (ESA) and Japan Aerospace Exploration Agency (JAXA), which consists of two spacecraft named the Mercury Planetary Orbiter (MPO) and Mercury Magnetospheric Orbiter (MMO, or Mio). These spacecraft together aim to carry out detailed investigations of Mercury's interior, surface, exosphere, and magnetosphere (Milillo et al., 2020; Murakami et al., 2020; Benkhoff et al., 2010). The mission made its first planetary flyby maneuver at Earth on 10 April 2020 (Mangano et al., 2021), during which several instruments collected measurements. The MPO and the MMO were attached during the Earth flyby, and therefore, their measurements could be deemed as one observation point. The two spacecraft will be separated when they are scheduled to insert into Mercury's orbit by late 2025 or early 2026.

This study uses measurements collected by the magnetometer (MAG) onboard MPO (Heyner et al., 2021), and the low energy electron by Mercury Electron Analyzer (MEA) (Sauvaud et al., 2010), which is part of the Mercury Plasma Particle Experiment (MPPE) onboard MMO (Saito et al., 2021). The MPO/MAG includes one outboard sensor and one inboard sensor, and it has a sampling rate of 128 Hz. Mio/MEA has a sampling rate of 4 s. The IMF and solar wind conditions are obtained from the OMNI dataset (King and Papitashvili, 2005), which has a time resolution of 1 minute.

2.2. Overview of Magnetosheath and Magnetopause

Figure 1 shows an overview of the dayside magnetopause crossing during BepiColombo's Earth flyby. BepiColombo traveled from the magnetosheath into the dayside magnetosphere. It crossed the magnetopause at a distance of $\sim 4.8 R_E$ (R_E is one Earth radius) dawnward from the subsolar magnetopause, which corresponded to a position of (11.2, -4.8, -0.3) R_E in the Geocentric Solar Magnetospheric (GSM) coordinate. During the 30 minutes interval around the magnetopause crossing ($\sim 00:05$ to $00:35$ UT) analyzed here, the IMF was southward with a strong radial component, i.e., the B_x was the dominant component ($B_x/B_t > 0.7$ in Figure 1h). The average electron density in the magnetosheath was estimated to be $\sim 10 \text{ cm}^{-3}$ based on the onboard-calculated partial moment from Mio/MEA between $00:05$ and $00:28$ UT. The magnetosheath plasma β was high with a value of ~ 8.0 , which was the ratio of the thermal pressure to the magnetic pressure. The thermal pressure in the magnetosheath was calculated by assuming that the pressure balance existed across the dayside magnetopause and the thermal pressure inside the dayside magnetosphere was negligible compared to the magnetic pressure.

3. Magnetopause Reconnections and FTE-type Flux Ropes

3.1. Identification of FTE-type Flux Ropes

The FTE-type flux ropes were identified after the measured magnetic field was rotated into boundary normal coordinates (the LMN coordinate). The minimum variance analysis (MVA) (Sonnerup and Cahill Jr., 1967; Sonnerup and Scheible, 1998) was performed on the magnetic field measurements across the magnetopause current sheet from $00:32:30$ to $00:33:25$ UT to obtain the LMN coordinate. The MVA results produced $L =$

[0.10, 0.24, 0.97] (maximum variance direction), $M = [0.12, 0.96, -0.25]$ (intermediate variance direction), $N = [0.99, -0.14, -0.06]$ (minimum variance direction), and the eigenvalue ratios were $\lambda_{max}/\lambda_{int} \sim 54.3$, $\lambda_{int}/\lambda_{min} \sim 3.9$. The λ_{max} , λ_{int} and λ_{min} are the maximum, intermediate, and minimum eigenvalues. Both of the ratios were larger than 3 indicating that the LMN coordinate of the magnetopause was well determined [Sonnerup & Scheible, 1998]. The FTE-type flux ropes were identified with bipolar signatures in the normal magnetic field (B_N) and clear magnetic field rotation (Russell and Elphic, 1978). The identification of flux ropes also required the signature of a strong magnetic field along their central axis, i.e. the intermediate variance direction (see Figure 2 for an example, and e.g. Slavin *et al.* [2009]; Akhavan-Tafti *et al.* [2018]). Six FTE-type flux ropes were identified in this manner in the magnetosheath just upstream of the dayside magnetopause and marked with green arrows in Figure 1e and listed in Table 1.

The first possible FTE-type flux rope shown in Figure 2 was centered at $\sim 00:11:04$ UT when the IMF clock angle was $\sim 210^\circ$, and B_x/B_t was ~ 0.75 . This flux rope traveled southward as inferred from the polarities of the B_N variation (negative to positive, Figure 2c). The flux rope corresponded to clear enhancement in B_M (Figure 2b) and field rotation in the plane of B_{max} - B_{int} (Figure 2e). However, the enhancement in the B_t strength preceded the reversal in B_N could indicate that the magnetic flux was piled up or this structure was a magnetosheath structure other than a flux rope. About 2 minutes later, the clock angle increased to $\sim 260^\circ$. This IMF orientation persisted for about 12 minutes, during which no FTE-type flux ropes were observed. At $\sim 00:26:06$ UT, the clock angle decreased from $\sim 260^\circ$ to $\sim 210^\circ$ while the ratio of B_x/B_t increased to ~ 0.90 . At this point, 5 FTE-type flux ropes successively appeared up to the point where the magnetopause was crossed. The travelling direction for these 5 flux ropes was inferred to be northward, again based on the B_N variations. The first flux rope traveled southward indicating that the primary magnetopause X-line was initially located northward of the spacecraft. Later, the northward motion of the 5 flux ropes indicated that the primary magnetopause X-line(s) had shifted southward.

3.2. Reconnection X-lines from Maximum Magnetic Shear Model

To further investigate reconnection during BepiColombo's dayside magnetopause traversal, the maximum magnetic shear model (Trattner *et al.*, 2007, 2017) was employed to deduce the location of reconnection X-lines. The magnetic shear angle plots during the intervals centered at 00:09, 00:20, 00:28 UT are shown in Figure 3. Figures 3a and 3b correspond to a distorted feature of the anti-parallel reconnection region, which has recently been termed a "Knee" event (Trattner *et al.*, 2021). The bent shape of the anti-parallel reconnection region is associated with the field line draping in the magnetosheath during the dominant B_x (significantly sunward) component in this period. Figure 3c did not provide the predicted X-line. This was because a continuous X-line along the maximum magnetic shear location was difficult to obtain under the situation of a $B_x/B_t \geq 0.9$, which was due to the lack of comprehensive study on how the significant radial IMF draping around the magnetopause influences magnetic reconnection.

In Figure 3a, BepiColombo was located southward of the predicted X-line. From Figure 3a to Figure 3b, the predicted X-line crossed the location of BepiColombo and was then located to the south of BepiColombo. The

changes in X-line locations from Figures 3a to 3b were due to the IMF clock angle decreasing around 10° together with the B_x/B_t increasing from 0.78 to 0.86.

The travelling directions for the FTE-type flux ropes were consistent with the predicted locations of the reconnection X-line by the maximum magnetic shear model during the changing solar wind conditions for this magnetopause encounter. Figure 3a corresponded to the only southward traveling FTE-type flux rope, while the other five northward traveling FTE-type flux ropes were observed during the conditions shown in Figures 3b and 3c. It needs to note that the FTE-type flux ropes and reconnection exhausts should correspond to strong lateral motion as the predicted X-lines were significantly along ~~with~~ the north-south direction. The reconnection exhausts would correspond to a strong duskward component when the spacecraft was located southward of the X-line and a strong dawnward component when it was northward of the X-line. Although the maximum magnetic shear model faces challenges in determining the draping magnetic field lines in the magnetosheath during the intervals of the dominant B_x component (Trattner et al., 2007, 2012), the model predictions are consistent with our observations during BepiColombo's crossing.

3.3. FTE-type Flux Rope Modeling

This study employed a force-free flux rope model (Kivelson and Khurana, 1995) to fit the FTE-type flux ropes. This flux rope model starts from the periodic pinch solution (Schindler et al., 1973) of Ampere's law ($\nabla \times \vec{B} = \mu_0 \vec{j}$), where \vec{B} is the magnetic field vector, \vec{j} is the current density vector, and μ_0 is the magnetic permeability in vacuum. Kivelson and Khurana (1995) further include the axial magnetic field component (B_{int}) in the periodic pinch solution. The flux rope model introduced by Kivelson and Khurana (1995) does not consider the gradient of the magnetic field along the axis of the flux rope. The self-consistent solution of the flux rope model is

$$\begin{cases} B_{max} = \left(\frac{B_T}{\chi}\right) \sqrt{1 + \epsilon^2} \sinh\left(\frac{x_{min}}{T}\right) \\ B_{int} = \left(\frac{B_T}{\chi}\right) \sqrt{1 + \left(\frac{B_{int0}\chi}{B_T}\right)^2} \\ B_{min} = \left(\frac{B_T}{\chi}\right) \epsilon \sin\left(\frac{x_{max}}{T}\right) \end{cases} \quad (1)$$

In the equation, the x_{min} and x_{max} are the positions in the flux rope along ~~with the directions of~~ \vec{n}_{min} and \vec{n}_{max} . The \vec{n}_{min} , \vec{n}_{int} and \vec{n}_{max} refer to the local coordinate of the flux rope, ~~which are determined from the MVA on the flux rope~~. The T is the vertical scale of the flux rope ~~along with in~~ the \vec{n}_{max} direction and the B_T is the magnetic field intensity near the boundary of the flux rope along ~~with the~~ \vec{n}_{min} direction. The B_{int0} is the B_{int} in the background. The χ is

$$\chi = \epsilon \cos\left(\frac{x_{max}}{T}\right) + \sqrt{1 + \epsilon^2} \cosh\left(\frac{x_{min}}{T}\right) \quad (2)$$

In this equation, the parameter ϵ is associated with the shape of the flux rope, i.e., from flattened to circular profiles. The axial flux content (Φ_{axial}) is calculated by integrating the axial field (B_{int}) over the entire flux rope area,

$$\Phi_{axial} = \int B_{int} dS \quad (3)$$

During the fitting, we assume that the traveling speed of flux ropes was 100 km/s, which corresponds to the average Alfvén speed in the subsolar magnetosheath. The traveling speed is required in calculating the scales and magnetic flux content for the flux ropes. The least-squares of the minimization of the magnetic field differences (X^2) is employed to define the best fit, which is calculated from

$$X^2 = \frac{\sum_{i=1}^{N_{point}} \sum_j^{max,int,min} \left[\left(B_j(i) - B'_j(i) \right) / B_t(i) \right]^2}{N_{point}} \quad (4)$$

where B_{max} , B_{int} , B_{min} , and B_t are the components and magnitude of the measured magnetic fields and B'_{max} , B'_{int} , and B'_{min} are the components of the magnetic fields from the model. The N_{point} is the number of data points. We set up a threshold of $X^2 < 0.1$ to be the successful modeling.

Different from the circular profile of flux ropes resulting from the Lundquist force-free flux rope model (Lundquist, 1950; Burlaga, 1988; Lepping et al., 1990), this force-free model can result in either flattened or circular profiles of flux ropes. We use the semi-minor and semi-major to refer to the flattened features. The semi-major corresponds to the scale of flux rope along ~~with the~~ \vec{n}_{min} direction, which is close to the L direction of the magnetopause. The semi-minor correspond to the scale of flux rope along ~~with the~~ \vec{n}_{max} direction, which is close to the N direction of the magnetopause. This flux rope model is successfully applied for the flux ropes in Earth's plasma sheet (Kivelson and Khurana, 1995), on Earth's magnetopause (Zhang et al., 2008), and in Mercury's plasma sheet (Zhao et al., 2019).

Out of the 6 FTE-type flux ropes, 4 were successfully modeled. As an example, the modeling curves of the flux rope centered at 00:28:13 UT are shown in Figures 4a to 4d. In the figures, the dashed lines overlapping with the solid measured magnetic fields represent the modeling curves from the flux rope model. It can be seen clearly that the two curves are close to each other and this flux rope is well fitted by the model. The modeling results for the 4 flux ropes are summarized in Table 1. The plasma density of $\sim 10 \text{ cm}^{-3}$ in the magnetosheath corresponds to an ion inertial length (d_i) of $\sim 70 \text{ km}$. The two FTE-type flux ropes at 00:26:06 UT and 00:26:26 UT are in the scales of several d_i . The magnetic flux contents of these two flux ropes are small ($\sim 20 \text{ kWb}$). In addition, these two flux ropes correspond to the largest and smallest core fields. The other two FTE-type flux ropes at 00:28:13 UT and 00:30:26 UT are in the scales of more than $10 d_i$. These two flux ropes contain much higher magnetic flux ($\sim 300 \text{ kWb}$ and $\sim 188 \text{ kWb}$). The analysis of the flux rope at $\sim 00:28:13 \text{ UT}$ corresponding to the highest magnetic flux content is shown in the next section. Moreover, the flux ropes at 00:26:06 UT, 00:26:26 UT, and 00:30:26 UT are close to circular profiles with the semi-minor slightly smaller than the semi-major. The flux rope at $\sim 00:28:13 \text{ UT}$ ~~includes the strongest~~ corresponds to the most flattened profile.

3.4. Coalescence Event

Figures 4a to 4d show the magnetic field measurements of the FTE-type flux rope centered at $\sim 00:28:13 \text{ UT}$ in the LMN coordinate. This FTE-type flux rope corresponds to the fifth green arrow counting from the leftside in Figure 1e. Figure 4c showed that the B_N included two successive bipolar signatures, which implied that two smaller scale flux ropes were merging. Indeed, the hodogram in the B_{max} - B_{int} plane in Figure 4f confirmed the field rotations of two flux ropes, named “FR#A” and “FR#B”. Figure 4e further illustrated the merging of the two flux ropes and the

trajectory of BepiColombo. The magenta arrow and shaded region in Figure 4e indicated the possible secondary reconnection between FR#A and FR#B. This FTE-type flux rope with the highest flux content possibly resulted from the coalescence of two smaller-scale flux ropes.

In order to study how well aligned FR#A and FR#B were, we applied the MVA on FR#A from 00:28:03 to 00:28:09 and FR#B from 00:28:09 to 00:28:16, separately. The eigenvalue ratios were $\lambda_{max}/\lambda_{int} \sim 1.91$ and $\lambda_{int}/\lambda_{min} \sim 21.7$ for FR#A. The eigenvalue ratios were $\lambda_{max}/\lambda_{int} \sim 3.34$ and $\lambda_{int}/\lambda_{min} \sim 12.6$ for FR#B. The large values of $\lambda_{int}/\lambda_{min}$ indicated that the \vec{n}_{min} were well determined for both flux ropes. The \vec{n}_{min} was $[-0.20, -0.58, -0.79]$ for FR#A and \vec{n}_{min} was $[0.23, -0.55, -0.80]$ for FR#B. The \vec{n}_{min} were close to each other with a separation angle of 25° . The \vec{n}_{min} obtained for the coalescence event was $[-0.04, -0.49, -0.87]$, which were 12° and 17° away from the \vec{n}_{min} of FR#A and FR#B, separately. The small separations of the \vec{n}_{min} should indicate that FR#A and FR#B were well aligned. It needs to note that the coalescence signature was only observed in this FTE-type flux rope centered at $\sim 00:28:13$ UT. The successive bipolar signatures of the B_N were not found in other 5 FTE-type flux ropes.

3.5. Magnetopause Reconnection and Secondary Magnetic Reconnection

In Figure 5, the properties of the secondary current sheet in the coalescence event and the magnetopause current sheet are studied. For the secondary current sheet, the eigenvalue ratios were $\lambda_{max}/\lambda_{int} \sim 6.4$, $\lambda_{int}/\lambda_{min} \sim 11.0$ resulting from the MVA. Both of the eigenvalue ratios were larger than 3 indicating the local coordinate of the secondary current sheet was well established. The magnetic field measurements of the magnetopause current sheet were shown in the LMN coordinate.

In the reconnecting current sheet, the dimensionless reconnection rate can be determined from the ratio of the normal magnetic field component (B_{normal}) to the reconnecting magnetic field (B_{inflow}) in the inflow region (Sonnerup, 1974; Sonnerup et al., 1981; Fuselier and Lewis, 2011; Phan et al., 2001; Sun et al., 2020b). In the secondary current sheet (Figures 5a to 5d), the B_{normal} was ~ 5 nT, which corresponded to the B_{min} averaged from 00:28:08.8 to 00:28:09.6 UT. Here the average B_t from 00:28:09.8 to 00:28:10.4 UT was taken as the B_{inflow} (~ 36 nT). The dimensionless reconnection rate was ~ 0.14 if the reconnection occurred in the secondary current sheet. Meanwhile, the intensity of the guide field (B_{int} , Figure 5b) was ~ 32 nT across the current sheet, which was ~ 0.89 when normalized to the B_{inflow} . In the magnetopause current sheet, the B_{normal} was 8.3 nT, which corresponded to the averaged B_N from 00:32:56 to 00:33:05 UT (Figure 5g). The B_{inflow} in the magnetosphere side adjacent to the magnetopause was ~ 46.1 nT, which corresponded to the averaged B_t from 00:33:06 to 00:33:15 UT (Figure 5h). Thus, the dimensionless reconnection rate was calculated to be ~ 0.18 . The guide field across the magnetopause was ~ 13 nT (B_M , Figure 5f), which was 0.28 normalized to the B_{inflow} .

However, it is needed to point out that the estimation of reconnection rate based on B_N/B_{inflow} could be imprecise. For example, the uncertainties of the normal direction and the fluctuations in the field strength could influence the accuracy of the reconnection rates. As noted by Sonnerup and Scheible (1998) and Khrabrov and Sonnerup (1998), there were uncertainties in the eigenvectors determined by the MVA, which could be either statistical error or error due to the magnetic structure was not perfectly stationary and one-dimensional. By employing the method

introduced by Khrabrov and Sonnerup (1998), we obtained an uncertainty of ~ 0.93 nT for the B_{normal} of the secondary current sheet and ~ 0.04 nT for the magnetopause current sheet.

However, it was not certain that magnetic reconnection was occurring in the secondary current sheet or the magnetopause current sheet. There was no complimentary evidence for the magnetic reconnection since the measurements from BepiColombo were limited during the Earth flyby. The low energy electron measurements (Mio/MEA) were limited in the field of view and the time resolution was ~ 4 seconds. The MEA could not provide a complete distribution relative to the background magnetic field and its time resolution was much longer than the time scale of the secondary current sheet. Therefore, the conclusions obtained about magnetic reconnection are tentative and further analysis about a similar event is needed, especially those measurements taken from the MMS.

4. Conclusions and Discussions

Our analysis of the subsolar magnetopause observations during BepiColombo's Earth flyby has produced several conclusions.

First, the BepiColombo's dayside magnetopause crossing took place during an interval when magnetosheath had a high plasma β (~ 8) and the IMF had a strong radial component ($B_x/B_t > 0.7$). The traveling of the FTE-type flux rope suggests that the X-line crosses the location of BepiColombo, ~~which~~ Although there is a possibility that the first and only southward travelling FTE-type flux rope is a magnetosheath structure, is in close agreement with the predictions of the maximum magnetic shear model suggest that the X-line crosses the location of BepiColombo as well. The X-line motion is associated with the rotation and the x component increase of the IMF. BepiColombo crosses the magnetopause near the magnetic equator, and 10 April 2020 is close to the spring equinox, which indicates a small Earth's dipole tilt influence. These observations of the possible crossing of the X-line provide clear evidence of magnetic reconnection occurrence near the magnetic equator under a strong radial IMF.

Second, the properties of the FTE-type flux ropes are obtained by employing a force-free flux rope model introduced by Kivelson and Khurana (1995). The FTE-type flux ropes correspond to scales ranging from several d_i to around $20 d_i$, and the FTE-type flux rope with a large scale and the highest magnetic flux content exhibits clear coalescence signatures. These observations strongly support the theories in which the FTE-type flux ropes grow in scales and magnetic flux contents through coalescence.

Third, magnetic reconnection in the coalescence event and the magnetopause current sheet is investigated. The reconnection rate of the secondary reconnection (0.14) is comparable with the reconnection rate of dayside magnetopause (0.18). However, the secondary reconnection corresponds to a larger normalized guide field (0.89) and a magnetopause reconnection (0.28). However, there is no complimentary evidence that magnetic reconnection is occurring in the secondary current sheet and magnetopause current sheet. Therefore, the conclusions about magnetic reconnection are tentative.

The large guide field of the secondary magnetic reconnection during the coalescence observed by BepiColombo is likely a common feature. For example, Zhou et al. (2017) reported a coalescence event with a strong guide field. We suggest that these large guide fields should be included in future simulations, which investigate the particle energizations due to coalescence. The large guide fields may influence the reconnection rate as suggested by

Pritchett and Coroniti (2004) and Ricci et al. (2004), and therefore affect the energization of particles during the coalescence. Furthermore, a recent investigation also suggests that a large guide field might limit the ability of Fermi acceleration during the coalescence (Montag et al., 2017).

Finally, the FTE-type flux rope containing the coalescence signature has a scale of $\sim 20 d_i$. Therefore, the secondary reconnecting current sheet embedded within the FTE-type flux rope is likely with a scale smaller than $20 d_i$. We want to note that the secondary reconnection during the coalescence of flux ropes share some similarities with the electron-only reconnection associated with the magnetosheath turbulence, whose reconnecting current sheet has scales smaller than $10 d_i$ and is accompanied by a large guide field as revealed by MMS measurements (Phan et al., 2018; Stawarz et al., 2019) and simulations (Califano et al., 2020). Therefore, it is likely that the secondary reconnection associated with coalescence is electron-only magnetic reconnection, which certainly deserves a detailed study.

Data availability

The measurements from Mio/MEA and MPO/MAG analyzed in this study are available in the supporting information. The data archiving is underway. Mio/MEA data will be able to be accessed from the AMDA science analysis system (<http://amda.cdpp.eu>) provided by the Centre de Données de la Physique des Plasmas (CDPP) supported by CNRS, CNES, Observatoire de Paris, and Université Paul Sabatier Toulouse. MPO/MAG data will be available from <https://archives.esac.esa.int/psa/#!/Home%20View>. OMNI dataset is available at <https://omniweb.gsfc.nasa.gov/>.

Author contributions

W. J. S. led the work, identified the events, conducted the data analysis of the dataset, and wrote the manuscript. W. J. S., J. A. S., and R. N. jointly designed the work. D. H. and J. Z. D. M. provided knowledge of the MPO-MAG instrument and the MPO-MAG data. S. A. and N. A. provided knowledge of the Mio-MEA instrument and the Mio-MEA data. K. J. T. provided Figure 3 and the relevant descriptions. J. T. Z. performed force-free fittings of the flux ropes. All authors discussed and contributed to the manuscript.

Competing interests

The authors declare no competing interests.

Acknowledgment

The BepiColombo project is supported by ESA and JAXA. W. J. S. and J. A. S. were supported by NASA Grants NNX16AJ67G and 80NSSC18K1137. N.A. and S.A. acknowledge the support of CNES for the BepiColombo

mission. The research at LASP (K. J. T.) is supported by NASA grants NNG04EB99C and 80NSSC20K0688. W. J. S. thanks Dr. Gangkai Poh for helpful discussions.

References.

- 310 Akhavan-Tafti, M., Slavin, J. A., Le, G., Eastwood, J. P., Strangeway, R. J., Russell, C. T., Nakamura, R., Baumjohann, W., Torbert, R. B., Giles, B. L., Gershman, D. J., and Burch, J. L.: MMS Examination of FTEs at the Earth's Subsolar Magnetopause, *J Geophys Res Space Phys*, 123, 1224-1241, 10.1002/2017JA024681, 2018.
- Belenkaya, E. S.: Reconnection modes for near-radial interplanetary magnetic field, *J Geophys Res Space Phys*, 103, 26487-26494, <https://doi.org/10.1029/98JA02270>, 1998.
- 315 Benkhoff, J., van Casteren, J., Hayakawa, H., Fujimoto, M., Laakso, H., Novara, M., Ferri, P., Middleton, H. R., and Ziethe, R.: BepiColombo—Comprehensive exploration of Mercury: Mission overview and science goals, *Planet Space Sci*, 58, 2-20, <https://doi.org/10.1016/j.pss.2009.09.020>, 2010.
- Biskamp, D. and Welter, H.: Coalescence of Magnetic Islands, *Phys Rev Lett*, 44, 1069-1072, 10.1103/PhysRevLett.44.1069, 1980.
- 320 Burch, J. L., Moore, T. E., Torbert, R. B., and Giles, B. L.: Magnetospheric Multiscale Overview and Science Objectives, *Space Sci Rev*, 199, 5-21, 10.1007/s11214-015-0164-9, 2016.
- Burlaga, L. F.: Magnetic clouds and force-free fields with constant alpha, *J Geophys Res Space Phys*, 93, 7217-7224, 10.1029/JA093iA07p07217, 1988.
- Califano, F., Cerri, S. S., Faganello, M., Laveder, D., Sisti, M., and Kunz, M. W.: Electron-Only Reconnection in Plasma Turbulence, *Front Phys*, 8, 10.3389/fphy.2020.00317, 2020.
- 325 Dorelli, J. C. and Bhattacharjee, A.: On the generation and topology of flux transfer events, *J Geophys Res Space Phys*, 114, <https://doi.org/10.1029/2008JA013410>, 2009.
- Drake, J. F., Swisdak, M., Che, H., and Shay, M. A.: Electron acceleration from contracting magnetic islands during reconnection, *Nature*, 443, 553, 10.1038/nature05116, 2006.
- 330 Dungey, J. W.: Interplanetary Magnetic Field and the Auroral Zones, *Phys Rev Lett*, 6, 47-48, 10.1103/PhysRevLett.6.47, 1961.
- Fear, R. C., Trenchi, L., Coxon, J. C., and Milan, S. E.: How Much Flux Does a Flux Transfer Event Transfer?, *J Geophys Res Space Phys*, 122, 12,310-312,327, doi:10.1002/2017JA024730, 2017.
- Fermo, R. L., Drake, J. F., Swisdak, M., and Hwang, K. J.: Comparison of a statistical model for magnetic islands in large current layers with Hall MHD simulations and Cluster FTE observations, *J Geophys Res Space Phys*, 116, 10.1029/2010JA016271, 2011.
- 335 Fuselier, S. A. and Lewis, W. S.: Properties of Near-Earth Magnetic Reconnection from In-Situ Observations, *Space Sci Rev*, 160, 95, 10.1007/s11214-011-9820-x, 2011.
- Gou, X. C., Shi, Q. Q., Tian, A. M., Sun, W. J., Dunlop, M. W., Fu, S. Y., Zong, Q. G., Facskó, G., Nowada, M., Pu, Z. Y., Mailyan, B., Xiao, T., and Shen, X. C.: Solar wind plasma entry observed by cluster in the high-latitude magnetospheric lobes, *J Geophys Res Space Phys*, 121, 4135-4144, <https://doi.org/10.1002/2015JA021578>, 2016.
- 340 Heyner, D., Auster, H. U., Fornaçon, K. H., Carr, C., Richter, I., Mieth, J. Z. D., Kolhey, P., Exner, W., Motschmann, U., Baumjohann, W., Matsuoka, A., Magnes, W., Berghofer, G., Fischer, D., Plaschke, F., Nakamura, R., Narita, Y., Delva, M., Volwerk, M., Balogh, A., Dougherty, M., Horbury, T., Langlais, B., Manda, M., Masters, A., Oliveira, J. S., Sánchez-Cano, B., Slavin, J. A., Vennerstrøm, S., Vogt, J., Wicht, J., and Glassmeier, K. H.: The BepiColombo Planetary Magnetometer MPO-MAG: What Can We Learn from the Hermean Magnetic Field?, *Space Sci Rev*, 217, 52, 10.1007/s11214-021-00822-x, 2021.
- 345 Hoilijoki, S., Ganse, U., Pfau-Kempf, Y., Cassak, P. A., Walsh, B. M., Hietala, H., von Alfthan, S., and Palmroth, M.: Reconnection rates and X line motion at the magnetopause: Global 2D-3V hybrid-Vlasov simulation results, *J Geophys Res Space Phys*, 122, 2877-2888, <https://doi.org/10.1002/2016JA023709>, 2017.
- 350 Imber, S. M., Slavin, J. A., Boardsen, S. A., Anderson, B. J., Korth, H., McNutt Jr., R. L., and Solomon, S. C.: MESSENGER observations of large dayside flux transfer events: Do they drive Mercury's substorm cycle?, *J Geophys Res Space Phys*, 119, 5613-5623, 10.1002/2014ja019884, 2014.
- 355 Jasinski, J. M., Akhavan-Tafti, M., Sun, W., Slavin, J. A., Coates, A. J., Fuselier, S. A., Sergis, N., and Murphy, N.: Flux Transfer Events at a Reconnection-Suppressed Magnetopause: Cassini Observations at Saturn, *J Geophys Res Space Phys*, 126, e2020JA028786, <https://doi.org/10.1029/2020JA028786>, 2021.

- Jasinski, J. M., Slavin, J. A., Arridge, C. S., Poh, G., Jia, X., Sergis, N., Coates, A. J., Jones, G. H., and Waite Jr., J. H.: Flux transfer event observation at Saturn's dayside magnetopause by the Cassini spacecraft, *Geophys Res Lett*, 43, 6713-6723, doi:10.1002/2016GL069260, 2016.
- 360 Kacem, I., Jacquey, C., Génot, V., Lavraud, B., Vernisse, Y., Marchaudon, A., Le Contel, O., Breuillard, H., Phan, T. D., Hasegawa, H., Oka, M., Trattner, K. J., Farrugia, C. J., Paulson, K., Eastwood, J. P., Fuselier, S. A., Turner, D., Eriksson, S., Wilder, F., Russell, C. T., Øieroset, M., Burch, J., Graham, D. B., Sauvaud, J. A., Avakov, L., Chandler, M., Coffey, V., Dorelli, J., Gershman, D. J., Giles, B. L., Moore, T. E., Saito, Y., Chen, L. J., and Penou, E.: Magnetic Reconnection at a Thin Current Sheet Separating Two Interlaced Flux Tubes at the Earth's
- 365 Magnetopause, *J Geophys Res Space Phys*, 123, 1779-1793, 10.1002/2017JA024537, 2018.
- Khrabrov, A. V. and Sonnerup, B. U. Ö.: Error estimates for minimum variance analysis, *J Geophys Res Space Phys*, 103, 6641-6651, <https://doi.org/10.1029/97JA03731>, 1998.
- King, J. H. and Papitashvili, N. E.: Solar wind spatial scales in and comparisons of hourly Wind and ACE plasma and magnetic field data, *J Geophys Res Space Phys*, 110, 10.1029/2004ja010649, 2005.
- 370 Kivelson, M. G. and Khurana, K. K.: Models of flux ropes embedded in a harris neutral sheet: Force-free solutions in low and high beta plasmas, *J Geophys Res Space Phys*, 100, 23637-23645, <https://doi.org/10.1029/95JA01548>, 1995.
- Kuo, H., Russell, C. T., and Le, G.: Statistical studies of flux transfer events, *J Geophys Res Space Phys*, 100, 3513-3519, <https://doi.org/10.1029/94JA02498>, 1995.
- 375 Lai, H. R., Wei, H. Y., Russell, C. T., Arridge, C. S., and Dougherty, M. K.: Reconnection at the magnetopause of Saturn: Perspective from FTE occurrence and magnetosphere size, *J Geophys Res Space Phys*, 117, 10.1029/2011ja017263, 2012.
- Lee, L. C. and Fu, Z. F.: A theory of magnetic flux transfer at the Earth's magnetopause, *Geophys Res Lett*, 12, 105-108, 10.1029/GL012i002p00105, 1985.
- 380 Lee, L. C., Ma, Z. W., Fu, Z. F., and Otto, A.: Topology of magnetic flux ropes and formation of fossil flux transfer events and boundary layer plasmas, *J Geophys Res Space Phys*, 98, 3943-3951, <https://doi.org/10.1029/92JA02203>, 1993.
- Lepping, R. P., Jones, J. A., and Burlaga, L. F.: Magnetic field structure of interplanetary magnetic clouds at 1 AU, *J Geophys Res Space Phys*, 95, 11957-11965, 10.1029/JA095iA08p11957, 1990.
- 385 Lockwood, M., Cowley, S. W. H., Smith, M. F., Rijnbeek, R. P., and Elphic, R. C.: The contribution of flux transfer events to convection, *Geophys Res Lett*, 22, 1185-1188, 10.1029/95gl01008, 1995.
- Luhmann, J. G., Walker, R. J., Russell, C. T., Crooker, N. U., Spreiter, J. R., and Stahara, S. S.: Patterns of potential magnetic field merging sites on the dayside magnetopause, *J Geophys Res Space Phys*, 89, 1739-1742, <https://doi.org/10.1029/JA089iA03p01739>, 1984.
- 390 Lundquist, S.: Magnetohydrostatic fields, *Ark. Fys.*, 2, 361-365, 1950.
- Mangano, V., Dósa, M., Fränz, M., Milillo, A., Oliveira, J. S., Lee, Y. J., McKenna-Lawlor, S., Grassi, D., Heyner, D., Kozyrev, A. S., Peron, R., Helbert, J., Besse, S., de la Fuente, S., Montagnon, E., Zender, J., Volwerk, M., Chaufray, J.-Y., Slavin, J. A., Krüger, H., Maturilli, A., Cornet, T., Iwai, K., Miyoshi, Y., Lucente, M., Massetti, S., Schmidt, C. A., Dong, C., Quarati, F., Hirai, T., Varsani, A., Belyaev, D., Zhong, J., Kilpua, E. K. J., Jackson, B. V.,
- 395 Odstrcil, D., Plaschke, F., Vainio, R., Jarvinen, R., Ivanovski, S. L., Madár, Á., Erdős, G., Plainaki, C., Alberti, T., Aizawa, S., Benkhoff, J., Murakami, G., Quemerais, E., Hiesinger, H., Mitrofanov, I. G., Iess, L., Santoli, F., Orsini, S., Lichtenegger, H., Laky, G., Barabash, S., Moissl, R., Huvelin, J., Kasaba, Y., Saito, Y., Kobayashi, M., and Baumjohann, W.: BepiColombo Science Investigations During Cruise and Flybys at the Earth, Venus and Mercury, *Space Sci Rev*, 217, 23, 10.1007/s11214-021-00797-9, 2021.
- 400 Milillo, A., Fujimoto, M., Murakami, G., Benkhoff, J., Zender, J., Aizawa, S., Dósa, M., Griton, L., Heyner, D., Ho, G., Imber, S. M., Jia, X., Karlsson, T., Killen, R. M., Laurenza, M., Lindsay, S. T., McKenna-Lawlor, S., Mura, A., Raines, J. M., Rothery, D. A., André, N., Baumjohann, W., Berezhnoy, A., Bourdin, P. A., Bunce, E. J., Califano, F., Deca, J., de la Fuente, S., Dong, C., Grava, C., Fatemi, S., Henri, P., Ivanovski, S. L., Jackson, B. V., James, M., Kallio, E., Kasaba, Y., Kilpua, E., Kobayashi, M., Langlais, B., Leblanc, F., Lhotka, C., Mangano, V., Martindale, A., Massetti, S., Masters, A., Morooka, M., Narita, Y., Oliveira, J. S., Odstrcil, D., Orsini, S., Pelizzo, M. G.,
- 405 Plainaki, C., Plaschke, F., Sahraoui, F., Seki, K., Slavin, J. A., Vainio, R., Wurz, P., Barabash, S., Carr, C. M., Delcourt, D., Glassmeier, K. H., Grande, M., Hirahara, M., Huvelin, J., Korabely, O., Kojima, H., Lichtenegger, H., Livi, S., Matsuoka, A., Moissl, R., Moncuquet, M., Muinonen, K., Quémérais, E., Saito, Y., Yagitani, S., Yoshikawa, I., and Wahlund, J. E.: Investigating Mercury's Environment with the Two-Spacecraft BepiColombo
- 410 Mission, *Space Sci Rev*, 216, 93, 10.1007/s11214-020-00712-8, 2020.
- Montag, P., Egedal, J., Lichko, E., and Wetherton, B.: Impact of compressibility and a guide field on Fermi acceleration during magnetic island coalescence, *Phys Plasmas*, 24, 062906, 10.1063/1.4985302, 2017.

Murakami, G., Hayakawa, H., Ogawa, H., Matsuda, S., Seki, T., Kasaba, Y., Saito, Y., Yoshikawa, I., Kobayashi, M., Baumjohann, W., Matsuoka, A., Kojima, H., Yagitani, S., Moncuquet, M., Wahlund, J.-E., Delcourt, D.,
 415 Hirahara, M., Barabash, S., Korabely, O., and Fujimoto, M.: Mio—First Comprehensive Exploration of Mercury’s Space Environment: Mission Overview, *Space Sci Rev*, 216, 113, 10.1007/s11214-020-00733-3, 2020.
 Øieroset, M., Phan, T. D., Haggerty, C., Shay, M. A., Eastwood, J. P., Gershman, D. J., Drake, J. F., Fujimoto, M., Ergun, R. E., Mozer, F. S., Oka, M., Torbert, R. B., Burch, J. L., Wang, S., Chen, L. J., Swisdak, M., Pollock, C.,
 420 Dorelli, J. C., Fuselier, S. A., Lavraud, B., Giles, B. L., Moore, T. E., Saito, Y., Avannov, L. A., Paterson, W., Strangeway, R. J., Russell, C. T., Khotyaintsev, Y., Lindqvist, P. A., and Malakit, K.: MMS observations of large guide field symmetric reconnection between colliding reconnection jets at the center of a magnetic flux rope at the magnetopause, *Geophys Res Lett*, 43, 5536-5544, <https://doi.org/10.1002/2016GL069166>, 2016.
 Paschmann, G., Haerendel, G., Papamastorakis, I., Sckopke, N., Bame, S. J., Gosling, J. T., and Russell, C. T.: Plasma and magnetic field characteristics of magnetic flux transfer events, *J Geophys Res Space Phys*, 87, 2159-2168, <https://doi.org/10.1029/JA087iA04p02159>, 1982.
 425 Phan, T. D., Sonnerup, B. U. Ö., and Lin, R. P.: Fluid and kinetics signatures of reconnection at the dawn tail magnetopause: Wind observations, *J Geophys Res Space Phys*, 106, 25489-25501, <https://doi.org/10.1029/2001JA900054>, 2001.
 Phan, T. D., Eastwood, J. P., Shay, M. A., Drake, J. F., Sonnerup, B. U. Ö., Fujimoto, M., Cassak, P. A., Øieroset, M., Burch, J. L., Torbert, R. B., Rager, A. C., Dorelli, J. C., Gershman, D. J., Pollock, C., Pyakurel, P. S., Haggerty, C. C., Khotyaintsev, Y., Lavraud, B., Saito, Y., Oka, M., Ergun, R. E., Retino, A., Le Contel, O., Argall, M. R.,
 430 Giles, B. L., Moore, T. E., Wilder, F. D., Strangeway, R. J., Russell, C. T., Lindqvist, P. A., and Magnes, W.: Electron magnetic reconnection without ion coupling in Earth’s turbulent magnetosheath, *Nature*, 557, 202-206, 10.1038/s41586-018-0091-5, 2018.
 435 Pi, G., Shue, J.-H., Grygorov, K., Li, H.-M., Němeček, Z., Šafránková, J., Yang, Y.-H., and Wang, K.: Evolution of the magnetic field structure outside the magnetopause under radial IMF conditions, *J Geophys Res Space Phys*, 122, 4051-4063, <https://doi.org/10.1002/2015JA021809>, 2017.
 Poh, G., Slavin, J. A., Lu, S., Le, G., Ozturk, D. S., Sun, W.-J., Zou, S., Eastwood, J. P., Nakamura, R., Baumjohann, W., Russell, C. T., Gershman, D. J., Giles, B. L., Pollock, C. J., Moore, T. E., Torbert, R. B., and
 440 Burch, J. L.: Dissipation of Earthward Propagating Flux Rope Through Re-reconnection with Geomagnetic Field: An MMS Case Study, *J Geophys Res Space Phys*, 124, 7477-7493, <https://doi.org/10.1029/2018JA026451>, 2019.
 Pritchett, P. L. and Coroniti, F. V.: Three-dimensional collisionless magnetic reconnection in the presence of a guide field, *J Geophys Res Space Phys*, 109, 10.1029/2003ja009999, 2004.
 Raeder, J.: Flux Transfer Events: 1. generation mechanism for strong southward IMF, *Ann. Geophys.*, 24, 381-392, 10.5194/angeo-24-381-2006, 2006.
 445 Ricci, P., Brackbill, J. U., Daughton, W., and Lapenta, G.: Collisionless magnetic reconnection in the presence of a guide field, *Phys Plasmas*, 11, 4102-4114, 10.1063/1.1768552, 2004.
 Rijnbeek, R. P., Cowley, S. W. H., Southwood, D. J., and Russell, C. T.: A survey of dayside flux transfer events observed by ISEE 1 and 2 magnetometers, *J Geophys Res Space Phys*, 89, 786-800, 10.1029/JA089iA02p00786, 1984.
 450 Russell, C. T. and Elphic, R. C.: Initial ISEE magnetometer results: magnetopause observations, *Space Sci Rev*, 22, 681-715, 10.1007/BF00212619, 1978.
 Russell, C. T. and Walker, R. J.: Flux transfer events at Mercury, *J Geophys Res Space Phys*, 90, 11067-11074, 10.1029/JA090iA11p11067, 1985.
 455 Saito, Y., Delcourt, D., Hirahara, M., Barabash, S., André, N., Takashima, T., Asamura, K., Yokota, S., Wieser, M., Nishino, M. N., Oka, M., Futaana, Y., Harada, Y., Sauvaud, J.-A., Louarn, P., Lavraud, B., Génot, V., Mazelle, C., Dandouras, I., Jacquy, C., Aoustin, C., Barthe, A., Cadu, A., Fedorov, A., Frezoul, A.-M., Garat, C., Le Comte, E., Lee, Q.-M., Médale, J.-L., Moirin, D., Penou, E., Petiot, M., Peyre, G., Rouzaud, J., Séran, H.-C., Němeček, Z., Šafránková, J., Marcucci, M. F., Bruno, R., Consolini, G., Miyake, W., Shinohara, I., Hasegawa, H., Seki, K.,
 460 Coates, A. J., Leblanc, F., Verdeil, C., Katra, B., Fontaine, D., Illiano, J.-M., Berthelier, J.-J., Techer, J.-D., Fraenz, M., Fischer, H., Krupp, N., Woch, J., Bührke, U., Fiethe, B., Michalik, H., Matsumoto, H., Yanagimachi, T., Miyoshi, Y., Mitani, T., Shimoyama, M., Zong, Q., Wurz, P., Andersson, H., Karlsson, S., Holmström, M., Kazama, Y., Ip, W.-H., Hoshino, M., Fujimoto, M., Terada, N., Keika, K., and BepiColombo Mio, M. T.: Pre-flight Calibration and Near-Earth Commissioning Results of the Mercury Plasma Particle Experiment (MPPE) Onboard MMO (Mio), *Space Sci Rev*, 217, 70, 10.1007/s11214-021-00839-2, 2021.
 465 Saunders, M. A., Russell, C. T., and Sckopke, N.: Flux transfer events: Scale size and interior structure, *Geophys Res Lett*, 11, 131-134, 10.1029/GL011i002p00131, 1984.

Sauvaud, J. A., Fedorov, A., Aoustin, C., Seran, H. C., Le Comte, E., Petiot, M., Rouzaud, J., Saito, Y., Dandouras, J., Jacquy, C., Louarn, P., Mazelle, C., and Médale, J. L.: The Mercury Electron Analyzers for the Bepi Colombo mission, *Adv Space Res*, 46, 1139-1148, <https://doi.org/10.1016/j.asr.2010.05.022>, 2010.

Schindler, K., Pfirsch, D., and Wobig, H.: Stability of two-dimensional collision-free plasmas, *Plasma Phys*, 15, 1165-1184, 10.1088/0032-1028/15/12/001, 1973.

Shi, Q. Q., Zong, Q. G., Zhang, H., Pu, Z. Y., Fu, S. Y., Xie, L., Wang, Y. F., Chen, Y., Li, L., Xia, L. D., Liu, Z. X., Fazakerley, A. N., Reme, H., and Lucek, E.: Cluster observations of the entry layer equatorward of the cusp under northward interplanetary magnetic field, *J Geophys Res Space Phys*, 114, 10.1029/2009JA014475, 2009.

Shi, Q. Q., Zong, Q. G., Fu, S. Y., Dunlop, M. W., Pu, Z. Y., Parks, G. K., Wei, Y., Li, W. H., Zhang, H., Nowada, M., Wang, Y. B., Sun, W. J., Xiao, T., Reme, H., Carr, C., Fazakerley, A. N., and Lucek, E.: Solar wind entry into the high-latitude terrestrial magnetosphere during geomagnetically quiet times, *Nat Commun*, 4, 1466, 10.1038/ncomms2476, 2013.

Slavin, J. A., Imber, S. M., Boardsen, S. A., DiBraccio, G. A., Sundberg, T., Sarantos, M., Nieves-Chinchilla, T., Szabo, A., Anderson, B. J., Korth, H., Zurbuchen, T. H., Raines, J. M., Johnson, C. L., Winslow, R. M., Killen, R. M., McNutt Jr., R. L., and Solomon, S. C.: MESSENGER observations of a flux-transfer-event shower at Mercury, *J Geophys Res Space Phys*, 117, 10.1029/2012ja017926, 2012.

Slavin, J. A., Lepping, R. P., Wu, C.-C., Anderson, B. J., Baker, D. N., Benna, M., Boardsen, S. A., Killen, R. M., Korth, H., Krimigis, S. M., McClintock, W. E., McNutt Jr., R. L., Sarantos, M., Schriver, D., Solomon, S. C., Trávníček, P., and Zurbuchen, T. H.: MESSENGER observations of large flux transfer events at Mercury, *Geophys Res Lett*, 37, 10.1029/2009gl041485, 2010.

Slavin, J. A., Acuña, M. H., Anderson, B. J., Baker, D. N., Benna, M., Boardsen, S. A., Gloeckler, G., Gold, R. E., Ho, G. C., Korth, H., Krimigis, S. M., McNutt, R. L., Raines, J. M., Sarantos, M., Schriver, D., Solomon, S. C., Trávníček, P., and Zurbuchen, T. H.: MESSENGER Observations of Magnetic Reconnection in Mercury's Magnetosphere, *Science*, 324, 606, 10.1126/science.1172011, 2009.

Song, P. and Russell, C. T.: Model of the formation of the low-latitude boundary layer for strongly northward interplanetary magnetic field, *J Geophys Res Space Phys*, 97, 1411-1420, <https://doi.org/10.1029/91JA02377>, 1992.

Sonnerup, B. U. Ö.: Magnetopause reconnection rate, *J Geophys Res*, 79, 1546-1549, 10.1029/JA079i010p01546, 1974.

Sonnerup, B. U. Ö. and Cahill Jr., L. J.: Magnetopause structure and attitude from Explorer 12 observations, *J Geophys Res*, 72, 171-183, 10.1029/JZ072i001p00171, 1967.

Sonnerup, B. U. Ö. and Scheible, M.: Minimum and maximum variance analysis, in: *Analysis methods for multi-spacecraft data*, edited by: Paschmann, G., and Daly, P. W., ESA Publication, Noordwijk, Netherlands., 185-220, 1998.

Sonnerup, B. U. Ö., Paschmann, G., Papamastorakis, I., Sckopke, N., Haerendel, G., Bame, S. J., Asbridge, J. R., Gosling, J. T., and Russell, C. T.: Evidence for magnetic field reconnection at the Earth's magnetopause, *J Geophys Res Space Phys*, 86, 10049-10067, 10.1029/JA086iA12p10049, 1981.

Stawarz, J. E., Eastwood, J. P., Phan, T. D., Gingell, I. L., Shay, M. A., Burch, J. L., Ergun, R. E., Giles, B. L., Gershman, D. J., Contel, O. L., Lindqvist, P. A., Russell, C. T., Strangeway, R. J., Torbert, R. B., Argall, M. R., Fischer, D., Magnes, W., and Franci, L.: Properties of the Turbulence Associated with Electron-only Magnetic Reconnection in Earth's Magnetosheath, *Astrophys J*, 877, L37, 10.3847/2041-8213/ab21c8, 2019.

Sun, W., Dewey, R. M., Aizawa, S., Huang, J., Slavin, J. A., Fu, S., Wei, Y., and Bowers, C. F.: Review of Mercury's dynamic magnetosphere: Post-MESSENGER era and comparative magnetospheres, *Sci China Earth Sci*, 65, 25-74, 10.1007/s11430-021-9828-0, 2022.

Sun, W. J., Slavin, J. A., Smith, A. W., Dewey, R. M., Poh, G. K., Jia, X., Raines, J. M., Livi, S., Saito, Y., Gershman, D. J., DiBraccio, G. A., Imber, S. M., Guo, J. P., Fu, S. Y., Zong, Q. G., and Zhao, J. T.: Flux Transfer Event Showers at Mercury: Dependence on Plasma β and Magnetic Shear and Their Contribution to the Dungey Cycle, *Geophys Res Lett*, 47, e2020GL089784, <https://doi.org/10.1029/2020GL089784>, 2020a.

Sun, W. J., Slavin, J. A., Dewey, R. M., Chen, Y., DiBraccio, G. A., Raines, J. M., Jasinski, J. M., Jia, X., and Akhavan-Tafti, M.: MESSENGER Observations of Mercury's Nightside Magnetosphere Under Extreme Solar Wind Conditions: Reconnection-Generated Structures and Steady Convection, *J Geophys Res Space Phys*, 125, e2019JA027490, 10.1029/2019ja027490, 2020b.

Tang, B. B., Wang, C., and Li, W. Y.: The magnetosphere under the radial interplanetary magnetic field: A numerical study, *J Geophys Res Space Phys*, 118, 7674-7682, <https://doi.org/10.1002/2013JA019155>, 2013.

Toledo-Redondo, S., Hwang, K.-J., Escoubet, C. P., Lavraud, B., Fornieles, J., Aunai, N., Fear, R. C., Dargent, J., Fu, H. S., Fuselier, S. A., Genestreti, K. J., Khotyaintsev, Y. V., Li, W. Y., Norgren, C., and Phan, T. D.: Solar

Wind—Magnetosphere Coupling During Radial Interplanetary Magnetic Field Conditions: Simultaneous Multi-Point Observations, *J Geophys Res Space Phys*, 126, e2021JA029506, <https://doi.org/10.1029/2021JA029506>, 2021.

525 Trattner, K. J., Mulcock, J. S., Petrinec, S. M., and Fuselier, S. A.: Probing the boundary between antiparallel and component reconnection during southward interplanetary magnetic field conditions, *J Geophys Res Space Phys*, 112, <https://doi.org/10.1029/2007JA012270>, 2007.

Trattner, K. J., Petrinec, S. M., Fuselier, S. A., and Phan, T. D.: The location of reconnection at the magnetopause: Testing the maximum magnetic shear model with THEMIS observations, *J Geophys Res Space Phys*, 117, <https://doi.org/10.1029/2011JA016959>, 2012.

530 Trattner, K. J., Fuselier, S. A., Petrinec, S. M., Burch, J. L., Ergun, R., and Grimes, E. W.: Long and Active Magnetopause Reconnection X-Lines During Changing IMF Conditions, *J Geophys Res Space Phys*, 126, e2020JA028926, <https://doi.org/10.1029/2020JA028926>, 2021.

535 Trattner, K. J., Burch, J. L., Ergun, R., Eriksson, S., Fuselier, S. A., Giles, B. L., Gomez, R. G., Grimes, E. W., Lewis, W. S., Mauk, B., Petrinec, S. M., Russell, C. T., Strangeway, R. J., Trenchi, L., and Wilder, F. D.: The MMS Dayside Magnetic Reconnection Locations During Phase 1 and Their Relation to the Predictions of the Maximum Magnetic Shear Model, *J Geophys Res Space Phys*, 122, 11,991-912,005, <https://doi.org/10.1002/2017JA024488>, 2017.

540 Walker, R. J. and Russell, C. T.: Flux transfer events at the Jovian magnetopause, *J Geophys Res Space Phys*, 90, 7397-7404, [10.1029/JA090iA08p07397](https://doi.org/10.1029/JA090iA08p07397), 1985.

Wang, Y. L., Elphic, R. C., Lavraud, B., Taylor, M. G. G. T., Birn, J., Russell, C. T., Raeder, J., Kawano, H., and Zhang, X. X.: Dependence of flux transfer events on solar wind conditions from 3 years of Cluster observations, *J Geophys Res Space Phys*, 111, [doi:10.1029/2005JA011342](https://doi.org/10.1029/2005JA011342), 2006.

545 Wang, Y. L., Elphic, R. C., Lavraud, B., Taylor, M. G. G. T., Birn, J., Raeder, J., Russell, C. T., Kawano, H., Zong, Q.-G., Zhang, H., Zhang, X. X., and Friedel, R. H.: Initial results of high-latitude magnetopause and low-latitude flank flux transfer events from 3 years of Cluster observations, *J Geophys Res Space Phys*, 110, [doi:10.1029/2005JA011150](https://doi.org/10.1029/2005JA011150), 2005.

Zhang, H., Khurana, K. K., Kivelson, M. G., Angelopoulos, V., Pu, Z. Y., Zong, Q.-G., Liu, J., and Zhou, X.-Z.: Modeling a force-free flux transfer event probed by multiple Time History of Events and Macroscale Interactions during Substorms (THEMIS) spacecraft, *J Geophys Res Space Phys*, 113, [10.1029/2008ja013451](https://doi.org/10.1029/2008ja013451), 2008.

550 Zhao, J. T., Sun, W.-J., Zong, Q. G., Slavin, J. A., Zhou, X. Z., Dewey, R. M., Poh, G. K., and Raines, J. M.: A Statistical Study of the Force Balance and Structure in the Flux Ropes in Mercury's Magnetotail, *J Geophys Res Space Phys*, 124, 5143-5157, [10.1029/2018ja026329](https://doi.org/10.1029/2018ja026329), 2019.

555 Zhong, J., Wei, Y., Lee, L. C., He, J. S., Slavin, J. A., Pu, Z. Y., Zhang, H., Wang, X. G., and Wan, W. X.: Formation of Macroscale Flux Transfer Events at Mercury, *Astrophys J*, 893, L18, [10.3847/2041-8213/ab8566](https://doi.org/10.3847/2041-8213/ab8566), 2020.

Zhou, M., Berchem, J., Walker, R. J., El-Alaoui, M., Deng, X., Cazzola, E., Lapenta, G., Goldstein, M. L., Paterson, W. R., Pang, Y., Ergun, R. E., Lavraud, B., Liang, H., Russell, C. T., Strangeway, R. J., Zhao, C., Giles, B. L., Pollock, C. J., Lindqvist, P. A., Marklund, G., Wilder, F. D., Khotyaintsev, Y. V., Torbert, R. B., and Burch, J. L.: Coalescence of Macroscopic Flux Ropes at the Subsolar Magnetopause: Magnetospheric Multiscale Observations, *Phys Rev Lett*, 119, 055101, [10.1103/PhysRevLett.119.055101](https://doi.org/10.1103/PhysRevLett.119.055101), 2017.

560




Figure 1. The electrons and magnetic field measurements of the dayside magnetopause during BepiColombo's Earth flyby. (a) the time-energy spectrogram of normalized electron counts from Mio/MEA, (b) magnetic x component B_x , (c) y component B_y , (d) z component B_z , (e) the magnetic component normal to the magnetopause B_N , (f) the magnetic field intensity B_t , (g) the clock angle (θ), (h) B_x/B_t of the IMF, (i) solar wind number density (n_p), (j) solar wind Alfvénic Mach number (M_A). All quantities are in the Geocentric Solar Magnetospheric (GSM) coordinate. The green arrows in (e) indicate the six FTE-type flux ropes. "S" indicates southward traveling and "N" northward traveling. The θ in (f) is defined as $\arctan(B_y/B_z)$, ranging from 0° to 360° . In (g), the black line is from the measurements of MPO/MAG, and the blue line is from OMNI.

575 **Table 1. List and properties of FTE-type flux ropes observed during BepiColombo’s dayside magnetopause crossing**

#	Time	Duration (s)	Travelling Direction	Core Field Intensity (nT)	Scale (km) ^b	Flux Content (kWb)	χ^2
1	00:11:04	~ 12	Southward	— ^a	—	—	—
2	00:26:06	~ 7	Northward	~23.9	462, 388 (0.84)	~13.7	~0.04
3	00:26:26	~ 6	Northward	~60.8	565, 524 (0.93)	~22.5	~0.04
4	00:26:35	~ 4	Northward	—	—	—	—
5	00:28:13	~ 20	Northward	~41	1745, 1281 (0.73)	~300	~0.08
6	00:30:26	~ 15	Northward	~45.2	1853,1745 (0.94)	~188	~0.08

^a “—” indicate that the values are not determined by the flux rope model. See the text for more information on the flux rope modeling.

580 ^b Scale contains semi-minor, semi-major, and the ratio between semi-minor and semi-major refers to the flattened profile. See the text for more information.





Figure 2. The southward traveling FTE-type flux rope centered at ~ 00:11:04 UT. (a) magnetic field component in the L direction, B_L , (b) magnetic field component in the M direction, B_M , (c) magnetic field component in the N direction, B_N , (d) B_t . This LMN is the local coordinate of the magnetopause. (e) and (f) are the hodograms of the magnetic field measurements under the local coordinate of the flux rope. The “B” and “E” indicate the beginning and the end of the data points.

Figure 3. Magnetic shear angle plots on the magnetopause surface during BepiColombo's dayside magnetopause crossing, which are obtained through the maximum magnetic shear model [Trattner *et al.*, 2007]. (a), (b), (c) correspond to the IMF averaged from 00:05 to 00:13 UT, 00:16 to 00:24 UT and 00:24 to 00:33 UT, respectively. The black circle represents the terminator plane separating the dayside magnetopause from the tailward magnetopause. The grey line represents the predicted magnetopause reconnection line. White areas correspond to the magnetic shear angle is within 3° of 180° . The black dots are the location of BepiColombo ("BC"). In (c), the predicted X-line is not provided. This is because a continuous X-line along the maximum magnetic shear location is difficult to obtain under the situation of a $B_x/B_t \geq 0.9$,






Figure 4. Overview of the flux rope centered at ~ 00:28:13 UT with the coalescence feature. (a) B_L , (b) B_M , (c) B_N , (d) B_t . The dashed lines are obtained from the flux rope model. This LMN is the local coordinate of the magnetopause. See the text for more information. (e) An illustration of the coalescence event and the BepiColombo's trajectory. The secondary reconnection site is marked by the magenta region. (f) and (g) are the hodograms of the magnetic field measurements under the local coordinate of the flux rope. The "B" and "E" indicate the beginning and the end of the data points. FR#A and FR#B indicate two flux ropes.




Figure 5. The magnetic field measurements under their separately local coordinate for the reconnecting current sheet of the coalescence event and the magnetopause current sheet. (a) to (d) are for the secondary current sheet of the coalescence event. The \vec{n}_{\min} , \vec{n}_{int} and \vec{n}_{\max} refer to the local coordinate of the secondary current sheet. (e) to (h) are for the magnetopause current sheet. The LMN local coordinate of the magnetopause current sheet is used. The eigenvalues and corresponding eigenvectors result from the MVA.

625