

Temporal and altitudinal variability of the Spread F observed by the VHF radar over Christmas Island

5 Ricardo Yvan de La Cruz Cueva¹, Eurico Rodrigues de Paula², Acácio Cunha Neto²

¹Physics Department, State University of Maranhão, São Luís, Maranhão, Brazil.

²DIHPA- Heliophysics, Planetary Sciences and Aeronomy Division, National Institute for Space Research, São José dos Campos, São Paulo, Brazil.

10 *Correspondence to:* Ricardo Y. C. Cueva (navivacu@gmail.com)

Abstract. The goal of this work is to study the time and altitude echoes characteristics under different solar and seasonality conditions using the VHF radar RTI images. The occurrence of equatorial spread F depends on the existence of conditions that can seed the Raileight-Taylor instability, and these conditions can change with solar flux, seasonality, longitude distributions, and day-to-day variability. So, the equatorial spread F is observed as its time and altitude occurrence. The VHF 15 radar of Christmas Island (2.0° N, 157.4° W, 2.9°N dip latitude) has been operational in the equatorial region for some time, allowing long-term observations. The occurrence of echoes during solar minimum conditions are observed all throughout the night, since the post reversal westward electric field is weaker than the solar maximum and the possibilities for the vertical plasma drift to become positive are larger. On other hand, echoes during solar maximum will be controlled by dynamics near the time of the Pre-reversal Peak (PRE). Our results indicate peak time occurrence of echoes along this period shows a well- 20 defined pattern, with echoes being distributed as closer to local sunset during solar maximum and around/closer midnight during solar minimum conditions, meanwhile, the peak altitude occurrence of echoes shows a slightly regular pattern with higher altitude occurrences during solar maxima and lower altitudes during solar minimum conditions.

25

30

1 Introduction

35 The contemporaneous understanding of the formation of F-region plasma irregularities depends mainly on the Rayleigh-Taylor (RT) instability process, due to its appearance at the bottomside of the F-region, then becoming unstable to finally generate plasma bubbles. These recently formed plasma bubbles evolve in nonlinear process then extend into high altitudes into the F-region. The small-scale (centimeter to a few tens of meters) irregularities formed in this process are the responsibles for radar backscatter, which can be observed as structures in the range-time-intensity (RTI) image of the radar.

40 The pioneering ionospheric radar work of Woodman and LaHoz (1976) attributed the term “plumes” to describe radar echoes reaching the topside ionosphere. They observed *a* slope in the formation of the plumes, then explained using numerical simulation by Ossakow (1981) and Zalesak et al. (1982).

45 The RT instability (and ESF) is controlled by a number of parameters including the prereversal enhancement (PRE) of the zonal equatorial electric field, zonal and meridional neutral winds, longitudinal conductivity gradients, flux tube integrated conductivities, and, possibly, variations in initial (or seed) perturbations (Abdu, 2001; Fejer et al., 1999). It has been noted that ESF bubbles at pre-midnight and post-midnight hours could be driven by different mechanisms (Dao et al., 2011; Yizengaw et al., 2013). The mechanisms that should control the appearance or suppression of equatorial plasma irregularities are different for the pre- and post midnight periods, due to the ambient conditions that prevail along night.

50 Yizengaw et al. (2009) showed that h'F presents a peak at post-midnight hours that indicate the existence of some electrodynamic force that drives the F layer upward creating conditions for irregularities development.

55 The effects of solar and geomagnetic activities on spread-F vary with latitude and longitude. Cueva et al. (2013) examined data from three equatorial stations along solar minimum and maximum conditions. Their results showed that there was an increase in the spread-F occurrence rate with solar flux. Although many researchers have discussed the characteristics of spread-F irregularities at equatorial and low latitudes, some issues are needed of better understanding in their spatial and temporal variability of spread-F and plasma bubbles. So, the analysis of long term data were performed in this work covering high and low solar conditions with spread F echoes observations over the Central Pacific region using the VHF radar installed in Christmas Island. In this study we present results from data analysis of echoes distribution using the 50-MHz

60 Christmas Island radar along 2003 and 2012 time period. The observations allowed us to determine how the echoes vary with local time and height throughout different seasons and solar flux conditions.

65 **2 Measurements and Analysis**

2.1 VHF radar measurements

The Christmas Island VHF radar provides data of meter-scale F-region irregularities routinely, being initially operated by Stanfor Research Institute - SRI International (2002-2007) and then operated by the US Air Force Research Laboratory 70 (AFRL). The system uses a 100 m x 100 m coaxial collinear (COCO) antenna array. Two stationary beams are used for measurements. One beam is pointed North (azimuth 0° and elevation 84.5°) and the other one is pointed to the east (azimuth 90° and elevation 60.5°). The coherent radar detects fluctuations related to the plasma instabilities called field-aligned irregularities, then detection of such irregularities requires the antenna to be pointing perpendicular to the geomagnetic field line (Tsunoda et al., 2000; Tsunoda et al., 1979). Then, the north beam antenna was choosen due to be pointed in the north 75 direction to reach perpendicularly to the magnetic field line. More technical details of this radar can be found at Miller et al. (2009). Its geographic position is 2.0° N, 157.4° W, 2.9°N dip latitude, and its magnetic inclination (declination) varied from 4.69° (9.36°E) in 2003 to 4.61° (9.38°E) in 2012.

It's worth mentioning that measurements available to this study cover different solar conditions when F10.7 varied from 200 80 SFU (high solar flux conditions) to 66 SFU (low solar flux conditions), as shown in Figure 1. Data measurements of spread F echoes to this study are between the period of January 2003 and December 2012. All our data are presented as altitude integration from 200 km to 1000 km height as function of signal to noise ratio (presented in Figure 2), and the horizontal dashed lines (at 20 LT and 00 LT) representing time threshold to assist observation of time-echoes distribution. Lack of data are also presented as black space in the figure, mainly for 2014 (March equinox and June solstice).

85

Our interest focus in the local occurrence of F-region echoes (5-meter-scale irregularities) as one of the most interesting and challenging phenomenon for space weather and climatological models. The physical mechanism responsible for this phenomenon are complexes and not fully understood. So, we organized our data attempting to present the difference in seasonal and solar flux conditions as a function of time and height of irregularities observed in the VHF-Radar. For this 90 study, we limit our focus to quiet-time irregularities.

95 **2.2 Data Analysis**

It is well known that high geomagnetic activities directly cause drastic perturbations in the zonal electric field, in the equatorial and low latitude regions, affecting the growth and development of ionospheric irregularities. These perturbations can be categorized as prompt penetration (PP) and disturbance dynamo (DD) electric field (Abdu et al., 2018; Astafyeva et 100 al., 2018; and Shreedevi & Choudhary, 2017). These perturbed electric fields occurring in the post sunset period can enhance/weaken the regular eastward vertical plasma drift, then affecting the uplift of the F layer (Fejer et al., 1991), and as a consequence affecting the generation of irregularities (Aarons. 1991; Abdu, 2012).

In sequence, to avoid the disturbed geomagnetic periods and their effects on irregularity generations, we classify the data 105 with low geomagnetic conditions using the 3-hour Planetary K index (known as Kp). Each measurement was tagged with the value of Kp for the time, of the measurement, plus the previous 3 Kp values. We limited our study to quiet geomagnetic conditions, to be those when none of the three Kp indexes exceeded 3.

The solstice is when the Sun reaches the most southerly or northerly point in the sky, while an equinox is when the Sun 110 passes over Earth's equator. For example, June solstice, or June 21, is the longest day of the year in the northern hemisphere. So, to sort our measurements according the four seasons Spring, Summer, Fall and Winter we use 91 days of data centered on each day 21 of March, June, September and December, respectively. We used the quiet time radar echoes, for each season, to obtain the occurrence rate of echoes. We establish that a good representation of irregularity occurrence is given by echoes distribution above 0 dB divided by the total number of observations. Our criterion is a good commitment among 115 being able to identify the occurrence of spread F echoes and to eliminate the effects of non-geophysical echoes.

The sample rate of the VHF radar is estimated for every 15-minute intervals starting at 18:00 LT, right before sunset until 05:00 LT near sunrise. To construct maps of irregularity occurrence rate in function of height and local time we had computed for every 15-km height intervals starting at 200 km up to 1000 km altitude.

120

3 Results and Discussions

During solar maximum the spread F events occur near the time when upward drift is large which is promptly after local sunset (Fejer et al., 1999), while during solar minimum the upward drift is usually short, the spread F exists throughout the 125 whole night, so upward and downward ionospheric conditions may play a role in the morphology of irregularities. Stoneback et al. (2011) showed the role of vertical drift during the extended solar minimum and how it varies from sunset until

postmidnight period. These previous work observations increase the need of further study of climatology of echoes evolution in time and altitude.

130 In the way of understanding this climatology of spread F evolution along seasonality and solar activity we analyzed radar echoes occurrence as function of time and altitude along solar maximum and extended solar minimum periods, since the evening vertical drifts and layers heights increase noticeably with solar activity, and along nighttime.

3.1 On the height variability of echo occurrence rates

135 Peak altitude profiles of the occurrence rate of F-region echoes are shown in Figure 3, top panels. They were organized by seasons (March equinox, June Solstice, September equinox and December solstice, from left to right respectively) along the period of 2003 to 2012. Horizontal dashed lines were placed at 250 and 350 km height to assist observation (hereafter called altitude threshold).

140 Comparing the peak of altitude echoes along seasons, we can observe higher occurrence rates of all years over June solstice and September equinox than March equinox and December solstice seasons. This observation match with previous observation by Cueva et al. (2013), that shown the peak occurrence of equatorial spread F for this region being around July-August months.

145 When analyzing solar minimum years 2006 and 2008) we can lay down our attention to the peak echoes altitude, which was slightly higher, in altitude, in June solstice than in September equinox. For the occurrence rates of peak time we observed being bigger in September equinox, and peak altitude occurrence before midnight (as in bottom panel), than in June solstice with peak altitude occurrence around midnight hours (as in bottom panel). For years of solar maximum (2003 and 2012) we
150 can mention that peak altitude distribution is the highest, mainly during June and september seasons, nevertheless present minor percentage of occurrence than solar minimum years (2006 to 2008). The minimum occurrence of peak altitude occurs in March equinox, which is the period of scarce spread F echoes over Christmas Island region. During solar maximum period spread F echoes have less occurrence than in solar minimum period, reaching higher altitudes as observed in June solstice 2003 when the peak altitude was higher than the threshold altitude of 350 km. During September equinox higher
155 plumes are frequently observed than in other periods which agrees with results presented by Cueva et al.(2013).

3.2 On the time variability of echo occurrence rates

Time variation in the occurrence rates of F-region echoes for the period in study is shown in Figure 3, lower panels, also
160 separated by seasons (March equinox, June Solstice, September equinox and December solstice, from left to right

respectively). The vertical dashed lines represent local sunset and local midnight. As we can observe the percentage of occurrence of echoes presents solar flux dependence. During solar maximum radar echoes are confined to a few hours after sunset, on the other hand during solar minimum echoes are more broaden out in time and can arise late in the evening after sunset and more closely to midnight hours. As we get closer to solar minimum period the amplitude of echoes occurrence increases due to high probability to occur echoes along all night. This can be observed during years 2006 to 2008 with more amplitude than echoes observed during solar maximum, similar finding was mentioned by Niranjan et al. (2003) when analyzed spread F data from 1997-2000 period, and also by Burke et al. (2004) and Dao et al. (2011) using satellite data from different geographical locations.

165 170 Seasonal dependence of echoes along solar cycle is also observed. September equinox has more conditions to develop irregularities over the region as explained before, as well higher echoes occurrence either for solar minimum and maximum periods. For March equinox and December solstice we have less probability of echoes occurrence as can be seen in the Figure 3, moreover amplitude of echoes occurrence is always lower in solar maximum than in solar minimum.

175 **4 Conclusions**

The variability over seasonality observed in the amplitude of peak echoes occurrence, either for altitude or time, is suitable for the seasonal spread F occurrence over the Pacific region. During high solar activity spread F were observed more often after sunset and rare/uncommon observations around mignight hours. The RT instability occurs at the magnetic equator after

180 185 sunset when the eastward electric fields increase and structures reaching to higher altitudes are due to vertical ExB drift at the equator, is well acknowledged for high solar flux periods. However, during the low solar cycle period observed (years 2006 to 2009) spread F didn't reached higher altitudes, its appearance was very frequent around midnight hours, and last for many hours. The mechanism that govern its appearance is not longer the prereversal enhancement because it just happen around the sunset terminator. The generation mechanism for the post-midnight irregularities at quiet time during solar minimum conditions is still not clear, or not completely understood. Some authors also found similar occurrence, in solar minimum period, of plasma density irregularities mostly after midnight (Heelis et al., 2010, Li et al., 2011 and Dao et al., 2011). So, occurrence of post-midnight events were observed to present negative correlation with solar activity, decreasing from solar minimum to solar maximum.

190 Under quiet magnetically conditions, and solar minimum there are some possible seeding mechanisms competing that increase the probability for spread F generation along all night (pre-midnight and post-midnight), as well as uplifting the F layer. For example, gravity waves, launched from active convection region in the troposphere, could propagate into the ionosphere (Takahashi et al., 2009, 2010; Maurya et al., 2020; Correia et al., 2020) and contribute to the instability seeding. Another is the Medium-scale traveling ionospheric disturbances (MSTID) activity providing perturbations in the electric

195 fields for the low latitude F region to be unstable at postmidnight hours, that can seed the RT instability at the magnetic equator (Otsuka et al., 2009, Yokoyama et al., 2011 and Narayanan et al., 2019). Another mechanism could be the uplift of the F layer around midnight (Nicolls et al., 2006) caused by decreasing westward electric field in conjunction with sufficient recombination and plasma flux. However, the causes of midnight F-layer increase are not yet clearly established.

200 For the extended solar minimum period, during June solstice and December solstice months, we observed post-midnight echoes similar as previously reported by Otsuka et al. (2012) during 2005 to 2009 period. September equinox also presents post-midnight events for solar minimum period. Our findings are summarized in Figure 4. On top panel is presented UT (LT=UT+14) in the vertical axis for the time peak echoes occurrence along solar cycle separated by seasonality. We can clearly observe the peak time echoes occurrences being closer to the time of PRE during high solar activity years (see 2003, 205 2004 and 2011 and 2012) and around midnight during solar minimum conditions (see years 2007 to 2009). December solstice season during high solar conditions is not following this trend, and further study must be necessary in this point. The representation of the ionospheric sunset and the local midnight are plotted as horizontal dashed lines, and the error bars are the standard deviations.

210 According to our observations, during solar minimum the error bars (standard deviation) must be higher than during solar maximum periods due to the probability of spread F echoes occurrences, which are spread out in time (with maximum observations around local midnight) under minimum conditions and localized around to local sunset under maximum solar conditions.

215 Bottom panel on Figure 4 shows altitude peak variation along solar cycle, also separated by seasonality. Altitude parameter seems to follow a very good trend, being higher altitudes for solar maximum conditions and lower altitudes for solar minimum conditions. Again December solstice doesn't match very well with this trend. The error bars show the distribution of peak altitude echoes observations. We observe the size of error bars decreasing from solar maximum to solar minimum conditions. The altitude parameter is an important parameter since it is one key process in the generation mechanism for 220 ionospheric irregularities. Peak altitude echoes of June solstice reaches higher altitude difference from solar maximum to solar minimum periods, when compared with March and September equinoxes which were closer to 300 km most of the solar cycle period.

225 So, for Christmas Island sector we can conclude that spread F echoes occurs along all solar flux conditions. The PRE being the main mechanism for spread F generation, consequently occurrences arising closer to sunset terminator, with higher structures and short duration for solar maximum conditions. Spread F occurrence over December solstice season needs more study since it doesn't follow the peak time occurrence for solar maximum condition. For solar minimum conditions the mechanisms necessary for spread F generations are not clear, being the seeding of the RT instability and the uplift of the F

layer. Anyway, the spread F occurrences are happening along all night with high occurrence mainly around local midnight,

230 with peak altitude echoes distribution remaining around 300kms, and with long time duration.

5 Data availability

All raw data belong to AFRL Geospace Environment Applications and Impacts Program at Kirtland AFB. Data requirements

235 will be made directly to AFRL directorate.

6 Author contributions

Ricardo Y.C. Cueva came up with the idea, prepared all data analysis, then prepared the article draft and final version. E.R.

240 de Paula participated advising and revieweing the manuscript. Acácio C. Neto gave support with data analysis and equipments.

245 7 Competing interest

The authors declare that they have no conflict of interest.

8 Acknowledgement

250 The authors are very grateful to R.T. Tsunoda and K.M. Groves for providing the VHF radar data from Christmas Island equatorial station, also scknowledgements for the AFRL Geospace Enviroment Applications and Impacts Program at Kirtland AFB. The author E. R. de Paula thanks the support from CNPq 302531/2019-0 as well as the INCT GNSS-NavAer grants 2014/465648/2014-2 CNPq and 2017/50115-0 FAPESP. The author R. Y. C. Cueva thanks the PIBIC/UEMA program for constant support.

255

260

9 References

265

Aarons, J. (1991). The role of the ring current in the generation or inhibition of equatorial F layer irregularities during magnetic storms. *Radio Science*, 26(04), 1131-1149.

270 Abdu, M.A: Outstanding problems in the equatorial ionosphere–thermosphere electrodynamics relevant to spread-F, *Journal of Atmospheric and Solar Terrestrial Physics* 63, 869–884, 2001.

Abdu, M. A. (2012). Equatorial spread F/plasma bubble irregularities under storm time disturbance electric fields. *Journal of Atmospheric and Terrestrial Physics*, 75–76, 44–56.

275 Abdu, M. A., Nogueira, P. A. B., Santos, A. M., de Souza, J. R., Batista, I. S., & Sobral, J. H. A. (2018). Impact of disturbance electric fields in the evening on prereversal vertical drift and spread F developments in the equatorial ionosphere. *Annales Geophysicae*, 36, 609–620. <https://doi.org/10.5194/angeo-36-609-2018>.

280 Astafyeva, E., Zakharenkova, I., Hozumi, K., Alken, P., Coisson, P., Hairston, M. R., & Coley, W. R. (2018). Study of the Equatorial and Low-Latitude Electrodynamical and Ionospheric Disturbances During the 22–23 June 2015 Geomagnetic Storm Using Ground-Based and Spaceborne Techniques. *Journal of Geophysical Research: Space Physics*, 123(3), 2424–2440. <https://doi.org/10.1002/2017JA024981>.

285 Burke W.J., L.C. Gentile, C.Y. Huang, C.E. Valladares and S.Y. Su: Longitudinal variability of equatorial plasma bubbles observed by DMSP and ROCSAT-1, *Journal of Geophysical Research*, 109, A12301, doi:10.1029/2004JA010583, 2004.

Correia, E., Raunheitte, L. T. M., Bageston, J. V., and D'Amico, D. E.: Characterization of gravity waves in the lower ionosphere using very low frequency observations at Comandante Ferraz Brazilian Antarctic Station, *Ann. Geophys.*, 38, 385–394, <https://doi.org/10.5194/angeo-38-385-2020>, 2020.

290

Cueva R.Y.C., de Paula E.R., and Kherani A.E.: Statistical analysis of radar observed F region irregularities from three longitudinal sectors, *Ann. Geophys.*, 31, 2137-2146, 2013.

295 Dao E., M.C. Kelley, P. Roddy, J. Retterer, J.O. Ballenthin, O. de La Beaujardiere, and Y.-J. Su: Longitudinal and Seasonal dependence of nighttime equatorial plasma density irregularities during solar minimum detected on the C/NOFS satellite, *Geophysical Research Letter*, 38, L10104, doi: 10.1029/2011GL047046, 2011.

Fejer, B. G., de Paula, E. R., González, S. A., & Woodman, R. F. (1991). Average vertical and zonal F region plasma drifts over Jicamarca. *Journal of Geophysical Research: Space Physics*, 96(A8), 13901–13906. doi:10.1029/91ja01171.

300

Fejer B.G., Scherliess L, and de Paula E.R.: Effect of the vertical plasma drift velocity on the generation and evolution of equatorial spread F, *J. Geophys. Res.*, 104, 19859-19869, 1999.

Heelis R A, Stoneback R, Earle G D, Haaser R A & Abdu M A, Medium-scale equatorial plasma irregularities observed by 305 coupled ion-neutral dynamics investigation sensors aboard the Communication Navigation Outage Forecast System in a prolonged solar minimum, *J Geophys Res (USA)*, 115 (2010) A10321, doi: 10.1029/2010JA015596.

Li G, Ning B, Abdu M A, Yue X, Liu L, Wan W & Hu L, On the occurrence of postmidnight equatorial F region 310 irregularities during the June solstice, *J Geophys Res (USA)*, 116 (2011) A04318, doi: 10.1029/2010JA016056.

310

Maurya, A.K., Shrivastava, M.N. & Kumar, K.N. Ionospheric monitoring with the Chilean GPS eyeball during the South American total solar eclipse on 2nd July 2019. *Sci Rep* 10, 19380 (2020). <https://doi.org/10.1038/s41598-020-75986-7>.

Miller E.S., J.J. Makela, and M.C. Kelley: Seeding of equatorial plasma depletions by polarization electric fields from 315 middle latitudes: Experimental evidence, *Geophys. Res. Lett.*, 36, L18105, 2009.

Narayanan, V. L., Patra, A. K., Gurubaran, S., Pavan Chaitanya, P., & Emperumal, K. (2019). Coincident airglow, VHF radar, and ionosonde observations of electrified medium-scale traveling ionospheric disturbances in the equatorial latitudes. *Geophysical Research Letters*, 46. <https://doi.org/10.1029/2019GL083266>.

320

Nicolls MJ, Kelley MC, Vlasov MN, Sahai Y, Chau JL, Hysell DL, Fagundes PR, Becker-Guedes F, Lima WLC (2006) Observations and modeling of postmidnight uplifts near the magnetic equator. *Ann Geophys* 24:1317–1331.

Niranjan K., P. S. Brahmanandam, P. Ramakrishna Rao, G. Uma, D. S. V. V. D. Prasad, and P. V. S. Rama Rao: Post 325 midnight spread-F occurrence over Waltair (17.7° N, 83.3° E) during low and ascending phases of solar activity, *Annales Geophysicae*, 21: 745–750, 2003.

Ossakow, S. L.: Spread-F theories, *J. Atmos. Terr. Phys.*, 43, 437–452, 1981.

330 Otsuka Y, Ogawa T & Effendy, VHF radar observations of nighttime F-region field-aligned irregularities over Kototabang, Indonesia, *Earth, Planet Space (Japan)*, 61 (2009) pp 431-437.

Otsuka, Y., K. Shiokawa, M. Nishioka, and Effendy: VHF radar observations of post-midnight F-region field-aligned irregularities over Indonesia during solar minimum, *Indian J. Radio Space Phys.* Vol 41, pp 199-207, 2012.

335

Shreedevi, P. R., & Choudhary, R. K. (2017). Impact of oscillating IMF B_z during 17 March 2013 storm on the distribution of plasma over Indian low-latitude and mid-latitude ionospheric regions. *Journal of Geophysical Research: Space Physics*, 122, 11,607–11,623. <https://doi.org/10.1002/2017JA023980>.

340 Stoneback R.A., R.A. Heelis, A.G. Burrell, W.R. Coley, B.G. Fejer and E. Pacheco: Observations of quiet time vertical ionos drift in the equatorial ionosphere during the solar minimum period of 2009, *Journal of Geophysical Research*, 116, A12327, doi:10.1029/2011JA016712, 2011.

Takahashi, H., Taylor, M. J., Pautet, P.-D., Medeiros, A. F., Gobbi, D., Wrasse, C. M., Fechine, J., Abdu, M. A., Batista, I. 345 S., Paula, E., Sobral, J. H. A., Arruda, D., Vadas, S. L., Sabbas, F. S., and Fritts, D. C.: Simultaneous observation of ionospheric plasma bubbles and mesospheric gravity waves during the SpreadFEx Campaign, *Ann. Geophys.*, 27, 1477–1487, <https://doi.org/10.5194/angeo-27-1477-2009>, 2009.

Takahashi, H., et al. (2010), Equatorial ionosphere bottom-type spread F observed by OI 630.0 nm airglow imaging, 350 *Geophys. Res. Lett.*, 37, L03102, doi:10.1029/2009GL041802.

Tsunoda RT, Baron MJ, Owen J, Towle DM. ALTAIR: an incoherent scatter radar for equatorial spread F studies. *Radio Sci* 14:1111, 1979.

355 Tsunoda, Roland T., Ecklund, Warner L., Johnston, Paul E. Radar measurements of electric fields in the topside of the equatorial electrojet: First results. *Geophysical Research Letters*, 27 (18). 2861-2864, doi:10.1029/2000gl003775, 2000.

Woodman, R. F. and La Hoz, C.: Radar observations of F region equatorial irregularities. *J. Geophys. Res.*, 81, 5447–5466, 1976.

360

Yizengaw E., J. Retterer, E.E. Pacheco, P. Roddy, K. Groves, R. Caton, and P. Baki: Postmidnight bubbles and scintillation in the quiet-time June solstice, *Geophysical Research Letter*, 40, 1-6, doi: 10.1002/2013GL058307, 2013.

Yizengaw E., Maldwin, M.B.; Sahai Y.; and de Jesus R. Strong postmidnight equatorial ionospheric observations during
365 magnetically quiet period, Journal of Geophysical Research, vol. 114, A12308, 2009.

Yokoyama T, Yamamoto M, Otsuka Y, Nishioka M, Tsugawa T, Watanabe S & Pfaff R F, On post-midnight low-latitude
ionospheric irregularities during solar minimum, 1: Equatorial Atmosphere Radar and GPS-TEC observations in Indonesia, J
Geophys Res (USA), 116 (2011) A11325, doi: 10.1029/2011JA016797.

370

Zalesak, S. T., Ossakow, S. L., and Chaturvedi, P. K.: Nonlinear equatorial spread F – the effect of neutral winds and
background Pedersen conductivity. J. Geophys. Res., 87, 151–166, 1982.


375

380


385

390


395




405 **Figure 1: Solar flux index F10.7cm covering period used in this study, which covers solar conditions where F10.7 varied from 200 SFU to 66 SFU.**



425 **Figure 2: VHF radar data is presented as altitude integration from 200 km to 1000 km height as function of signal to noise ratio (dBKm), the horizontal lines represent local sunset and local midnight to help observation of echoes distribution.**



440 **Figure 3: Peak altitude (top panels) and time (bottom panels) variations along the years 2003, 2006, 2008 and 2012 years. Also
divided by seasons.**



450 **Figure 4: Peak time (top panel) and altitude (bottom panel) echoes occurrences along solar cycle, divided by seasons. The error bars show the standard deviation of observations.**