

Sounding of sporadic E layers from CSES radio occultation and comparing with ionosonde measurements

Chengkun Gan¹, Jiayu Hu², Xiaomin Luo³, Chao Xiong^{4,5}, Shengfeng Gu¹

¹GNSS Research Center, Wuhan University, Wuhan 430079, China

5 ²School of Geodesy and Geomatics, Wuhan University, Wuhan 430079, China

³School of Geography and Information Engineering, China University of Geosciences (Wuhan), Wuhan 430074, China

⁴Department of Space Physics, School of Electronic Information, Wuhan University, Wuhan 430072, China

⁵Hubei LuoJia Laboratory, Wuhan 430079, China

Correspondence to: Shengfeng Gu (gsf@whu.edu.cn)

10 **Abstract.** GNSS radio occultation (RO) plays an important role in ionospheric electron density inversion and sounding of sporadic E layers. As the China's first electromagnetic satellite, China Seismo Electromagnetic Satellite (CSES) has collected the RO data from both GPS and BDS-2 satellites since March 2018. In this study, we extracted the **carrier to noise density ratio (CNR)** **signal to noise ratio (SNR)** data of CSES and calculated the standard deviation of normalized **CNR-SNR**. A new criterion is developed to determine the Es events, that is when the mean value of the absolute value of the difference
15 between the normalized **CNR SNR** is greater than 3 times of the standard deviation. The statistics show that sporadic E layers have strong seasonal variations with highest occurrence rates in summer season at middle latitudes. It is also found that the occurrence height of Es is mainly located at 90-110 km, and the period of local time 15:00-18:00 is the high incidence period of Es. In addition, the geometric altitudes of a sporadic E layer detected in CSES radio occultation profiles and the virtual heights of a sporadic E layer obtained by the **Wuhan Zuoling Tai station (ZLT)** ionosonde show **four three**
20 different space-time matching criterions. Our results reveal that there is a good agreement between both parameters which is reflected in the significant correlation.

1. Introduction

The name Sporadic E and its abbreviation Es refer to thin layers of metallic ion plasma which accumulates in the dynamo region of the Earth's ionosphere, mostly between 100 and 125 km, where ion motion is controlled mainly by collisions with
25 the neutrals, thus the ions move with the winds while electrons remain strongly magnetized (Haldoupis 2012). The formation of sporadic E layer was traditionally attributed to the "windshear theory" (Whitehead 1961; Axford 1963; Whitehead 1989), in which vertical shears in the horizontal wind play a key role in forming these layers from long-lived metallic ions through ion-neutral collisional coupling and geomagnetic Lorentz forcing, vertical shear converge metallic ions into thin sheets of enhanced electron density. More recently, researchers have found multiple factors can contribute to the occurrence of Es,
30 including tidal wind, the Earth's geomagnetic field, and meteoric deposition of metallic material in the background

thermosphere, resulting in variations of Es occurrence with respect to local time, altitude, latitude, longitude, and season (Haldoupis, 2011; Yeh et al, 2014, Didebulidze et al., 2020). Meanwhile, the ionospheric E region has a relatively higher electrical conductivity and therefore plays a crucial role in the ionosphere electron dynamics at both E- and F-region altitudes (Yue et al. 2015).

35

Variance in the signal to noise ratio (SNR) caused by strong gradients in the index of refraction has been suggested to identify and sound sporadic E layers (Wu et al., 2005; Arras et al., 2008; Yeh et al., 2012; Hocke et al., 2001; Yue et al., 2015; Tsai et al. 2018). However, in terms of judgment criteria, many scholars propose different selection methods. Chu (2014) set thresholds for signal phase amplitude and carrier phase delay ratio when screening Es, and the ratio of disturbance amplitude to normalized SNR must be greater than 0.01 then it can be counted as Es event. Wu et al. (2005) directly used the normalized SNR data sequence as the characteristic parameter to detect Es. Arras et al. (2017) and Tsai et al. (2018) used the value of 0.2 as the threshold of the normalized SNR standard deviation sequence. It is considered that Es event occurs when the peak exceeds 0.2. Xue et al. (2018) used 0.1 as the standard deviation threshold to detect single-layer and multi-layer Es events at the same time. Based on GPS radio occultation (RO) techniques, some investigations established global distribution of Es layers information to analyze the climatology of global Es occurrence rates. (Arras et al., 2008; Wickert et al., 2004; Yeh et al., 2012; Tilo et al., 2014; Arras et al. 2017).

40

45

Since the invention of ionosonde in the nineteen thirties, Es has been investigated extensively from ground, by means of analyzing ionosonde and incoherent scatter radar observations (Whitehead 1989; Mathews 1998). Ionosondes provide reliable measurements on sporadic E parameters and on the altitude of each layer. The altitudes are given in virtual heights, and the lower boundary of the sporadic E layer ($h'Es$). Arras et al. (2017) compared sporadic E altitudes and their intensity with ground based ionosonde data provided by the Digisonde located at Pruhonic close to Prague, Czech Republic (geographic 50°N, 14.5°E) to confirm the derived sporadic E parameters. Wuhan Zuoling Tai station (ZLT) ionosonde (geographic 30.5°N, 114.4°E) is located in central China. It is a representative location due to its low geomagnetic latitude and the longest observational record, which has been well-maintained during the past several decades and its data are of high quality (Zhou et al. 2021).

50

55

China's first electromagnetic satellite, China Seismo Electromagnetic Satellite (CSES), also known as ZH01(01), was successfully launched on Feb 2, 2018 (Shen et al. 2018). The CSES is a 3-axes-stabilized satellite, based on the Chinese CAST2000 platform, with a mass of about 730 kg and peak power consumption of about 900 W. Scientific data are transmitted in the X-Band at 120 Mbps. The orbit is circular Sun-synchronous, at an altitude of about 507 km, inclination of about 97.4°, descending node at 14:00 LT. All payloads of CSES are designed to work in the region within the latitude of $\pm 65^\circ$ (Shen et al. 2018). In recent years, a few studies were published concerning the performance of different payloads of CSES. Ambrosi et al. (2018) investigated the seismo-associated perturbations of the Van Allen belts using the High

60

65 Energetic Particle Detector (HEPD) of the CSES satellite mission. Concerning the performance of the Electric Field Detector (EFD) on board, Huang et al. (2018) studied several natural electromagnetic emissions during the six-month orbit test phase, and the preliminary analysis suggested that the EFD show good performance. Cao et al. (2018) studied the data from the search coil magnetometer (SCM) mounted on CSES that designed to measure the magnetic field fluctuation of low frequency electromagnetic waves ranging from 10 Hz to 20 kHz, they concluded that the performance of SCM can satisfy the
70 requirement of scientific objectives of CSES mission. As one of the main payloads, the GNSS occultation receiver (GOR) had the occultation observation function of both GPS and BDS-2 (Lin et al. 2018). Yan et al. (2020) provided a comprehensive comparison of in situ electron density (Ne) and temperature (Te) measured by Langmuir probe (LAP) on board the CSES with other space-borne and ground-based observations. Their results suggested that the CSES in situ plasma parameters are reliable with a high scientific potential for investigation of geophysics and space. Wang et al. (2019)
75 compared CSES ionospheric RO data with Constellation Observing System for Meteorology, Ionosphere and Climate (COSMIC) measurements. Results indicated that NmF2 and hmF2 between CSES and COSMIC is in extremely good agreement, and co-located electron density profiles (EDPs) between the two sets are generally in good agreement above 200 km.

80 Though the performance of CSES has already been analyzed for different payloads, there are still rooms for an in-depth analysis of GOR, especially for the region with an altitude below 200 km, e.g., E-layer. In addition, as demonstrated by previous studies, the RO measurements can provide a very valuable data for the global sounding of sporadic E layers. In this study we assessed the GRO performance of CSES in the investigating of lower ionosphere, especially the occurrence and properties of sporadic E layers on a global scale.

85

This paper is organized as follows. We first realize the algorithm of sounding sporadic E layers with almost nine months CSES GOR data. Then, we show the results and discussions on global Es-event morphology. Afterward, the comparison of Es altitudes from RO profiles with those from Wuhan ZLT ionosonde measurements reveal a large correspondence between both measurement techniques are introduced. Finally, we present the conclusion.

90 2. Methods

The GOR payload on board CSES can receive the dual frequencies from GPS (L1: 1575.42±10 MHz; L2: 1227.6±10 MHz) and BDS-2 (L1: 1561.98±2 MHz; L2: 1207.14±2 MHz) (Wang et al, 2019). Based on GNSS RINEX format data, we first
~~calculates~~ calculate the electron density profile by occultation inversion algorithm (Lei et al, 2007; Yue et al, 2011), ~~then~~ and
extracts the signal to noise density ratio (SNR) ~~carrier to noise density ratio (CNR)~~ data of L1 and the corresponding time
95 information according to the observation data. Considering the resolution of time and altitude, moving average of 31 points (corresponding to 70-120 km in the vertical direction) is used to calculate the background trend term of SNR ~~CNR~~ data.

After that, we calculate the normalized SNR data and the standard deviation of normalized SNR data. A new criterion is developed to determine whether Es occurs. That is, when the mean value of the absolute value of the difference between the normalized SNR is greater than 3 times the standard deviation, we consider the Es occurs. If more than one value of the normalized SNR sequence meets the conditions, multi-layer Es occurs.

2.1 Sounding of sporadic E layers

Carrier to noise ratio, denoted as CNR or C/N (dB Hz), provides highly desirable information about the quality of the received GNSS signal. Signal to noise ratio, denoted as SNR, is directly related to the C/N measurement and the coherent integration time T_{coh} as:

$$C/N = 10 \log_{10} \left(\frac{SNR}{T_{coh}} \right) \quad (1)$$

where the coherent integration time T_{coh} is used to compute the cross ambiguity function (CAF) as the most effective way to detect weak signals in traditional receivers. The T_{coh} is a known parameter since it is chosen by the receiver.

Signal to noise ratio, denoted as SNR or S/N (dB), which can be estimated to obtain the carrier-to-noise ratio (C/N0) measurement, provides highly desirable information about the quality of the received GNSS signal. (Gómez-Casco, D et al. 2018). The SNR is very sensitive to the electron density changing with altitude, e.g., the sporadic E layer. These vertically small variations in the electron density would lead to phase fluctuation of the GNSS signal which can be observed as a reduction or increase of the signal power at the receiver (Hajj et al., 2002). According to Equation (1), C/N and SNR have the same trend, which can indicate the quality of observation signal, therefore, we can apply C/N data instead of SNR data for sounding of sporadic E layers. According to RINEX Version 2.10 documentation, the numerical magnitude of SNR on L1 and L2 is stored in the S1 and S2 observations in the Level-1 original observations data product of CSES, respectively.

Because SNR data itself also has a certain long-term variation, we need to extract the background trend item in SNR data to obtain the disturbance information after removing the background trend. In this study, the moving average method is used to extract the background trend term of SNR data. The formula is as follows:

$$\overline{X}_k = \frac{X_{k-\frac{N-1}{2}} + \dots + X_k + \dots + X_{k+\frac{N-1}{2}}}{N} \quad (2)(1)$$

where, X_k and \overline{X}_k is the k-th data of the original SNR sequence and after smoothing, and N is the size of the smoothing window. Considering that the original data processed in this study is the original occultation observation data with a sampling rate of 1Hz, so we choose 31 data points as the size of the smooth window.

It is inconvenient to analyze SNR data due to the large value of SNR data, therefore, it has to be first normalized. The calculation formula is as follows:

$$SNR1 = \frac{SNR}{SNR0} \quad (3)(2)$$

where SNR is the original data sequence, SNR0 is the background trend item sequence, and SNR1 is the normalized data sequence.

130

Note that there is no strict standard to judge whether single-layer Es or multi-layer Es occurs. In this study, 70-120 km is selected as the interval to sound the occurrence of Es events. The standard deviation of normalized SNR sequence is calculated:

$$\overline{SNR1} = \sum_{i=1}^n SNR1_i \quad (4)(3)$$

$$std = \sqrt{\sum_{i=1}^n (SNR1_i - \overline{SNR1})^2 / (n - 1)} \quad (5)(4)$$

where $\overline{SNR1}$ is the normalized SNR sequence mean; $SNR1_i$ is the normalized SNR sequence; n is the number of normalized SNR sequences. It is thought that Es occurred once the difference of $SNR1_i$ from the mean is greater than 3 times the standard deviation. While there are multi-layer Es occurs in a single occultation event if multiple $SNR1_i$ meets the judgment criteria.

140

We selected two representative occultation events from CSES observation data as examples to verify the correctness of our Es detection algorithm. The detection of single-layer Es event is shown in Figure 1. The left figure shows the electron density profile of G06 satellite at 06:56UT on August 14, 2018 and the SNR profile. The right figure shows the electron density profile, normalized SNR profile within 60-160 km at the same time, in which the red dotted line is $\overline{SNR1} \pm 3std$ boundary vertical line, it can be seen that there is a $SNR1_i$ whose value exceeds the boundary line and corresponds to the height of abnormal electron density in the figure. According to the normalized SNR sequence, the Es height detected in the figure is 96.49 km. The detection of multi-layer Es events is shown in Figure 2. The left figure shows the electron density profile and the SNR profile of G17 satellite at 20:58 on August 27, 2018. The right figure shows the electron density profile, normalized SNR profile within 60-160 km at the same time. The red dotted line is $\overline{SNR1} \pm 3std$ boundary vertical line, and the Es heights detected in the figure are 73.63 km and 102.76 km respectively.

150

~~The principle of occultation inversion is based on the spherical symmetry assumption.~~ Under the assumptions of spherical symmetry (i.e., assuming only vertical electron density gradients), straight-line propagation and an earth's spherical shape, we calculate the electron density profile by occultation inversion algorithm mainly referring to Lei et al. (2007). These assumptions especially the assumptions of spherical symmetry are frequently not fully accurate for smaller-scale ionospheric phenomena, the calculated electron density values are not accurate and can only describe the approximate numerical distribution. Nevertheless, this study does not attempt to retrieve the absolute accurate electron density values of Es, but show the electron density differences at Es peaks compared to those electron density profiles without the Es phenomenon. Our new criterion is developed to mainly use the normalized SNR to determine the Es events, the electron density profile is only a reference to illustrate the effect of relatively higher electron density at Es on the normalized SNR variation, it is

160

further verified that variance in SNR can be suggested to identify and sound sporadic E layers. There is a certain deviation in the low altitude range by these assumptions, and the electron density calculated by inversion will also have an impact. Compared with the electron density itself, the signal-to-noise ratio is more sensitive to the electron density gradient, the SNR peak height does not fully correspond to the local peak of electron density., and the CNR data does not fully correspond,

165 Therefore, it will affect the inversion height comparison.





Figure 1 Schematic diagram of G06 single-layer Es sounding. The left figure shows the electron density profile of G06 occultation event and the SNR profile at 06:56 on August 14, 2018, the right figure shows the electron density profile and normalized SNR profile within 60-160 km at the same time, and the red dotted line is $\overline{SNR1} \pm 3std$ boundary vertical.



170


Figure 2 Schematic diagram of G17 multi-layer Es sounding. The left figure shows the electron density profile of G17 occultation event and the SNR profile at 06:56 on August 14, 2018, the right figure shows the electron density profile and normalized SNR profile within 60-160 km at the same time, and the red dotted line is $\overline{SNR1} \pm 3std$ boundary vertical.


3. Discussions on global Es-event morphology

175 The GOR measurement of CSES from March 1 to December 1 in 2018 are used in the data analysis. With nearly nine
months of data from CSES, there are 104531 and 12642 electron density profiles obtained from GPS and BDS-2 data of
CSES, respectively. The inversion algorithm is utilized based on the FUSING (FUSing IN Gnss) software (Shi et al. 2018;
Zhao et al. 2018; Gu et al. 2020; Gu et al. 2021). Originally, the FUSING software is developed for high precision real-time
GNSS data processing and multi-sensor navigation, and now it can also be used for atmospheric modeling (Lou et al. 2019;
180 Luo et al. 2020; Luo et al. 2021).



3.1 Distribution of Es occurrence rate for seasons and altitude

The nine-month data have been divided into spring (March, April and May), summer (June, July and August) and autumn
(September, October and November). For each season, we use the altitude resolution of 1 km to count the number of
occultation events which sound Es events in each altitude interval. Due to the resolution of observation values, we do not
185 distinguish the occultation events of sounding Es in different layers. Considering the error caused by the integrity of the
original observation data in different seasons and different days, we count the total number of days with observation data in
each season, then calculate the ratio of the number of occultation events with Es events in different height intervals to the
total number of days in the season, that is counting the number of occultation events with Es events per day. Since CSES has
both GPS and BDS-2 observations, we count the average number of daily occultation events which sound Es events of
190 different satellite systems. The results are as follows:





195 **Figure. 3 Height distribution of Es average daily occurrence rate for three different seasons, from top to bottom are the results with spring, summer and autumn, respectively. The blue dotted line diagram shows the total number of occultation events with Es, the red and green bar chart shows the number of occultation events with Es events per day.**






Figure 3 Height distribution of Es average daily occurrence rate for three different seasons, from top to bottom are the results of spring, summer and autumn, respectively. The blue dotted line diagram shows Es occurrence rate, the red and green bar chart shows the number of occultation events with Es events per day.

205

There are the results with of spring, summer and autumn from top to bottom, respectively. Due to the lack of observation data of CSES for about 20 days in summer, it is not very appropriate to compare seasonal differences only plotting the total number of occultation events with Es. So, as shown in the blue dotted line diagram of Fig.3, we also calculate the ratio of the number of occultation events with Es events in different height intervals to the total number of occultation events in the season.

210

It can be seen from Fig. 3 that the Es average daily occurrence rate has obvious seasonal variation, the height of Es occurrence in spring, summer and autumn is mainly 90-110 km, the height with the largest daily average incidence of Es in spring is 98 km, with a daily average of 2.88, the height with the largest daily average incidence in summer is 99 km, with a daily average of 3.36, and in autumn the height is 101 km, with a daily average of 2.71. The lower total number of occultation events with Es in summer is due to the lack of observation data of CSES for about 20 days in summer. The


215


results show overall distribution of three seasons shifts to a slightly higher height, and more Es also appears above 110 km. The results show significantly more Es appears above 110 km than below 90km overall distribution of three seasons. The reasons, firstly, there are less observation data of CSES at lower altitude, and this situation is reflected in the blue dotted line diagram of Fig. 3; secondly, due to the time resolution, some initial lower altitude values are discarded when using the sliding window to calculate the SNR background trend term, Es occurring at a lower height is also discarded at the same time.

220 **3.2 Distribution of global Es occurrence rate for seasons**

The global longitude and latitude regions are divided into grids with a resolution of $5^{\circ} \times 5^{\circ}$. The number of occultation events in each grid and the number of occultation events with Es events are counted, and the ratio of the number of occultation events with Es to the total number of occultation observations is taken as the Es occurrence frequency of the grid.



225 **In order to reduce the impact of accidental errors, we further optimized the statistical method, the Es occurrence rate for the grid is calculated only when the number of occultation events in the grid is greater than 10.** Finally, the global longitude-latitude distribution characteristics of Es occurrence frequency in this season are obtained. The statistical results are as follows:





230

Figure 4 The geographical distribution of Es occurrence rate for three different seasons in $5^{\circ} \times 5^{\circ}$ geographic latitude/longitude grid, from top to bottom are the results of spring, summer and autumn, respectively.



235




Figure 4 The geographical distribution of Es occurrence rate for three different seasons in $5^{\circ} \times 10^{\circ}$ geographic latitude/longitude grid, from top to bottom are the results of spring, summer and autumn, respectively.

240 There are the results of spring, summer and autumn from top to bottom, respectively. In general, Es preferably occurs at midlatitudes of the summer hemisphere. The overall occurrence frequency of global Es in spring and autumn is lower than that in summer. This phenomenon may be due to the strong solar radiation in summer and the ionization of more metal atoms in the ionosphere, which increases the source of Es and promotes the formation of Es. Therefore, the occurrence rate in mid-latitude of the hemisphere in summer is higher than that in other latitudes (Chu et al. 2014). There is no significant difference in the frequency of Es between the northern and southern hemispheres in spring and autumn, and it shows an almost symmetrical trend along the equator. In spring and autumn, the direct point of the sun is near the equator. Because the magnetic line of force here is almost horizontal, it is difficult to form ion aggregation even if the ionization rate increases, so the occurrence rate is relatively high in the low latitude area of the magnetic equator (Arras et al. 2017; Xue et al. 2018). The Es rates at polar regions are always low. We can also find an occurrence depression around the American area (the longitude sector of 70° – 120° W) in the mid-latitude in summer, where the Es occurrence rates were lower than anywhere else along the zone bands, this is consistent with the phenomenon found by Tsai et al. 2018.

3.3 Distribution of Es occurrence rate for latitude and altitude

To comprehensively analyze the distribution of Es incidence with latitude and altitude, the latitude-altitude region is divided into grids with a resolution of $5^{\circ} \times 10^{\circ} \times 1\text{km}$. Similarly, the ratio of the number of occultation events corresponding to Es events in the grid to the total number of days with observed data in the season is calculated, the daily average number of Es events is taken as the occurrence frequency of Es for statistical analysis. The results are as follows:




Figure 5 The distribution of Es occurrence rate for three different seasons in $5^{\circ} \times 1\text{km}$ geographic latitude/altitude grid, from left to right are the results with of spring, summer and autumn, respectively.

260





Figure 5 The distribution of Es occurrence rate for three different seasons in $10^{\circ} \times 1\text{km}$ geographic latitude/altitude grid, from left to right are the results with of spring, summer and autumn, respectively.

265 There are the results ~~with of~~ spring, summer and autumn from left to right, respectively. It can be seen from the figure that the incidence of Es latitude altitude shows obvious seasonal changes. The incidence of Es in summer in the northern hemisphere is significantly higher than that in spring and autumn in the same latitude range and height range. The latitude range of Es high incidence is 20° - 50° north-south latitude, mainly around 30° . The occurrence height of Es is mainly concentrated in 90-110 km.

270 3.4 Distribution of Es occurrence rate for local time and latitude


In order to comprehensively analyze the distribution of Es incidence with local time and latitude, the local-time-latitude region is divided into grids with a resolution of $1\text{h} \times 5^{\circ}$. ~~Similarly, the ratio of the number of occultation events corresponding to Es events in the grid to the total number of days with observed data in the season is calculated, the daily average number of Es events is taken as the occurrence frequency of Es for statistical analysis.~~ In order to exclude the effect

275 of single-day observation integrity on the distribution of Es incidence with local time. We use the ratio of the number of occultation events with Es to the total number of occultation observations in the grid, at the same time, the Es occurrence rate for the grid is calculated only when the number of occultation events in the grid is greater than 10 to reduce the impact of accidental errors. The results are as follows:



280

Figure 6 The distribution of Es occurrence rate for three different seasons in $1h \times 5^\circ$ local time/geographic latitude grid, from top to bottom are the results with of spring, summer and autumn, respectively.



285




Figure 6 The distribution of Es occurrence rate for three different seasons in 1h×5° local time/geographic latitude grid, from top to bottom are the results with of spring, summer and autumn, respectively.

290

There are the results ~~with of~~ spring, summer and autumn from top to bottom, respectively. It can be seen from the figure that the incidence of Es shows obvious local time changes, the period of local time 15:00-18:00 is the high incidence period of Es in all seasons. This is because the occurrence of Es is directly related to the intensity of solar radiation, the electricity and aggregation of metal ions gradually occur, therefore, there is a delay between the high incidence period of Es and the 12:00 of local time with the strongest solar radiation.

295

4. Experiments of Comparing with Ionosonde measurements

In this study, we choose a certain space-time matching criterion to obtain the pairs of the geometric altitudes of a sporadic E layer detected in CSES radio occultation profiles and the virtual heights of a sporadic E layer obtained by the ZLT ionosonde for confirming the derived sporadic E parameter in height. Luo et al. (2019) choose a certain space-time matching criterion to evaluate the quality of the electron density profile from the FY-3C mission with respect to COSMIC mission. We modified their method to confirm the height of the derived sporadic E layer. We counted the data of Wuhan ZLT ionosonde from March 1 to December 16 in 2018 of the same period, and extracted the h'Es data. The space-time matching criterion is quantified as the size of the space-time window centered on the position and occurrence time of the sporadic E layer obtained by the ZLT ionosonde. The sporadic E layer detected in CSES radio occultation profiles falling into the space-time window and the sporadic E layer obtained by the ZLT ionosonde constitute the pairs participating in the comparative analysis. Here the space-time window is denoted as (B, L, T), where B and L represent the size of space window along latitude and longitude, respectively; T represents the size of the time window.

300

305

310 In this study, considering that the temporal resolution of the ionosonde is 15 minutes, four different space-time matching criteria are proposed with the window as (10°, 10°, 7.5 min), (5°, 10°, 7.5min), and (5°, 5°, 7.5min) and (2°, 5°, 7.5min), respectively. Among the other parameters, the height of sporadic E layer is an important parameter of the derived sporadic E layer. Thus, the correlation coefficient (CC), is derived for determining the height of sporadic E layer. The definition of the correlation coefficient is presented below.

$$315 \quad CC = \frac{\sum_{i=1}^N (x_i^C \cdot x_i^Z) - \frac{1}{N} \sum_{i=1}^N x_i^C \sum_{i=1}^N x_i^Z}{\sqrt{(\sum_{i=1}^N (x_i^C)^2 - \frac{1}{N} (\sum_{i=1}^N x_i^C)^2)(\sum_{i=1}^N (x_i^Z)^2 - \frac{1}{N} (\sum_{i=1}^N x_i^Z)^2)}} \quad (6)$$


where N represents the total number of data pairs in the matching group under a given spatiotemporal matching windows; $x_i^C (i=1,2,3 \dots, n)$ represents the geometric altitudes of *i*th sporadic E layer detected in CSES radio occultation profiles; $x_i^Z (i=1,2,3 \dots, n)$ represents the virtual heights of *i*th sporadic E layer obtained by the ZLT ionosonde.

320 The data of ionosonde is stored in SAO format files, this data file contains different types of parameters, such as station information and detection time, ionospheric characteristic parameters for automatic measurements, echo traces (virtual height, amplitude, doppler, frequency) at different height layers of the ionosphere (E, F1, F2), electron density profiles, virtual height and critical frequency of Es trace, etc. The SAO format description can refer to <https://ulcar.uml.edu/~iag/SAO-4.htm>. In order to facilitate the reading and use of data, SAOExplorer software
325 (<http://ulcar.uml.edu/SAO-X/SAO-X.html>) has been developed by the Center for Atmospheric Research at the University of Massachusetts Lowell, USA, to display and measure Digisonde Ionospheric frequency map observed by a series of ionospheric altimeters.


Figure 7 shows an example of simultaneous detecting Es by CSES and ZLT ionosonde, the left figure shows the electron
330 density profile of G13 satellite at 07:02 on August 15, 2018 and the electron density profile of ZLT ionosonde at 07:00 on August 15, 2018, the right figure shows the electron density profile in the range of 0-200 km. In the figure, the geodetic coordinates of Es detected by CSES is (39.5°N, 115.2°E, 102.990 km) and the geodetic coordinates of Es detected by ZLT is (30.5°N, 114.4°E, 102.994 km).

335 Figure 7 shows an example of simultaneous detecting Es by CSES and ZLT ionosonde, the top left figure shows the electron density profile and the SNR1 profile of G27 satellite at 17:42 on May 17, 2018, and the electron density profile of ZLT ionosonde at 17:45 on May 17, 2018, the top right figure shows the electron density profile of ZLT ionosonde and the SNR1 profile in the range of 0-220 km. In the figure, the geodetic coordinates of Es detected by CSES is (33.0°N, 112.3°E, 99.2 km) and the geodetic coordinates of Es detected by ZLT is (30.5°N, 114.4°E, 102.5 km). The bottom figure shows the


340 ionogram image of Wuhan ZLT ionosonde to show Es situation. We can obtain the virtual height of Es is 102.5km, we can also obtain Es layer critical frequency and frequency map at about 100km.



345 **Figure 7** An example of simultaneous detecting Es by CSES and ZLT ionosonde, the left figure shows the electron density profile of G13 satellite at 07:02 and ZLT ionosonde at 07:00 on August 15, 2018, the right figure shows the electron density profile in the range of 0-200 km.



foF2	3.650
foF1	N/A
foF1p	N/A
foE	N/A
foEp	0.36
fxI	4.55
foEs	3.10
fmin	1.60
<hr/>	
MUF(D)	10.78
M(D)	2.95
D	N/A
<hr/>	
h'F	290.0
h'F2	290.0
h'E	N/A
h'Es	102.5
<hr/>	
hmF2	331.4
hmF1	N/A
hmE	110.0
yF2	76.0
yF1	N/A
yE	20.0
B0	66.3
B1	3.44
<hr/>	
C-level	44
<hr/>	
Auto:	
Artist5	
500200	



D	100	200	400	600	800	1000	1500	3000	[km]
MUF	4.3	4.3	4.4	4.7	5.0	5.6	7.0	10.8	[MHz]

WU430_2018137174500.SBF / 190Eh512h 50 kHz 2.5 km / DPS-4D WU430 030 / 30.5 N 114.4 E

Ion2Png v. 1.3.16

350

Figure 7 An example of simultaneous detecting Es by CSES and ZLT ionosonde, the top left figure shows the electron density profile and the SNR1 profile of G27 satellite at 17:42 on May 17, 2018, and the electron density profile of ZLT ionosonde at 17:45 on May 17, 2018, the top right figure shows the electron density profile of ZLT ionosonde and the SNR1 profile in the range of 0-220 km. The bottom figure shows the ionogram image of Wuhan ZLT ionosonde.

355

Figure 8 presents the comparison of the geometric altitudes of a sporadic E layer detected in CSES radio occultation profiles and the virtual heights of a sporadic E layer obtained by the ZLT ionosonde. We also show the regression line as the solid black line and corresponding statistical coefficients in every subgraph. In the fourth subgraph, there are two points at 111.25 km of the ionosonde data, corresponding to 106.37 km and 106.40 km of CSES data respectively, with small difference and almost overlapping in the figure.

360

These figures reveals that there is a good agreement between both parameters, which can also be seen from the high correlation larger than 0.7. The comparison among different windows conclude that the correlation increased slightly as a stricter space-time matching window involved, but with less pairs or couples. Compared with results from Arras et al. (2017), we also found a height offset between both measurement techniques mainly concentrated in 100-110 km of ionosonde altitude, and the calculation results of different space-time windows are different.

365

The mean offset values in 100-110km are 2.36km, 2.25km, and 2.90km, which correspond to space-time matching windows

(10°, 10°, 7.5 min), (5°, 10°, 7.5min), and (5°, 5°, 7.5min), respectively. This may result from the different height parameters used for both techniques: the geometric heights provided by the RO technique and the virtual height which is influenced by the ionization below the sporadic E layer calculated from ionosonde recordings (Arras et al. 2017).





Figure 8 Comparison of the geometric altitudes of Es detected in CSES radio occultation profiles and the virtual heights of Es obtained by the ZLT ionosonde. From top to bottom and from left to right are the results with space-time matching window (10°, 10°, 7.5 min), (5°, 10°, 7.5min), (5°, 5°, 7.5min) and (2°, 5°, 7.5min), respectively.

370

375



380 **Figure 8 Comparison of the geometric altitudes of Es detected in CSES radio occultation profiles and the virtual heights of Es obtained by the ZLT ionosonde. From top to bottom are the results of space-time matching window (10°, 10°, 7.5 min), (5°, 10°, 7.5min) and (5°, 5°, 7.5min), respectively. The black solid line is the regression line.**

5. Conclusions

The RO plays an important role in sounding of sporadic E layers. As China's first electromagnetic satellite, CSES has already providing service for more than three years up to now. In this study, the Level-1 data of CSES and Wuhan ZLT ionosonde from March 1 to December 1 in 2018 are collected in sounding of sporadic E layers used to study the comparison of heights.

We calculate the geodetic longitude; latitude and elevation of each occultation tangent point in the occultation inversion process, and count the corresponding time information, then extract the SNR data of L1 observations in the occultation inversion period. The occurrence of Es is judged according to the judgment criteria of $|SNR1_i - \overline{SNR1}| > 3std$. Single layer or multi-layer es is judged according to the number of data whose sequence meets the judgment criteria. Combined with the electron density profile of occultation inversion, the correctness of our Es detection algorithm is verified.

According to the Es results we detected, we drew distribution of Es occurrence rate for seasons and altitude, distribution of global Es occurrence rate for seasons. It is concluded that the occurrence height of Es is mainly located at 90-110 km, and there are obvious seasonal and latitudinal changes in the occurrence rate of Es. There is no significant difference in the occurrence frequency of Es in the northern and southern hemispheres in spring and autumn, and it is almost symmetrical along the equator. Summer in the northern hemisphere is the time period of high incidence of Es, and the latitude range of high incidence of Es is 20°-50° in the northern and southern latitudes, mainly around 30°. The period of local time 15:00-18:00 is the high incidence period of Es.

Finally, the comparison of the geometric altitudes of sporadic E layers detected in CSES radio occultation profiles and the virtual heights of sporadic E layers obtained by the ZLT ionosonde was carried out with for different space-time matching window, i.e., (10°, 10°, 7.5 min), (5°, 10°, 7.5min), and (5°, 5°, 7.5min) and (2°, 5°, 7.5min). For these four windows, the number of CSES matched pairs was 37, 26, 14 and 5, respectively. The correlation coefficients of altitudes were 0.707, 0.736, and 0.748 and 0.884, respectively. The comparison of Es altitudes from RO profiles with those from coinciding ground based ionosonde measurements revealed a large correspondence between both measurement techniques.

Data availability

CSES Radio Occultation data can be downloaded from <http://www.leos.ac.cn> and email author Shengfeng Gu. The Wuhan ZLT ionosonde observations can be downloaded from <https://data.meridianproject.ac.cn/>.

Author contributions

XL, CX and SG designed the research; CG and JH performed the research; CG, JH and SG analyzed the data; CG drafted the paper. XL, CX and SG put forward valuable modification suggestions. All authors contributed by providing the necessary data and discussions and writing the paper.

415 Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

CSES Radio Occultation data can be downloaded from <http://www.leos.ac.cn>. The authors express their thanks. We also acknowledge the use of data of Wuhan ZLT ionosonde from the Chinese Meridian Project.

420 Financial support

This research has been supported by the National Key R&D Program of China (grant no. 2018YFC1503502). This work is also supported by the National Natural Science Foundation of China (No. 42104029).

References

- Ambrosi, G., Bartocci, S., Basara, L. et al. Battiston, R., Burger, W. J., Carfora, L., Castellini, G., Cipollone, P., Conti, L.,
425 Contin, A., Donato, C. D., Santis, C. D., Follega, F. M., Guandalini, C., Ionica, M., Iuppa, R., Laurenti, G., Lazzizzera, I.,
Lolli, M., Manea, C., Marcelli, L., Masciantonio, G., Mergé, M., Osteria, G., Pacini, L., Palma, F., Palmonari, F., Panico, B.,
Patrizii, L., Perfetto, F., Picozza, P., Pozzato, M., Puel, M., Rashevskaya, I., Ricci, E., Ricci, M., Ricciarini, S. B., Scotti, V.,
Sotgiu, A., Sparvoli, R., Spataro, B. and Vitale, V.: The HEPD particle detector of the CSES satellite mission for
investigating seismo-associated perturbations of the Van Allen belts, *Sci. China Technol. Sci.* 61, 643–652,
430 <https://doi.org/10.1007/s11431-018-9234-9>, 2018.
- Arras, C., Wickert, J., Beyerle, G., Heise, S., Schmidt, T., and Jacobi, C.: A global climatology of ionospheric irregularities
derived from GPS radio occultation, *Geophys. Res. Lett.*, 35: L14809. <https://doi.org/10.1029/2008G L034158>, 2008.
- Arras, C., and Wickert, J.: Estimation of ionospheric sporadic E intensities from GPS radio occultation measurements, *J.*
Atmos. Sol-Terr. Phys., 171:60–63. <https://doi.org/10.1016/j.jastp.2017.08.006>, 2017.
- 435 Axford, W. I.: The formation and vertical movement of dense ionized layers in the ionosphere due to neutral wind shears, *J.*
Geophys. Res., 68, 769–779. <https://doi.org/10.1029/JZ068i003p00769>, 1963.

- Cao, J. B., Zeng, L., Zhan, F., Wang, Z. G., Wang, Y., Chen, Y., Meng, Q. C., Ji, Z. Q., Wang, P. F., Liu, Z. W., and Ma, L. Y.: The electromagnetic wave experiment for CSES mission: Search coil magnetometer, *Sci. China. Tech. Sci*, 61, 653–658, <https://doi.org/10.1007/s11431-018-9241-7>, 2018.
- 440 Chu, Y. H., Wang, C. Y., Wu, K. H., Chen, K. T., Tzeng, K. J., Su, C. L., Feng, W., and Plane, J. M. C.: Morphology of sporadic E layer retrieved from COSMIC GPS radio occultation measurements: wind shear theory examination, *J. Geophys. Res.*, 119:2117–2136. <https://doi.org/10.1002/2013JA019437>, 2014.
- Didebulidze, G. G., Dalakishvili, G., and Todua, M.: Formation of Multilayered Sporadic E under an Influence of Atmospheric Gravity Waves (AGWs), *Atmosphere.*, 11(6), 653. <https://doi.org/10.3390/atmos11060653>, 2020.
- 445 Gómez-Casco, D., López-Salcedo, J. A., and Seco-Granados, G.: C/N0 estimators for high-sensitivity snapshot GNSS receivers, *GPS Solut.*, 22, 122. <https://doi.org/10.1007/s10291-018-0786-y>, 2018.
- Gu, S., Dai, C., Fang, W., Zheng, F., Wang, Y., and Zhang, Q.: Multi-GNSS PPP/INS tightly coupled integration with atmospheric augmentation and its application in urban vehicle navigation, *J. Geod.*, 95(6), –15. <https://doi.org/10.1007/s00190-021-01514-8>, 2021.
- 450 Gu, S., Wang, Y., Zhao, Q., Zheng, F., and Gong, X.: BDS-3 differential code bias estimation with undifferenced uncombined model based on triple-frequency observation, *J. Geod.*, 94, 45. https://doi.org/10.1007/s00190-020-01364-w_2, 2020.
- Haldoupis, C.: A Tutorial Review on Sporadic E Layers. In: Abdu M., Pancheva D. (eds) *Aeronomy of the Earth's Atmosphere and Ionosphere*. IAGA Special Sopron Book Series, vol 2. Springer, Dordrecht. https://doi.org/10.1007/978-94-007-0326-1_29, 2011.
- 455 Haldoupis, C.: Midlatitude Sporadic E. A Typical Paradigm of Atmosphere-Ionosphere Coupling, *Space. Sci. Rev.*, 168, 441–461. <https://doi.org/10.1007/s11214-011-9786-8>, 2012.
- Hajj, G. A., Kursinski, E. R., Romans, L. J., Bertiger, W. I., and Leroy., S. S.: A technical description of atmospheric sounding by GPS occultation, *J. Atmos. Sol-Terr. Phys.*, 64.4: [https://doi.org/10.1016/S1364-6826\(01\)00114-6](https://doi.org/10.1016/S1364-6826(01)00114-6), 2002.
- 460 Hocke, K., Igarashi, K., Nakamura, M., Wilkinson, P., Wu, J., Pavelyev, A., and Wikert, J.: Global sounding of sporadic E layers by the GPS/MET radio occultation experiment, *J. Atmos. Solar-Terr. Phys.*, 63, 1973–1980. [https://doi.org/10.1016/S1364-6826\(01\)00063-3](https://doi.org/10.1016/S1364-6826(01)00063-3), 2001.
- Huang, J. P., Lei, J. G., Li, S. X., Zeren, Z. M., Li, C., Zhu, X. H., and Yu, W. H.: The Electric Field Detector (EFD) onboard the ZH-1 satellite and first observational results, *Earth. Planet. Phys.*, 2(6), 469–478. <https://doi.org/10.26464/epp2018045>, 2018.
- 465 Lei, J., Syndergaard, S., Burns, A. G., Solomon, S. C., Wang, W., Zeng, Z., Roble, R. G., Wu, Q., Kuo, Y., Holt, J. M., Zhang, S., Hysell, D. L., Rodrigues, F. S., and Lin, C. H.: Comparison of COSMIC ionospheric measurements with ground-based observations and model predictions: Preliminary results, *J. Geophys. Res.*, 112, A07308, <https://doi.org/10.1029/2006JA012240>, 2007.

- 470 Lin, J., Shen, X. H., Hu, L. C., Wang, L. W., and Zhu, F. Y.: CSES GNSS ionospheric inversion technique, validation and error analysis, *Sci. China Tech. Sci.*, 61: 669–677, <https://doi.org/10.1007/s11431-018-9245-6>, 2018.
- Lou, Y., Luo, X., Gu, S., Xiong, C., Song, Q., Chen, B., Chen, B., Xiao, Q., Chen, D., Zhang, Z., Zheng, G.: Two Typical Ionospheric Irregularities Associated With the Tropical Cyclones Tembin (2012) and Hagibis (2014), *ns. J. Geophys. Res. Space. Phys.*, 124(7), 6237–6252. <https://doi.org/10.1029/2019JA026861>, 2019.
- 475 Luo, J., Wang, H., Xu, X., and Sun, F.: The Influence of the Spatial and Temporal Collocation Windows on the Comparisons of the Ionospheric Characteristic Parameters Derived from COSMIC Radio Occultation and Digisondes, *Adv. Space Res.*, 63:3088–3101. <https://doi.org/10.1016/j.asr.2019.01.024>, 2019.
- Luo, X., Gu, S., Lou, Y., Cai, L., and Liu, Z.: Amplitude scintillation index derived from C/N0 measurements released by common geodetic GNSS receivers operating at 1 Hz, *J. Geod.*, 94(2), 1–14. <https://doi.org/10.1007/s00190-020-01359-7>,
480 2020.
- Luo, X., Lou, Y., Gu, S., Li, G., Xiong, C., Song, W., and Zhao, Z.: Local ionospheric plasma bubble revealed by BDS Geostationary Earth Orbit satellite observations, *GPS Solut.*, 25(3), –10. <https://doi.org/10.1007/s10291-021-01155-6>, 2021.
- Mathews J.D.: Sporadic E: current views and recent progress, *J. Atmos. Sol.-Terr. Phys.* 60, 413. [https://doi.org/10.1016/S1364-6826\(97\)00043-6](https://doi.org/10.1016/S1364-6826(97)00043-6), 1998.
- 485 Shen, X. H., Zong, Q. G., and Zhang, X. M.: Introduction to special section on the China Seismo-Electromagnetic Satellite and initial results. *Earth Planet. Phys.*, 2(6), 439–443. <https://doi.org/10.26464/epp2018041>, 2018.
- Shi, C., Guo, S., Gu, S., Yang, X., Gong, X., Deng, Z., Ge, M., and Schuh, H.: Multi-GNSS satellite clock estimation constrained with oscillator noise model in the existence of data discontinuity, *J. Geod.*, 93, 515–528. <https://doi.org/10.1007/s00190-018-1178-3>, 2019.
- 490 Fytterer, T., Arras, C., Hoffmann, P., and Jacobi, C.: Global distribution of the migrating terdiurnal tide seen in sporadic E occurrence frequencies obtained from GPS radio occultations. *Earth Planet Sp.*, 66, 79. <https://doi.org/10.1186/1880-5981-66-79>, 2014.
- Tsai, L. C., Su, S. Y., Liu, C. H., Schuh, H., Wickert, J., and Alizadeh, M. M.: Global morphology of ionospheric sporadic E layer from the FormoSat-3/COSMIC GPS radio occultation experiment. *GPS Solut.*, 22, 118. <https://doi.org/10.1007/s10291-018-0782-2>, 2018.
- 495 Wang, X., Cheng, W., Zhou, Z., Xu, S., Yang, D., and Cui, J.: Comparison of CSES ionospheric RO data with COSMIC measurements, *Ann. Geophys.*, 37, 1025–1038, <https://doi.org/10.5194/angeo-37-1025-2019>, 2019.
- Whitehead, J. D.: Recent work on mid-latitude and equatorial sporadic-E, *J. Atmos. Terr. Phys.*, 51(5), 401–424. [https://doi.org/10.1016/0021-9169\(89\)90122-0](https://doi.org/10.1016/0021-9169(89)90122-0), 1989.
- 500 Whitehead, J. D.: The formation of the sporadic-E layer in the temperate zones, *J. Atmos. Terr. Phys.*, 20, 49–58. [https://doi.org/10.1016/0021-9169\(61\)90097-6](https://doi.org/10.1016/0021-9169(61)90097-6), 1961.

- Wickert, J., Pavelyev, A. G., Liou, Y. A., Schmidt, T., Reigber, C., Igarashi, K., Pavelyev, A. A., and Matyugov, S.: Amplitude variations in GPS signals as possible indicator of ionospheric structures, *Geophys. Res. Lett.*, 31, L24801, <https://doi.org/10.1029/2004GL020607>, 2014.
- 505 Wu, D. L., Ao, C. O., Hajj, G. A., de la Torre Juarez, M., and Mannucci, A. J.: Sporadic E morphology from GPS-CHAMP radio occultation, *J. Geophys. Res.*, 110, A01306. <https://doi.org/10.1029/2004JA010701>, 2005.
- Xue, Z. X., Yuan, Z. G., Liu, K., and Yu, X. D.: Statistical research of Es distribution in inland areas of China based on COSMIC occultation observations. *Chinese J. Geophys.* (in Chinese), 61(8): 3124-3133. <https://doi.org/10.6038/cjg2018L0670>, 2018.
- 510 Yan, R., Zhima, Z., Xiong, C., Shen, X., Huang, J., Guan, Y., Zhu, X., and Liu, C.: Comparison of electron density and temperature from the CSES satellite with other space-borne and ground-based observations. *J. Geophys. Res. Space. Phys.*, 125, e2019JA027747. <https://doi.org/10.1029/2019JA027747>, 2020.
- Yeh, W. H., Huang, C. Y., Hsiao, T. Y., Chiu, T. C., Lin, C. H., and Liou, Y. A.: Amplitude morphology of GPS radio occultation data for sporadic-E layers, *J. Geophys. Res.*, 117, A11304. <https://doi.org/10.1029/2012JA017875>, 2012.
- 515 Yeh, W. H., Liu, J. Y., Huang, C. Y., and Chen, S. P.: Explanation of the sporadic-E layer formation by comparing FORMOSAT-3/COSMIC data with meteor and wind shear information, *J. Geophys. Res. Atmos.*, 119, 4568–4579, <https://doi.org/10.1002/2013JD020798>, 2014.
- Yue, X., Schreiner, W. S., Rocken, C., and Kuo, Y. H.: Evaluation of the orbit altitude electron density estimation and its effect on the Abel inversion from radio occultation measurements. *Radio Sci.*, 46, RS1013. <https://doi.org/10.1029/2010RS004514>, 2011.
- 520 Yue, X., Schreiner, W. S., Zeng, Z., Kuo, Y. H., and Xue, X.: Case study on complex sporadic E layers observed by GPS radio occultations, *Atmos. Meas. Tech.*, 8, 225–236, <https://doi.org/10.5194/amt-8-225-2015>, 2015.
- Zhao, Q., Wang, Y., Gu, S., Zheng, F., Shi, C. Ge, M., and Schuh, H.: Refining ionospheric delay modeling for undifferenced and uncombined GNSS data processing, *J. Geod.*, 93, 545–560. <https://doi.org/10.1007/s00190-018-1180-9>, 2019.
- 525 Zhou, X., Yue, X., Liu, H. L., Lu, X., Wu, H., Zhao, X., and He, J.: A comparative study of ionospheric day-to-day variability over Wuhan based on ionosonde measurements and model simulations, *J. Geophys. Res. Space. Phys.*, 126, e2020JA028589. <https://doi.org/10.1029/2020JA028589>, 2021.