

Responses to the Comment and/or Suggestions from Referee 1

Comments on "Effects of the terdiurnal tide on the Sporadic-E layers (Es) development at low latitudes over the Brazilian sector" by Pedro Alves Fontes et al., Ann. Geophys. Discuss., <https://doi.org/10.5194/angeo-2022-17-RC1>, 2022.

Using the ionosonde and meteor radar observations and the MIRE model, this study investigated the effects of terdiurnal tide on the Es layers at low latitudes in the Southern Hemisphere. The paper suggested that the terdiurnal tide could help in the formation of the Es layers in the Brazilian sector. The results could improve our understanding of the formation of Es layers in the lower latitudes. However, they are major issues that need to be clarified first.

Thank you very much for the revision given by the referee. We have carried out a revision of the manuscript considering all the referee's comments.

Comments:

1. The production and loss terms used in the model should be given.

Response:

Ok. We include information and Table 1 with the equations and rate coefficients in this new version (See lines 124 to 129).

2. The paper mentioned that the wind data used in MIRE were obtained by the meteor radar. The altitude range for the meteor radar is from 80 to 100 km while the height range of interest in this study is between 86 and 140 km. The authors mentioned that they expanded the wind equation to 120 km. It is not clear to me how the authors obtain the neutral wind data from 100 to 140 km and the accuracy of the wind data in this altitude range. The neutral wind data is very important in simulating the formation of the Es layers.

Response:

We apologize for not making this clear. In fact, the measurements available from the meteor radar provide data from 80 to 100 km only. However, the height range of interest

in MIRE to simulate the E region and the Es layer dynamics is from 86 to 140 km. Thus, to extend the wind equations up to 140 km, we used a fitting function for each parameter as proposed in Resende et al. (2017a; 2017b). In this work, the Lorentz curve (according to the equation below) was used in this study for wind amplitude, considering the theory about the wind behavior (Lindzen and Chapman, 1969; Forbes and Garret, 1979).

$$U_{(x_0,y_0)}^{(24,12,8)}(z) = \mu_{(x_0,y_0)} + \frac{2 \cdot A_{(x_0,y_0)}}{\pi} \cdot \frac{\varphi_{(x,y)}}{4 \cdot (z - h_{(x_0,y_0)})^2 + \varphi_{(x,y)}^2},$$

where $\mu_{(x_0,y_0)}$, $A_{(x_0,y_0)}$, $\varphi_{(x,y)}$ and $h_{(x_0,y_0)}$ are the fitted parameters for the meridional and zonal components of the daytime (24 h), semidiurnal (12 h) and terdiurnal (8 h) tides. Additionally, the wind phases were obtained by a linear fitting of the observational data, for São João do Cariri region. And the vertical wavelengths can be used from the respective wave phase equations by multiplying the linear coefficient of the fitted curves by the corresponding tidal period 24 h for diurnal tide, 12 h for the semidiurnal tide, and 8 for the terdiurnal (Buriti et al., 2008). It is important to mention here that Conceição-Santos et al. (2019) proposed a modification in the $\lambda_{x,y}$ of the extrapolated winds, keeping them constant between 100 and 120 km, and tending to infinity between 120 and 140 km. A detailed description about the radar system and the methodology used to compute the neutral wind parameters is available in Resende et al. (2017a). Finally, we used the Es layer density profiles of the MIRE for the months of December (summer), April (autumn), July (winter) and October (spring) as representative of the seasons.

All these information was included in lines 143 to 156 in this new version.

3. I think that to study the seasonal variation, multiple years of data are needed. This paper only used one year of data. I suggest that the authors use a larger dataset.

Response:

We appreciate and understand what the authors pointed out. However, in this study, we believe that one year is enough to have an idea of the Es layer behavior, as seen in Resende et al. (2017a; 2017b) and Conceição-Santos et al. (2019). The tidal wind pattern does not have significant changes each year. The GSWM model, for example, gives us the monthly parameters representing all the years. Furthermore, the inclusion of more years of data

takes a long time to analyze since the interpretation and reduction of the ionosonde data are not carried out from automatic program. Finally, the purpose of this study is to show that in addition to the diurnal and semidiurnal tides, there is also an influence of the terdiurnal tide in the Es layer formation over the Brazilian station. We highly respect the reviewer's opinion, but we believe that our results are sufficient to analyze the development of the Es layers in this low latitude sector.

4. Figures 1 to 5 are not new to me. At least, the authors should make comparisons with previous studies.

Response:

We thank the reviewer for the suggestions. Regarding Figures 1 and 2, we have merged them into a single figure (classified as Figure 1 now). In the discussion about Figure 1, we added other references to improve text. Also, we discussed the importance of analyzing the Es layers concerning the types, showing the mechanism of Es layer formation at latitude. Additionally, we included a discussion of the Esc and Esh types (lines 201 to 213).

Also, with respect to Figure 2 (before Figure 3), we have added discussions showing the relationship of the h'Es with solar activity and the terdiurnal tide (lines 231 to 240).

Figures 4 and 5 have been merged into a single figure. It is now Figure 3. In this figure, we kept the discussion by highlighting the months that best represented the 8-hour oscillations in the sum of the Es layer types in each season.

5. Figure 8: the comparisons should be made among observations, simulation1 (D+S), and simulation2 (D+S+T).

Response:

Ok! At the end of the discussion of this figure (now Figure 6), we have added a table where we show a clear comparison of the densities of the Es layers simulated in MIRE with the D+S and D+S+T components and the maximum daily average density for those months found in the ionosonde data. In the last column of Table 2, we also included the maximum average daily densities for the months that had the most pronounced peak. Additionally, we discussed these results in this new version (lines 363 to 377).

6. Terdiurnal tides in the low latitudes have been widely studied. The literature research could be done better.

Response:

We thank the reviewer for the suggestion. The references about the terdiurnal tide have been improved in the introduction, with the addition of two paragraphs (lines 40 to 73) where we show the two theories of tidal generation currently discussed in the literature. We also added new discussions to the results and added some more references on the topic.

Finally, we would like to thank Referee 1 for the suggestions and corrections to improve the manuscript.

References

- Buriti, R. A., Hocking, W. K., Batista, P. P., Medeiros, A. F., and Clemesha, B. R. Observations of equatorial mesospheric winds over Cariri (7.4° S) by a meteor radar and comparison with existing models, *Annales Geophysicae*, 26, 485–497, <https://doi.org/10.5194/angeo-26-485-2008>, 2008.
- Conceição-Santos, F., Muella, M. T. A. H., Resende, L. C. A., Fagundes, P. R., Andrioli, V. F., Batista, P. P., Pillat, V. G., and Carrasco, A. J. Occurrence and Modeling Examination of Sporadic- E Layers in the Region of the South America (Atlantic) Magnetic Anomaly, *Journal of Geophysical Research: Space Physics*, 124, 9676–9694, <https://doi.org/10.1029/2018JA026397>, 2019.
- Forbes, J. M., and Garrett, H. B. Theoretical studies of atmospheric tides. *Reviews of Geophysics*, 17, 1951-1981, <https://doi.org/10.1029/RG017i008p01951>, 1979.
- Lindzen, R. S., and Chapman, S. Atmospheric tides, *Space Science Reviews*, 10, 1415–1422, <https://doi.org/10.1007/BF00171584>, 1969.
- Resende, L. C. A., Batista, I. S., Denardini, C. M., Batista, P. P., Carrasco, A. J., Andrioli, V. F., and Moro, J. The influence of tidal winds in the formation of blanketing sporadic e-layer over equatorial Brazilian region, *Journal of Atmospheric and Solar-Terrestrial Physics*, 171, 64–71, <https://doi.org/10.1016/j.jastp.2017.06.009>, 2017a.
- Resende, L. C. A., Batista, I. S., Denardini, C. M., Batista, P. P., Carrasco, A. J., Andrioli, V. de F., and Moro, J. Simulations of blanketing sporadic E-layer over the Brazilian sector

driven by tidal winds, *Journal of Atmospheric and Solar-Terrestrial Physics*, 154, 104–114, <https://doi.org/10.1016/j.jastp.2016.12.012>, 2017b.