

Referee 2

1. General comments

This manuscript proposes a method for generalizing the mapping of flow lines and magnetic field in the magnetosheath. The proposed method aims to be computationally inexpensive and generalizable, which is desirable for many applications including statistical studies. However, a number of critical issues need to be addressed if the manuscript is to be considered for publication. The results presented are not general enough to be of use for actual applications; thus, the manuscript represents no (or very minor) scientific advancement. The novelty of the study needs to be explained, in particular by a more focused comparison with previous works. Furthermore, the method should be presented more clearly; it is currently difficult to tell if the proposed method is incorrect or if the presentation is unclear and contains too many mistakes (typos and inconsistencies). Please find below my detailed comments and suggestions for improvement.

Reply (ref.02.01):

- Thank you for suggestion that the novelty should be clear in the manuscript. We compare the grid pattern between the radial mapping by Soucek and Escoubet (2012) (figure 1, page 5) and our magnetopause-normal mapping (figure 3, page 6). The radial mapping has the problem of artificial converging flow pattern in the flank region (see figure 2 right panel, page 5). This is due to the strong grid non-orthogonality. Our method can properly transform the scalar-potential and there is no artificial converging flow pattern (figure 11, page 15).

2. Specific comments

Concerning the whole manuscript

Coordinate system: The authors introduce a new, non-parabolic coordinate system (u,v,ϕ) in which the potentials for the velocity and magnetic field are to be expressed. However, it is not mentioned whether the new coordinate system is orthogonal; in fact, from Fig. 3 it appears that the grid in the right panel is not. The parabolic coordinate system (used in KF94) is orthogonal by construction, and the gradient and Laplace equation in this system are defined and given explicitly (Eq. (8)-(9) in the KF94 paper). In this manuscript, however, the potentials are obtained from the shell and connector variables v and u in the new coordinate system (according to line 274-285 and 289). The authors do not define the gradient and curl in this system and thus the evaluation of Eq. (1) is not defined in the manuscript.

Reply (ref.02.02):

- The coordinates with u and v are orthogonal around the magnetopause, and the grid non-orthogonality is suppressed in our method (section 3.3, page 5 and figure 3, page 6;). To make it fully orthogonal (curvilinear

and locally orthogonal grids) one needs to find a conformal mapping. Mathematical studies conclude that there is no unique conformal mapping to the magnetosheath problem because the magnetosheath is bounded only by the magnetopause and the bow shock in the radial direction but not along the streamlines.

- The azimuthal coordinate ϕ is still orthogonal to the u and v coordinates (page 5, lines 111–112).
- The computation of the u and v variables and their gradient and curl is performed in the Cartesian so that the connection represented by the Christoffel symbol vanish in the computation. Computation in the Cartesian domain is also beneficial to the practical application because spacecraft trajectories are often represented in the Cartesian coordinates (page 14, lines 283–286).

3. *Generalization of the method: In section 3, the mapping algorithm is reduced to an axisymmetric geometry and the calculations are made specifically for the Farris (1991) bow shock and Shue (1997) models. Yet, it is stated in sections 1 and 4 that the method is easy to generalize to an arbitrary magnetopause shape. Would it be possible to express the derivations in more general terms? I suggest giving expressions on a form which allows for non-axisymmetric geometries and aberrated GSE coordinate systems (see for example the asymmetric magnetosheath thickness and aberrated x axis in <https://doi.org/10.1002/jgra.50465>), or otherwise indicate to the reader which modifications and transformations are necessary to generalize the method. Are there restrictions on the choice of magnetopause model? Does the method require an analytic expression for the magnetopause?*

Reply (ref.02.03):

- Non-axisymmetric case. It is possible to obtain the steady-state magnetosheath potential in a more general sense without referring to the KF94 solution. For example, for a non-axisymmetric geometry of magnetosheath (e.g., Dimmock and Nykyri, 2013), one needs to solve the Laplace equation for a given set of boundaries (bow shock and magnetopause). Various numerical solvers are known for solving the Laplace equation such as the Jacobi method, the Gauss-Seidel method, and the successive over-relaxation (SOR) method. These Laplace solvers are numerically more expensive than the mapping method, but the computation in 3-D is feasible with the contemporary computational resources (page 16, lines 327–332).
- Magnetopause model. There is no restriction regarding the choice of the magnetopause model. The magnetopause-normal direction needs to be computed either analytically using the gradient of the magnetopause function as ∇f_{mp} , or numerically for a user-defined magnetopause shape (page 16, lines 335–337).

4. *Figures 1 and 2 show examples of results of the mapping, but not in which way the proposed method is better than (or even different from) the method*

by Soucek (2012). They are also introduced very early in the manuscript, before the potential functions or the mapping procedure have been described, and do not contain much relevant information. I suggest replacing them with figures showing the steps of the mapping procedure and/or a comparison with the Soucek (2012) method.

Reply (ref.02.04):

- Figure 1 and figure 3 compare directly the quality of grid pattern between the method by Soucek and Escoubet (2012) and ours. The artificially introduced converging flow pattern in the radial mapping method is shown in figure 2 (page 6).
- Agreed. Stepwise figures are shown in the revision (figures 5–9).

5. *Title*

The phrase “Potential mapping method” can be interpreted as “Possible mapping method”. Perhaps the title could be rephrased to avoid misunderstanding.

Reply (ref.02.05):

- Agreed. We change the title into “Scalar-potential mapping of the steady-state magnetosheath model” to avoid confusion (title field, page 1).

6. *Abstract*

The abstract lacks a motivation for the study (a short background and a science question to be answered). It is very technical, especially the second sentence, and difficult to follow before reading the manuscript.

Reply (ref.02.06):

- Agreed. The abstract was rewritten by including the short background (page 1, lines 1–3) and the science question (page 1, lines 3–4).

7. *Section 1*

This section needs to be more concise and explain the novelty of the proposed method. In particular, the advantages compared to Soucek and Escoubet (2012) need to be clearly explained. Furthermore, since this study is very similar to the present work, it would be informative if the manuscript contained a comparison between the two methods in cases where the difference between them is the most important, together with an explanation as to why the proposed method is advantageous.

Reply (ref.02.07):

- Agreed. We have cut the introduction text and added paragraph addressing the question (page 2, lines 40–44).

8. *The introduction should be restructured. For example, lines 36-38 “While the radial mapping [...]” say the same thing as lines 52-53 (“While the radial mapping [...]”). The “gap” that this study fills is also stated twice (line 25-26 and line 48-49).*

Reply (ref.02.08):

- Agreed. We shortened the introduction (section 1, pages 1–2), and restructured the manuscript with the review of KF94 model (section 2, pages 2–4) and discussion about different mapping methods (section 3, pages 4–6).
9. *Line 52-53: “While the radial mapping is nearly boundary-fitted on the day-side, the orthogonality of mapping degrades in the flank to tail region.”: this sentence is essential as it mentions the difference between the proposed method and the previous one. However, the term “orthogonality of mapping” should be better explained and perhaps illustrated in a figure. Currently, it is not clear what “orthogonality” refers to.*

Reply (ref.02.09):

- Agreed. The orthogonality (or non-orthogonality) of grids is graphically displayed in figure 1 (page 5) and figure 3 (page 6).
10. *Section 2*

The title should be more informative, for example indicating that the section reviews theory from previous works.

Reply (ref.02.10):

- New section headers are:
 - Sec. 1 – Introduction
 - Sec. 2 – Revisiting the magnetosheath scalar potential
 - * 2.1 Parabolic coordinates
 - * 2.2 Velocity potential
 - * 2.3 Stream function
 - * 2.4 Magnetic scalar potential
 - Sec. 3 – Mapping method comparison
 - * 3.1 Mapping problem
 - * 3.2 Radial direction as reference
 - * 3.3 Magnetopause-normal direction as reference
 - * 4.1 Overview of the procedure
 - * 4.2 Setup
 - * 4.3 Step 1: Measuring the distance to magnetopause
 - * 4.4 Step 2: Computing the thickness of empirical magnetosheath

- * 4.5 Step 3: Computing the magnetosheath thickness in the KF system
 - * 4.6 Step 4: Mapping the position vector onto the KF system
 - * 4.7 Step 5: Evaluating the shell and connector variables
 - * 4.8 Step 6: Computing the potentials and stream function
- Sec. 4 – Magnetopause-normal mapping
 – Sec. 5 – Concluding remark

11. *This section reviews previous results from KF94. It should begin with an introductory sentence explaining what the section contains. It was not entirely clear when the review of previous work ends and the new work starts. Also, the level of detail in the section seems a bit unnecessary. Would the authors perhaps consider referring to the works by Kobel and Flückiger (1994) and Guicking et al. (2012), instead of writing out all the equations in the main text? Alternatively, detailed equations could be placed in an appendix to improve the flow in the text.*

Reply (ref.02.11):

- We added a short paragraph about the section organization in the paper at the end of section 1 (page 1, lines 45–47).
- It is better for the benefit to the readers if the manuscript contains all the necessary information (equations) in a coherent way, rather than simply citing the original references. The reason for this is that the coordinate system used by Kobel and Flückiger (1994) has the co-focal point as the origin. In our method, the bow shock and the magnetopause do not share the same focal point.

12. *Section 3*

Section 3.1: This section contains confusing terminology and probably some typos. These include:

- *Line 130: “magnetosheath-to-magnetopause distance”: the magnetosheath is a region bounded by the magnetopause, so this phrase does not make sense. Instead of magnetosheath, consider writing “position vector r ” (since α_{emp} is the distance from r to the magnetopause).*
- *“magnetosheath-normal direction”: → “magnetopause-normal direction” (line 136 and Figure 4 caption).*
- *“magnetopause thickness” → “magnetosheath thickness” (line 137, Figure 4 caption, line 250, possibly more places in the text).*

Reply (ref.02.12):

- Corrected into “computing the distance to the magnetopause” (page 7, line 140)

- Corrected into “the distance from the planet to the bow shock r_{bs} , the distance from the planet to the magnetopause r_{mp} , the relative distance to the magnetopause α_{emp} ” (page 7, lines 146–147).
- Corrected into “the magnetosheath thickness” (page 7, line 147–148).

13. *Line 138-139: are these the only input parameters, or should the magnetopause and bow shock shapes also be regarded as input?*

Reply (ref.02.13):

- Yes, assuming that the shape of bow shock and that of magnetopause are already known. We changed the sentence into “The position vector \mathbf{r} , the bow shock stand-off distance R_{bs} , the bow shock shape, the magnetopause stand-off distance R_{mp} , and the magnetopause shape are assumed to be known in our mapping” (page 7, lines 148–149).

14. *Also, the reason for defining a unit vector orthogonal to the magnetopause was not clear when reading the manuscript for the first (or second) time. The motivation for this choice should be emphasized in this section and better explained in the introduction.*

Reply (ref.02.14):

- The need for the magnetopause-normal direction is elaborated in section 3 (pages 4–6). The grids need to be as orthogonal as possible particularly around the magnetopause.

15. *Section 3.2-3.5: see above comments about generalization of the method. Equations (18)-(20) and (26)-(28) are specific to the Shue and Farris models; it should be clarified that they do not describe a generalized method.*

Reply (ref.02.15):

- Equations (18)–(20) in the original manuscript are packed into a subsection “Setup” (section 4.2, pages 7–8). An explanation was added at the end of section 4.2 that one needs to compute the radial distance from the planet to the bow shock or magnetopause as a function of the zenith angle when using different shapes (page 8, lines 177–179).
- Steps 1 and 2 in the original manuscript are compressed into section 4.2 in the revised manuscript. The procedure begins with Step 3 in the original manuscript, and is introduced as Step 1 in the revision (page 8, section 4.3).

16. *Section 3.6: The derivations are very similar to what has already been done in section 3.2-3.5. For the sake of getting a better flow in the text, perhaps it would be possible to reduce the number of equations, or make an appendix with the details?*

Reply (ref.02.16):

- We thoroughly checked the logical flow in the text. We moved Eq. (17) in the original manuscript onto Eq. (47) in appendix in the revised manuscript. We corrected Eqs. (13) and (36) in the revised manuscript. We deleted Eqs. (24) and (25) in the original manuscript. All the other equations are necessary and should appear as is.

17. *Section 3.8: Line 270-272: What is meant by the sentence “The mesh pattern [...]” Why is this important?*

Reply (ref.02.17):

- The sentence was deleted. It becomes important when discussing the conformal mapping of magnetosheath coordinates, but we conclude that it is beyond the scope of the manuscript.

18. *Section 4*

This section needs to be reworked when the above comments have been taken into account.

Reply (ref.02.18):

- We went through the conclusion section after revising the main text part. We changed “arbitrary shape” into “non-parabolic shape” (page 15, line 310) and also added more discussion on page 16, lines 327–337.

19. *Line 292: “wider range” – compared to what? (This also appears in the abstract.)*

Reply (ref.02.19):

- Corrected into “wide range” (page 1, line 12 and page 15, line 308).

20. *Line 294: “The method is applicable to an arbitrary shape of magnetosheath domain”: in its current state, the method is specific to the Shue and Farris models, and thus this sentence is too strong (see above comments about generalization of the method).*

Reply (ref.02.20):

- Changed into “non-parabolic shape” (page 15, line 310).

21. *Line 302-304: Here the authors mention non-orthogonality – is the coordinate system (u,v,ϕ) orthogonal in the magnetosheath?*

Reply (ref.02.21):

- No, not orthogonal between u and v any more due to the non-uniqueness or non-existence of conformal mapping in the magnetosheath. But our method restores the orthogonality near the magnetopause.

22. *Line 309-311: “In reality, non-axisymmetric [...]” – this should be expanded on or incorporated in other parts of the manuscript.*

Reply (ref.02.22):

- Done (page 16, lines 327–332).

23. *Technical corrections*

- *Line 4: “solution of Laplace equation” → “solution of the Laplace equation”*
- *Line 51: “magnetopause. But” → “magnetopause, but”*
- *Line 53: “orthogonality of mapping” → “orthogonality of the mapping”*
- *Line 63: “flow velocity is U is” → “flow velocity U is?”*
- *Line 164: “[...], the cylindrical distance” → “[...], and the cylindrical distance”*
- *Line 207: “such that relative distance” → “such that the relative distance”*
- *Line 277: “streamline” → “streamlines”*
- *Line 278: “nose of magnetopause” → “nose of the magnetopause”*
- *Line 293: “solar wind condition” → “solar wind conditions”*

Reply (ref.02.23):

- “solutions” (page 1, line 8).
- “magnetopause. But” was deleted.
- “orthogonality of mapping” was deleted.
- “flow velocity U is” (page 3, line 68).
- “and the cylindrical distance” (page 8, line 177).
- “such that the relative distance” (page 10, line 212–213).
- “streamlines” (page 14, line 291).
- “nose of the magneopause” (page 14, line 292).
- “solar wind conditions” (page 15, line 309).

Scalar-potential mapping of the steady-state magnetosheath model

Yasuhito Narita¹, Simon Toepfer², and Daniel Schmid¹

¹Space Research Institute, Austrian Academy of Sciences, Schmiedlstr. 6, 8042 Graz, Austria

²Institut für Theoretische Physik, Technische Universität Braunschweig, Mendelssohnstr. 3, 38106 Braunschweig, Germany

Correspondence: Y. Narita (yasuhito.narita@oeaw.ac.at)

Abstract. The steady-state magnetosheath model has various applications in studying the plasma and magnetic field profile around the planetary magnetospheres. In particular, the magnetosheath model is analytically obtained by solving the Laplace equation for parabolic boundaries (bow shock and magnetopause). We address the question “How can we utilize the magnetosheath model by transforming into a more general, empirical, non-parabolic magnetosheath geometry?” To achieve the goal, we develop the scalar-potential mapping method which provides a semi-analytic estimate of steady-state flow velocity and magnetic field in the empirical magnetosheath domain. The method makes use of a coordinate transformation from the empirical magnetosheath domain into the parabolic magnetosheath domain, and evaluate a set of the shell variable and the connector variable to utilize the solutions of Laplace equation obtained for the parabolic magnetosheath domain. Our model uses two invariants of transformation: the zenith angle in the magnetosheath and the ratio of the distance to the magnetopause to the thickness of magnetosheath along the magnetopause-normal direction. The use of magnetopause-normal direction makes a marked difference from the earlier model construction using the radial direction as reference. The plasma flow and magnetic field can be determined as a function of the upstream condition (flow velocity or magnetic field) in a wide range of zenith angle. The scalar-potential mapping method is computationally inexpensive by using the analytic expression as much as possible, is applicable to various planetary magnetosheath domains.

1 Introduction

Steady-state plasma flow and magnetic field can be regarded as a realization of potential field in the planetary magnetosheath region when the vorticity and the electric current are treated as ignored. In such a case, the potential is obtained by solving the Laplace equation, which was elegantly and analytically solved by Kobel and Flückiger (1994) for a parabolic shape of magnetosheath (hereafter KF). The KF potential was further extended to the stream function in the magnetosheath by Guicking et al. (2012). The KF solution made a series of breakthroughs in the magnetosheath research. One of the most successful applications is the ability to track the plasma parcel along the streamline in the modeled magnetosheath. The tracking method was extensively used to observationally study the mirror mode growth (e.g., Tatrallyay et al., 2002; Génot et al., 2011) and the streamwise turbulence evolution in the magnetosheath (Guicking et al., 2012). Predictive models of plasma flow and magnetic field serve as a useful tool when combined with the numerical simulation or the observational data.

The KF potential is obtained on the assumption that the planetary bow shock and magnetopause have a parabolic shape sharing the same focal point. Empirical models of the bow shock and magnetopause (fitted to the spacecraft data), on the other

hand, are not necessarily parabolically or co-focally shaped. For example, the empirical Earth bow shock model by Farris et al. (1991) and Cairns et al. (1995) has a parabolic shape but the focal point differs from that of the KF solution; the empirical magnetopause model by Shue et al. (1997) applies a power-law scaling to the parabolic shape such that the magnetic field lines appear stretched in the tail region. The gap between the KF parabolic magnetosheath and the empirical magnetosheath needs to be filled when applying the KF potential in the empirical magnetosheath.

Naively speaking, one wishes to find a conformal mapping (angle-preserving mapping) from the KF parabolic magnetosheath onto a non-parabolic shape of empirical magnetosheath such as the analytic extension of magnetopause shape (Narita et al., 2023). However, no general mathematical algorithm is known so far to obtain the conformal mapping when the spatial domain is not properly bounded. The problem lies in the fact that the magnetosheath is bounded only by two sides, i.e., the standing shock and the magnetopause in the radial direction to the planet, but not bounded along the flow in the tail region. The algorithms of numerical conformal mapping are so far proposed for spatially bounded domains (Papamichael and Whiteman, 1973; Chakravarthy and Anderson, 1979; Fornberg, 1980; Karageorghis et al., 1996) or domains with a closed shape of internal boundaries (Wei et al., 2014).

Here we address the question “How can we utilize the KF magnetosheath model by transforming into a more general, empirical, non-parabolic magnetosheath geometry?” To achieve the goal, we develop a mapping method which provides a semi-analytic estimate of steady-state flow velocity and magnetic field in the empirical magnetosheath domain. Our scalar-potential mapping method is computationally inexpensive by using the analytic expression as much as possible, and is applicable to various planetary magnetosheath domains.

This work is organized in the following fashion. After reviewing the magnetosheath model constructed by Kobel and Flücker (1994) (section 2 and discussing different mapping methods (section 3), the detailed procedure of the magnetopause-normal mapping is presented (section 4) with concluding remark (section 5).

2 Revisiting the magnetosheath scalar potential

2.1 Parabolic coordinates

In the KF parabolic coordinates, the shell variable v and the connector u play an important role in computing the flow velocity and magnetic field in the magnetosheath. These variables are explicitly evaluated using Cartesian coordinates and the radial distance from the focal point as

$$v = \sqrt{r_0 + (x_k - x_0)} \quad (1)$$

$$u = \sqrt{r_0 - (x_k - x_0)}, \quad (2)$$

where r_0 is the distance to the focus at x_0 :

$$r_0 = \sqrt{(x_k - x_0)^2 + y_k^2 + z_k^2}. \quad (3)$$

The focus is along the x axis, and is defined as

$$x_0 = \frac{1}{2}R_{\text{mp}}. \quad (4)$$

x_k , y_k , and z_k are the Cartesian representation of the KF magnetosheath model (i.e., with the pre-fixed bow shock and magne-
60 topause shapes) obtained by projecting the position vector onto the unit vectors e_x , e_y , and e_z :

$$x_k = \mathbf{r}^{(k)} \cdot \mathbf{e}_x \quad (5)$$

$$y_k = \mathbf{r}^{(k)} \cdot \mathbf{e}_y \quad (6)$$

$$z_k = \mathbf{r}^{(k)} \cdot \mathbf{e}_z. \quad (7)$$

To complete the variable set for computing the potentials and the stream function, the azimuthal angle ϕ is introduced as

$$65 \quad \phi = \text{atan}(z_k/y_k). \quad (8)$$

2.2 Velocity potential

In the frame of potential field theory, the **flow velocity** \mathbf{U} is obtained from the velocity potential (scalar potential) $\Phi^{(\text{vel})}$ as

$$\mathbf{U} = -\nabla\Phi^{(\text{vel})} = -\nabla \times (\Psi\mathbf{e}_\phi). \quad (9)$$

70 The symbol \mathbf{e}_ϕ is the unit vector in the azimuthal directions around the symmetry axis (Sun-to-planet direction). Kobel and Flückiger (1994) and Guicking et al. (2012) obtained the analytic expression of the velocity potential $\Phi^{(\text{vel})}$ using the shell variable v (iso-contour lines enveloping the magnetosphere) and the connector variable u (iso-contour lines connecting from the bow shock to the magnetopause).

$$75 \quad \Phi^{(\text{vel})} = -U_x \left(\frac{v_{\text{mp}}^2 v_{\text{bs}}^2}{v_{\text{bs}}^2 - v_{\text{mp}}^2} \right) \left(\frac{u^2 - v^2}{2v_{\text{bs}}^2} + \ln v \right) - \frac{1}{2}U_x (u^2 - v^2) + \Phi_0^{(\text{vel})}, \quad (10)$$

where U_x is the upstream flow velocity, v_{mp} the shell variable at the magnetopause, v_{bs} the shell variable at the bow shock, v the shell variable, u the connector variable, and $\Phi_0^{(\text{vel})}$ a free parameter (integration constant) which is set to zero without loss of generality. The **boundary shell** values v_{mp} and v_{bs} contain the information on the stand-off distances (R_{mp} and R_{bs}) in the subsolar region, and are defined by Kobel and Flückiger (1994) as

$$80 \quad v_{\text{mp}} = \sqrt{R_{\text{mp}}} \quad (11)$$

$$v_{\text{bs}} = \sqrt{2R_{\text{bs}} - R_{\text{mp}}}. \quad (12)$$

2.3 Stream function

Guicking et al. (2012) transformed the KF potential and obtained analytically the stream function Ψ as a function of the shell variable and the connector variable:

$$85 \quad \Psi = -\frac{1}{2}U_x \left(\frac{v_{\text{mp}}^2 v_{\text{bs}}^2}{v_{\text{bs}}^2 - v_{\text{mp}}^2} \right) \frac{u}{v} \left(\frac{v^2}{v_{\text{bs}}^2} - 1 \right) - \frac{1}{2}U_x uv. \quad (13)$$

Hereafter, one may set $U_x = -1$ so that the velocity potential $\Phi^{(\text{vel})}$ is normalized to the upstream velocity. Isocontour lines of the stream function represent the streamline.

2.4 Magnetic scalar potential

The magnetic field in the magnetosheath is derived from the scalar potential in the same fashion as the flow velocity, that is,

$$90 \quad \mathbf{B} = -\nabla\Phi^{(\text{mag})}. \quad (14)$$

The magnetic potential is a function of the shell variable v and the connector u (Kobel and Flückiger, 1994):

$$\begin{aligned} \Phi^{(\text{mag})} = & -\frac{v_{\text{mp}}^2 v_{\text{bs}}^2}{v_{\text{bs}}^2 - v_{\text{mp}}^2} \times \\ & \left[\left(B_y^{(\text{up})} \cos \phi + B_z^{(\text{up})} \sin \phi \right) u \left(\frac{1}{v} + \frac{v}{v_{\text{bs}}^2} \right) + \right. \\ & \left. B_x^{(\text{up})} \left(\frac{u^2 - v^2}{2v_{\text{bs}}^2} + \ln v \right) \right] \\ 95 \quad & -B_x^{(\text{up})}(-x) - B_y^{(\text{up})}y - B_z^{(\text{up})}z + \Phi_0^{(\text{mag})}, \end{aligned} \quad (15)$$

where $B_x^{(\text{up})}$ is the sunward component of the upstream magnetic field (corresponding to the GSE-X in near-Earth space), and $B_y^{(\text{up})}$ and $B_z^{(\text{up})}$ are two components of the upstream magnetic field perpendicular to the x direction. ϕ is the azimuthal angle of the position around the symmetry axis (the y direction is given by the angle $\phi = 0$). The integration constant is chosen as $\Phi_0^{(\text{mag})} = 0$. The magnetic potential cannot be further transformed into the form of stream function since the magnetic field distribution is essentially three-dimensional in the magnetosheath.

3 Mapping method comparison

3.1 Mapping problem

Our task is to find the shell variable v and the connector u in the empirical magnetosheath by finding a suitable mapping of the position vector from the empirical magnetosheath (denoted by \mathbf{r}) onto the KF parabolic system (denoted by $\mathbf{r}^{(k)}$). The evaluated v and u are then readily used to obtain the scalar potentials and the stream function. The flow velocity and the magnetic field in the empirical magnetosheath are obtained by computing the gradient of the respective potential.

110 A practically useful mapping procedure to utilize the KF potential is proposed by Soucek and Escoubet (2012) by using the radial direction as a reference. While the radial mapping can reasonably (i.e., with a relatively high accuracy) transform the KF potential into the empirical magnetosheath domain on the dayside, the mapping quality becomes degraded in the flank region due to the conversion effect associated with the non-orthogonal grid construction. Our approach makes a difference to the radial mapping in using the magnetopause-normal direction as a reference. The azimuthal coordinate ϕ is still orthogonal to the u and v coordinates. We briefly compare between the two mapping methods here.

3.2 Radial direction as reference

115 Soucek and Escoubet (2012) presented in their pioneering work an algorithm of radial mapping by transforming the KF magnetosheath model into a general, empirical magnetosheath shape by referring to the radial direction from the planet and scaling the radial position in the magnetosheath to the KF model. While the radial mapping can reasonably (i.e., with a relatively high accuracy) transform the KF potential into the empirical magnetosheath domain on the dayside, the mapping quality becomes degraded in the flank region due to the strongly non-orthogonal grids. Figure 1 displays a comparison of the radial grids between the KF magnetosheath model and the empirical magnetosheath model. The grids span the radial direction to the planet and transfinite interpolation between the bow shock and the magnetopause. The radial mapping has a drawback in a stronger grid non-orthogonality effect, which causes an artificial converging flow pattern in the flank region (velocity potential shown in Fig. 2 when the scalar potential is directly transformed. In the Soucek-Escoubet method, the problem of flow conversion effect was avoided by solving the MHD Rankine-Hugoniot relation and tracking the streamline iteratively between the KF parabolic magnetosheath model and the empirical magnetosheath model.




Figure 1. Grid pattern generated by the radial mapping for the Kobel-Flückiger parabolic magnetosheath (left panel) and the non-parabolic, empirical magnetosheath (right panel).

125 3.3 Magnetopause-normal direction as reference

Our mapping method makes a difference from the radial mapping method in that the magnetopause-normal direction is used as a reference to the magnetopause. Our method guarantees the grid orthogonality around the magnetopause both on the dayside and in the flank region. The magnetopause-normal grids are shown in Fig. 3 for the KF magnetosheath model (with parabolic




Figure 2. Velocity potential in the Kobel-Flückiger model (left panel) and its radial mapping onto the non-parabolic empirical boundaries (right panel).

130 boundaries) and the empirical magnetosheath model (with non-parabolic boundaries). Even though the exact conformal mapping is not available, the magnetopause-normal mapping method retains the grid orthogonality around the magnetopause. This feature (orthogonality around the magnetopause) plays a crucial role in mapping the scalar potentials. An example of the scalar-potential mapping by referring to the magnetopause-normal direction (our final results) are shown in section 4.8.




Figure 3. Mesh pattern used in the magnetopause-normal mapping in this work for the parabolic boundaries (left panel) and the non-parabolic, empirical boundaries (right panel).

4 Magnetopause-normal mapping

4.1 Overview of the procedure

135 The magnetopause-normal mapping is performed with two transformations. In the first transformation, the position vector is mapped from the empirical magnetosheath \mathbf{r} onto the KF magnetosheath model \mathbf{r}_k . This is achieved on the assumption that the distance to the magnetopause along the magnetopause-normal direction is the same when normalized to the magnetosheath thickness (defined as the distance from the magnetopause to the bow shock along the magnetopause-normal direction). The

azimuthal angle ϕ is the same between the empirical magnetosheath and the KF model. The first transformation is divided into
 140 computing the distance to the magnetopause (step 1), the thickness of the empirical magnetosheath (step 2), the thickness of
 the KF magnetosheath (step 3), and the mapping of the position vector onto the KF model(step 4).

In the second transformation, the mapped position vector is used to compute the shell variable v and the connector variable
 u (step 5) and to obtain the potentials and the stream function in the empirical magnetosheath using Eqs. (10), (13), and (15)
 (step 6). Here again, the azimuthal angle ϕ is treated as the same.

145 Figure 4 illustrates the mapping procedure and graphically explains the variables that need to be determined to perform
 the mapping such as the zenith angle of the nearest magnetopause θ_{mp} , the distance from the planet to the bow shock r_{bs} ,
 the distance from the planet to the magnetopause r_{mp} , the relative distance to the magnetopause α_{emp} , and the magnetosheath
 thickness $\alpha_{emp}^{(bs)}$. The position vector \mathbf{r} , the bow shock stand-off distance R_{bs} , the bow shock shape, the magnetopause stand-off
 distance R_{mp} , and the magnetopause shape are assumed to be known in our mapping.




Figure 4. Variables used in the magnetopause-normal mapping with the zenith angle of the nearest magnetopause θ_{mp} , the radial distance to
 the bow shock and magnetopause along the magnetosheath-normal direction (r_{bs} and r_{mp} , respectively), the distance from the magnetosheath
 to the magnetopause α_{emp} , the magnetopause thickness $\alpha_{emp}^{(bs)}$. The position vector is denoted by \mathbf{r} . The bow shock and magnetopause
 stand-off distances are denoted by R_{bs} and R_{mp} , respectively.

150 4.2 Setup

We begin with a position vector in the empirical magnetosheath domain, and express the position vector as $\mathbf{r} = xe_x + ye_y + ze_z$.
 Hereafter, we present the mapping procedure in the two-dimensional plane spanning the x and y directions for simplicity, but the
 computation in three dimensions is straightforward by representing the y component of position vector in the cylindrical fashion
 as $\rho \cos \phi$ and the z component into $\rho \sin \phi$ using the distance ρ to the x axis. The boundaries (bow shock and magnetopause)
 155 are specified by the users and do not need to be parabolic. In this paper, we use the following bow shock and magnetopause
 models.

- The empirical bow shock position expressed in GSE (Geocentric Solar Ecliptic) coordinates proposed and discussed by Farris et al. (1991); Cairns et al. (1995)

$$x = R_{\text{bs}} - b_{\text{emp}} y^2, \quad (16)$$

160 where R_{bs} is the bow shock stand-off distance and b_{emp} is the empirical flaring parameter. We note here that the original Farris empirical bow shock model is not a paraboloid model, it is an ellipsoid model (with an eccentricity of 0.81), describing the bow shock on the dayside. It is not a proper representation of the far flank bow shock. Also, the Cairns paraboloid bow shock model does not properly represent the far flank bow shock. The distant bow shock shape approaches that of a hyperboloid.

- 165 – The empirical magnetopause position by Shue et al. (1997):

$$x^2 + y^2 - \frac{4R_{\text{mp}}^4}{4R_{\text{mp}}^2 - y^2} = 0, \quad (17)$$

in the Cartesian representation and

$$r_{\text{mp}} = R_{\text{mp}} \sqrt{\frac{2}{1 + \cos \theta}}. \quad (18)$$

in the polar representation.

170 We use a specific exponent for the Shue model (with an alpha exponent of 0.5) in an effort to show that the analytic model is ‘simple’. The solar wind conditions for which this exponent is applicable is not often encountered (e.g., interplanetary magnetic field has the Bz component larger than +8 nT, with specific values of solar wind dynamic pressure).

In our setup, the radial distance from the planet to the bow shock is expressed as (see appendix)

$$175 \quad r_{\text{bs}} = \frac{1}{2b_{\text{emp}} \sin^2 \theta} \left(-\cos \theta + \sqrt{1 - (1 - 4b_{\text{emp}} R_{\text{bs}}) \sin^2 \theta} \right). \quad (19)$$

The radial distance to the magnetopause is given conveniently by Eq. (18). The Shue model reproduces the magnetopause stand-off distance R_{mp} in the subsolar direction ($\theta = 0$), and the cylindrical distance asymptotes to $2R_{\text{mp}}$ in the tail. It is worth noting here that one needs to compute the radial distance from the planet to the bow shock or magnetopause as a function of the zenith angle when using different shapes.

180 4.3 Step 1: Measuring the distance to magnetopause

In the first step, the distance from the given position in the magnetosheath to the nearest magnetopause is computed (see Fig. 5). We express the position vector along the magnetopause-normal direction such as

$$\mathbf{r} = \mathbf{r}_{\text{mp}} + \alpha_{\text{emp}} \mathbf{e}_{\text{mp}}, \quad (20)$$

where r_{mp} is the magnetopause position nearest to the position vector, and e_{mp} is the unit vector in the magnetopause-normal
 185 direction. The unit vector points away from the planet and satisfies the condition

$$r_{\text{mp}} \cdot e_{\text{mp}} > 0 \quad (21)$$

The symbol α_{emp} is the distance to the magnetopause along the magnetopause-normal direction e_{mp} in the empirical magne-
 tosheath.

The nearest magnetopause position is obtained by searching for the zenith angle θ_{mp} for the minimum distance from the
 190 sample position to the magnetopause. The distance D is defined as

$$D = \sqrt{(r_x - r_{\text{mp}} \cos \theta_{\text{mp}})^2 + (r_y - r_{\text{mp}} \sin \theta_{\text{mp}})^2}. \quad (22)$$

The search for the minimum distance is implemented in a brute-force fashion as a function of $\mu_{\text{mp}} = \cos \theta_{\text{mp}}$ in our study.




Figure 5. Measuring the distance to the empirical magnetopause (Step 1).

Having the nearest magnetopause at a distance of r_{mp} and a zenith angle of θ_{mp} , we are ready to compute the magnetopause-
 normal direction and the distance α_{emp} . To obtain the magnetopause-normal direction, we define the magnetopause shape
 195 function f_{mp} as

$$f_{\text{mp}} = x^2 + y^2 - \frac{4R_{\text{mp}}^4}{4R_{\text{mp}}^2 - y^2}, \quad (23)$$

and compute the normal direction by the gradient of f_{mp} as

$$\frac{\partial f_{\text{mp}}}{\partial x} = 2x \quad (24)$$

$$\frac{\partial f_{\text{mp}}}{\partial y} = 2y \left[1 - \frac{4R_{\text{mp}}^4}{(4R_{\text{mp}}^2 - y^2)^2} \right]. \quad (25)$$

200 The magnetopause-normal direction is obtained by normalizing the gradient vector $(\partial_x f_{\text{mp}}, \partial_y f_{\text{mp}})$ and representing with the
 basis vectors (e_x and e_y) as

$$e_{\text{mp}} = \frac{\text{sgn}}{\sqrt{(\partial_x f_{\text{mp}})^2 + (\partial_y f_{\text{mp}})^2}} \times (\partial_x f_{\text{mp}} e_x + \partial_y f_{\text{mp}} e_y) \quad (26)$$

evaluated at the magnetopause ($x = r_{\text{mp}} \cos \theta_{\text{mp}}$ and $y = r_{\text{mp}} \sin \theta_{\text{mp}}$). The magnetopause-normal vector \mathbf{e}_{mp} has a unit length, and the sign ($\text{sgn} = \pm 1$) is chosen such that the normal vector is pointing outward (Eq. 21). The distance α_{emp} to the magnetopause along the normal direction is obtained from Eq. (20) as

$$\alpha_{\text{emp}} = \frac{(x - r_{\text{mp}} \cos \theta_{\text{mp}}) + (y - r_{\text{mp}} \sin \theta_{\text{mp}})}{\mathbf{e}_{\text{mp}} \cdot \mathbf{e}_x + \mathbf{e}_{\text{mp}} \cdot \mathbf{e}_y} \quad (27)$$

Equation (27) is constructed to be robust against the singular behavior on the dayside ($\mathbf{e}_{\text{mp}} \cdot \mathbf{e}_y = 0$) and in distant tail ($\mathbf{e}_{\text{mp}} \cdot \mathbf{e}_x = 0$).

210 4.4 Step 2: Computing the thickness of empirical magnetosheath

In the second step, the magnetosheath thickness is computed using the position vector and the magnetopause normal direction (Fig. 6). For our mapping purpose, the distance α_{emp} is normalized to the magnetosheath thickness $\alpha_{\text{emp}}^{(\text{bs})}$ such that the relative distance $\alpha_{\text{emp}}/\alpha_{\text{emp}}^{(\text{bs})}$ serves as an invariant of the mapping from the empirical magnetosheath onto the KF magnetosheath. To achieve this, we combine Eq. (16) with Eq. (20), and analytically determine the thickness from the bow shock to the magnetopause in the empirical magnetosheath. That is, the thickness $\alpha_{\text{emp}}^{(\text{bs})}$ is obtained by rewriting the bow shock quadratic equation (Eq. 16) for the position vector using the variable $\alpha_{\text{emp}}^{(\text{bs})}$ (Eq. 20) extended to the bow shock location. The equation is again quadratic, and the solution is algebraically obtained as:

$$\alpha_{\text{emp}}^{(\text{bs})} = \frac{1}{2b_{\text{emp}} e_{\text{mp},y}^2} \times \left[-(e_{\text{mp},x} + 2b_{\text{emp}} y_{\text{mp}} e_{\text{mp},y})^2 + d_{\alpha} \right], \quad (28)$$

220 where d_{α} is an auxiliary variable defined as

$$d_{\alpha} = [(e_{\text{mp},x} + 2b_{\text{emp}} y_{\text{mp}} e_{\text{mp},y})^2 - 4b_{\text{emp}} e_{\text{mp},y}^2 \times (x_{\text{mp}} + b_{\text{emp}} y_{\text{mp}}^2 - R_{\text{bs}})]^{1/2}. \quad (29)$$

In the subsolar direction ($y_{\text{mp}} = 0$), the thickness is simplify given as

$$225 \alpha_{\text{emp}}^{(\text{bs})} = R_{\text{bs}} - R_{\text{mp}}. \quad (30)$$

Equations (28) becomes singular in the subsolar direction and Eq. (30) needs to be set separately to avoid the numerical digergence problem.

4.5 Step 3: Computing the magnetosheath thickness in the KF system

In the third step, the magnetosheath thickness is computed in the KF model (Fig. 7). We repeat the procedures of steps 1 and 2 for the KF system and determine the KF magnetosheath thickness as reference. We treat the zenith angle θ_{mp} and the relative distance $\alpha_{\text{emp}}/\alpha_{\text{emp}}^{(\text{bs})}$ as invariants of the mapping between the empirical magnetosheath and the KF system. The KF bow shock




Figure 6. Computing the magnetosheath thickness in the empirical model (Step 2).

location is given as

$$x = R_{\text{bs}} - b_{\text{k}}y^2, \quad (31)$$

where the KF bow-shock flaring parameter b_{k} is pre-fixed as (Kobel and Flückiger, 1994)

$$235 \quad b_{\text{k}} = \frac{1}{4R_{\text{bs}} - 2R_{\text{mp}}}. \quad (32)$$

The radial distance from the planet to the KF bow shock is

$$r_{\text{bs}}^{(\text{k})} = \frac{1}{2b_{\text{k}} \sin^2 \theta} \times \left(-\cos \theta + \sqrt{1 + (4b_{\text{k}}R_{\text{bs}} - 1) \sin^2 \theta} \right). \quad (33)$$

The KF magnetopause is defined in Kobel and Flückiger (1994) as

$$240 \quad x = R_{\text{mp}} - \frac{1}{2R_{\text{mp}}}y^2. \quad (34)$$

From Eq. (34) the radial distance from the planet to the KF magnetopause is computed as

$$r_{\text{mp}}^{(\text{k})} = \frac{R_{\text{mp}}}{\sin^2 \theta} \left(-\cos \theta + \sqrt{1 + \sin^2 \theta} \right). \quad (35)$$

To obtain the magnetopause-normal direction in the KF system, we compute the gradient of the magnetopause shape function:

$$245 \quad f_{\text{mp}}^{(\text{k})} = x - R_{\text{mp}} + \frac{1}{2R_{\text{mp}}}y^2. \quad (36)$$

The gradient is analytically given as

$$\frac{\partial f_{\text{mp}}^{(\text{k})}}{\partial x} = 1 \quad (37)$$

$$\frac{\partial f_{\text{mp}}^{(\text{k})}}{\partial y} = \frac{y}{R_{\text{mp}}} \quad (38)$$

The magnetopause-normal direction $e_{\text{mp}}^{(k)}$ is then obtained by applying Eqs. (37) and (38) to Eq. (26), which reads as

$$250 \quad e_{\text{mp}}^{(k)} = \frac{\text{sgn}}{\sqrt{(\partial_x f_{\text{mp}}^{(k)})^2 + (\partial_y f_{\text{mp}}^{(k)})^2}} \times \left(\partial_x f_{\text{mp}}^{(k)} e_x + \partial_y f_{\text{mp}}^{(k)} e_y \right) \quad (39)$$




Figure 7. Computing the magnetosheath thickness in the KF model (Step 3). The same zenith angle as that in step 2 is used.

The thickness in the KF system $\alpha_k^{(\text{bs})}$ is determined by combining the bow shock shape (Eq. 31) with the position vector at the bow shock:

$$255 \quad \mathbf{r}_{\text{bs}}^{(k)} = \mathbf{r}_{\text{mp}}^{(k)} + \alpha_k^{(\text{bs})} e_{\text{mp}}^{(k)}. \quad (40)$$

Equation (31) becomes again a quadratic equation with respect to the thickness $\alpha_k^{(\text{bs})}$, and the solution reads:

$$\alpha_k^{(\text{bs})} = \frac{1}{2b_k e_{\text{mp},y}^2} \times \left[-(e_{\text{mp},x} + 2b_k y_{\text{mp}} e_{\text{mp},y}) + d_\alpha^{(k)} \right] \quad (41)$$

where the auxiliary variable $d_\alpha^{(k)}$ is defined as

$$260 \quad d_\alpha^{(k)} = \left[(e_{\text{mp},x} + 2b_k y_{\text{mp}} e_{\text{mp},y})^2 - 4b_k e_{\text{mp},y}^2 (x_{\text{mp}} + b_k y_{\text{mp}}^2 - R_{\text{bs}}) \right]^{1/2}. \quad (42)$$

4.6 Step 4: Mapping the position vector onto the KF system

In the fourth step, the mapping of the position vector is performed from the empirical magnetosheath onto the KF system (Fig. 8). Assumption is made such that the relative distance to the magnetopause along the magnetopause-normal direction

265 **is the same between the two systems.** The distance from the magnetosheath position vector to the magnetopause along the magnetopause-normal direction in the KF system α_k is then determined by the relative distance in the empirical magnetosheath α_{emp} , the thickness of the empirical magnetosheath $\alpha_{\text{emp}}^{\text{bs}}$, and magnetosheath thickness in the KF system $\alpha_k^{(\text{bs})}$ as

$$\alpha_k = \alpha_{\text{emp}} \alpha_k^{(\text{bs})} / \alpha_{\text{emp}}^{(\text{bs})}. \quad (43)$$

The mapped position vector is then computed as

$$270 \quad \mathbf{r}^{(k)} = \mathbf{r}_{\text{mp}}^{(k)} + \alpha_k \mathbf{e}_{\text{mp}}^{(k)}, \quad (44)$$

using the nearest magnetopause position $\mathbf{r}_{\text{mp}}^{(k)}$ (Eq. 35), the magnetosheath-to-magnetopause distance α_k (Eq. 43), and the magnetopause-normal direction $\mathbf{e}_{\text{mp}}^{(k)}$ (Eq. 39).




Figure 8. Mapping the position vector onto the KF magnetosheath model (Step 4).

4.7 Step 5: Evaluating the shell and connector variables

In the fifth step, the shell variable v and the connector variable u are computed from the mapped position vector $\mathbf{r}^{(k)}$ using
 275 Eqs. (1) and (2), respectively. The variables v and u are the same as the parabolic coordinates used in the KF potential with a focus at $x_0 = R_{\text{mp}}/2$. In our algorithm, the focus is explicitly given in Eqs. (1), (2), and (3). The azimuthal angle around the symmetry axis ϕ is treated in the same way as in the KF model.




Figure 9. Evaluating the shell variable v and the connector variable u in the KF magnetosheath model (Step 5).

Figure 10 compares the iso-contours of the shell v and the connector u represented in the KF system (left panel) and the empirical magnetosheath (right panel) for a bow shock stand-off distance of $12.8 R_E$ (Génot et al., 2011), a bow shock flaring
 280 of $0.0223 R_E^{-1}$, (Farris et al., 1991; Cairns et al., 1995), and a magnetopause stand-off distance $9.8 R_E$ (Génot et al., 2011).




Figure 10. Iso-contour lines with $u = \text{const.}$ (center of curvature on the left side) and that with $v = \text{const.}$ (center of curvature on the right side) in the KF magnetosheath model (left panel) and the empirical magnetosheath model (right panel). The bow shock stand-off distance is 12.8 Earth radii and the magnetopause stand-off distance is 9.8 Earth radii.

The shell variable v is characterized by the lines with the curvature center on the right side in the panel, and contains the parabolic bow shock (at $v = v_{\text{bs}}$) and magnetopause (at $v = v_{\text{mp}}$) marked by thick lines. The connector variable u has the curvature center on the left side in the panel, and the iso-contour lines are orthogonal to the bow shock and magnetopause. [The computation of the \$u\$ and \$v\$ variables and their gradient and curl is performed in the Cartesian so that the connection represented by the Christoffel symbol vanish in the computation. Computation in the Cartesian domain is also beneficial to the practical application because spacecraft trajectories are often represented in the Cartesian coordinates.](#)

4.8 Step 6: Computing the potentials and stream function

The scalar potentials (velocity potential and magnetic potential) and the stream function are obtained from the shell v and the connector u using Eqs. (10), (13), and (15). The velocity potential (normalized to the upstream flow) is displayed in Fig. 11 left panel, and the stream function in right panel. The iso-contours of the velocity potential represent the lines of the same flow velocity. The iso-contours of the stream function represent the [streamlines](#) in the magnetosheath. The flow is deflected around the [nose of the magnetopause](#) (the subsolar point) and the streamlines are tangential to the magnetopause. [Due to the grid orthogonality around the magnetopause, the streamlines are constructed as tangential to the magnetopause shape, which qualifies the magnetopause-normal mapping method as a useful tool for the magnetosheath model.](#)

The magnetic potential and the derived magnetic field are displayed in Fig. 12. The magnetic potential and the magnetic field (the gradient of the potential multiplied by the minus sign) depend on the upstream field. Fig. 12 shows an example with an upstream field angle of 135 degree to the x axis (i.e., 45 degree to the upstream flow direction). The magnetic field is computed using the central difference scheme. Near the boundaries (bow shock or magnetopause), the mesh resolution is enhanced so that the mesh points do not cross the boundary when computing with the central difference scheme. The upstream field is deflected on the positive y side (right panel, lower half plane), and is draping the magnetopause on the negative y side (right panel, upper half plane).




Figure 11. Velocity potential (left panel) and stream function (right panel) in the empirical magnetosheath domain obtained by mapping onto the shell variable v and the connector variable u .




Figure 12. Magnetic potential for the upstream magnetic field with an angle of 135 degree to the x axis (45 degree to the upstream flow direction), (left panel) and sampled magnetic field vectors obtained by the negative gradient of the magnetic potential (right panel).

5 Concluding remark

Our potential mapping method may be regarded as an updated version of the radial mapping method (Soucek and Escoubet, 2012) by retaining the orthogonality near the magnetopause in the flank to tail region and also by computing the field through the potential mapping. Velocity potential, stream function, and magnetic potential are evaluated in the empirical magnetosheath.

The advantages of our methods are as follows.

1. The method makes extensive use of the exact solution of the Laplace equation (the Kobel-Flückiger potential and the Guicking stream function). The plasma flow and magnetic field can be determined semi-analytically in a [wide range](#) of zenith angle in the magnetosheath when the [solar wind conditions](#) and the boundary shapes are given.
2. The method is applicable to a [non-parabolic shape](#) of magnetosheath domain, opening the door to develop a tool to assist numerical simulations and spacecraft observations of not only the Earth but also the planetary magnetosheath domain.
3. The method is computationally inexpensive. In particular, if the shape of bow shock and magnetopause is analytically given, most of the computational steps in the potential mapping method have an analytic expression.

As stated in section 1, one ideally needs to find a conformal mapping from the KF magnetosheath model onto the empirical magnetosheath. While the conformal mapping is known both for the empirical bow shock and the empirical magnetopause, the conformal mapping of the entire magnetosheath domain still remains a challenge. There are two problems on this. First, the closing boundary (the u -contours) connecting between the bow shock and the magnetopause is not known, and moreover, the uniqueness of finding such a boundary is not guaranteed. Second, the gradients along u are not the same between the empirical bow shock and the empirical magnetopause such that a naive transfinite interpolation ends up with highly non-orthogonal grids in the magnetosheath.

Our method of computing the plasma flow and magnetic field should be compared against the observations and simulations. For example, THEMIS and ARTEMIS spacecraft (Angelopoulos, 2008) and MMS spacecraft (Burch et al., 2016) are providing a huge amount of data on both sides of the bow shock in the equatorial plane; Cluster spacecraft Escoubet et al. (2001) are collecting data in polar orbit; ACE spacecraft data Stone et al. (1998) may be used as an upstream monitor; and Earth flyby data of planetary missions (such as Cassini, BepiColombo) cover the far-distance tail region. In reality, non-axisymmetric structure arises in the magnetosheath. Our method has the possibility to be extended to three-dimensional, non-axisymmetric modeling by the use of magnetopause normal mapping. It is possible to obtain the steady-state magnetosheath potential in a more general sense without referring to the KF94 solution. For example, for a non-axisymmetric geometry of magnetosheath (e.g., Dimmock and Nykyri, 2013), one may numerically solve the Laplace equation for a given set of boundary shapes (bow shock and magnetopause). Various numerical solvers are known for solving the Laplace equation such as the Jacobi method, the Gauss-Seidel method, and the successive over-relaxation (SOR) method. These Laplace solvers are numerically more expensive than the mapping method, but the computation in 3-D is feasible with the contemporary computational resources. Or one may expand the magnetosheath magnetic field in different orthogonal functions. The KF solution makes use of the Bessel functions (Kobel and Flückiger, 1994). For flexible magnetopause and bow shock boundary models, a magnetosheath magnetic field model is constructed by making use of Legendre polynomials (Romashets and Vandas, 2019). There is no restriction regarding the choice of the magnetopause model. The magnetopause-normal direction needs to be computed either analytically using the gradient of the magnetopause function as ∇f_{mp} , or numerically for a user-defined magnetopause shape.

Additional references to analytic models of the magnetosheath magnetic field (using expansions in Legendre polynomials) that make use of flexible magnetopause and bow shock boundary models (e.g., Romashets and Vandas, JGR, [2019])

340 **Appendix: Planet-to-bow shock distance**

By introducing the zenith angle θ and inserting $x = r_{bs} \cos \theta$ and $y = r_{bs} \sin \theta$ in Eq. 16), we obtain the equation for the radial distance to the empirical bow shock:

$$b_{emp} r_{bs}^2 \sin^2 \theta + r_{bs} \cos \theta - R_{bs} = 0. \quad (45)$$

Equation (45) can be algebraically solved, and we take the positive value of the solution as presented in Eq. (19).

345 *Code and data availability.* No codes or data are used in this work.

Author contributions. YN, ST, and DS developed the idea of potential mapping method, checked mathematics, and wrote the manuscript. YN prepared the figures. All authors listed have made a substantial, direct, and intellectual contribution to the work and approved it for publication.

Competing interests. Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial
350 relationships that could be construed as a potential conflict of interest.

References

- Angelopoulos, V. The THEMIS mission, *Space Sci. Rev.*, 141, 5, 2008. <https://doi.org/10.1007/s11214-008-9336-1>
- Burch, J. L., Moore, T. E., Torbert, R. B., and Giles, B. L., Magnetospheric Multiscale overview and science objectives, *Space Sci. Rev.*, 199, 5–21, 2016. <https://doi.org/10.1007/s11214-015-0164-9>
- 355 Cairns, I. H., Fairfield, D. H., Anderson, R. R., Carlton, V. E. H., Paularenas, K. I., and Lazarus, A.: Unusual locations of Earth’s bow shock on September 24–25, 1987: Mach number effects, *J. Geophys. Res.*, 100, 47–62, 1995. <https://doi.org/10.1029/94JA01978>
- Chakravarthy, S., and Anderson, D.: Numerical conformal mapping, *Math. Comp.* 33, 953–969, 1979.
- Dimmock, A. P., and Nykyri, K: The statistical mapping of magnetosheath plasma properties based on THEMIS measurements in the magnetosheath interplanetary medium reference frame, *J. Geophys. Res. Space Physics*, 118, 4963–4976, 2013. <https://doi.org/10.1002/jgra.50465>
- 360
- Escoubet, C. P., Fehringer, M., and Goldstein, M.: Introduction: The Cluster mission, *Ann. Geophys.*, 19, 1197–1200, 2001. <https://doi.org/10.5194/angeo-19-1197-2001>
- Farris, M. H., Petrinec, S. M., and Russell, C. T.: The thickness of the magnetosheath – Constraints on the polytropic index, *Geophys. Res. Lett.*, 18, 1821–1824, 1991. <https://doi.org/10.1029/91GL02090>
- 365 Fornberg, B.: A numerical method for conformal mappings, *SIAM J. Sci. Stat. Comput.* 1, 386–400, 1980. <https://doi.org/10.1137/0901027>
- Génot, V., Broussillou, L., Budnik, E., Hellinger, P., Trávníček, P. M., Lucek, E., and Dandouras, I.: Timing mirror structures observed by Cluster with a magnetosheath flow model, *Ann. Geophys.*, 29, 1849–1860, 2011. <https://doi.org/10.5194/angeo-29-1849-2011>
- Guicking, L., Glassmeier, K.-H., Auster, H.-U., Narita, Y., and Kleindienst, G.: Low-frequency magnetic field fluctuations in Earth’s plasma environment observed by THEMIS, *Ann. Geophys.*, 30, 1271–1283, 2012. <https://doi.org/10.5194/angeo-30-1271-2012>
- 370 Karageorghis, A., Stylianopoulos, N. S., and Zachariades, H. A.: A numerical conformal mapping method for harmonic mixed boundary value problems, *J. Sci. Comp.*, 11, 167–178, 1996. <https://doi.org/10.1007/BF02088814>
- Kobel, E., and Flückiger, E. O.: A model of the steady state magnetic field in the magnetosheath, *J. Geophys. Res.*, 99, 23617–23622, 1994. <https://doi.org/10.1029/94JA01778>
- Narita, Y., Toepfer, S., and Schmid, D.: Magnetopause as conformal mapping, *Ann. Geophys.*, 41, 87–91, 2023. <https://doi.org/10.5194/angeo-41-87-2023>
- 375 Papamichael, N., and Whiteman, J. R.: A numerical conformal transformation method for harmonic mixed boundary value problems in polygonal domains, *J. Applied Math.*, 24, 304–316, 1973. <https://doi.org/10.1007/BF01595198>
- Romashets, E. P., and Vandas, M.: Analytic modeling of magnetic field in the magnetosheath and outer magnetosphere, *J. Geophys. Res. Space Physics*, 124, 2697–2710, 2019. <https://doi.org/10.1029/2018JA026006>
- 380 Schmid, D., Narita, Y., Plaschke, F., Volwerk, M., Nakamura, R., and Baumjohann, W.: Magnetosheath plasma flow model around Mercury, *Ann. Geophys.*, 39, 563–570, 2021. <https://doi.org/10.5194/angeo-39-563-2021>
- Shue, J.-H., Chao, J. K., Fu, H. C., Russell, C. T., Song, P., Khurana, K. K., and Singer, H. J.: A new functional form to study the solar wind control of the magnetopause size and shape, *J. Geophys. Res. Space*, 102, 9497–9511, 1997. <https://doi.org/10.1029/97JA00196>
- Soucek, J., and Escoubet, C. P.: Predictive model of magnetosheath plasma flow and its validation against Cluster and THEMIS data, *Ann. Geophys.*, 30, 973–982, 2012. <https://doi.org/10.5194/angeo-30-973-2012>
- 385 Stone, E. C., Frandsen, A. M., Mewaldt, R. A., Christian, E. R., Margolies, D., Ormes, J. F., and Snow, F. The Advanced Composition Explorer, *Space Sci. Rev.*, 86, 1–22, 1998. <https://doi.org/10.1023/A:1005082526237>

- Tátrallyay, M. and Erdős, G.: The evolution of mirror mode fluctuations in the terrestrial magnetosheath, *Planet. Space Sci.*, 50, 593–599, 2002. [https://doi.org/10.1016/S0032-0633\(02\)00038-7](https://doi.org/10.1016/S0032-0633(02)00038-7)
- 390 Toepfer, S., Narita, Y., and Schmid, D.: Reconstruction of the interplanetary magnetic field from the magnetosheath data: A steady-state approach, *Front. Physics*, 10, 1050859 <https://doi.org/10.3389/fphy.2022.1050859>
- Wei, L. K., Murid, A. H. M., and Hoe, Y. S.: Conformal mapping and periodic cubic spline interpolation, *Mathematika*, 30, 8–20, 2014.