

# Analysis of diurnal, seasonal and annual variations of fair weather atmospheric potential gradient at reduced number concentration of condensation nuclei from long-term measurements at Świder, Poland

Izabela Pawlak<sup>1</sup>, Anna Odzimek<sup>1</sup>, Daniel Kępski<sup>1</sup>, and José Tacza<sup>1</sup>

<sup>1</sup>Institute of Geophysics, Polish Academy of Sciences, Księcia Janusza 64, 01-452 Warsaw, Poland

**Correspondence:** Izabela Pawlak (izap@igf.edu.pl)

**Abstract.** The ground-level atmospheric potential gradient (PG) has been measured with a radioactive collector method in Stanisław Kalinowski Geophysical Observatory in Świder (52.12°N, 21.23°E), Poland, for several decades. Long-term measurements analysed previously revealed rather typical behaviour in the diurnal and seasonal variations of the PG of a land station controlled by pollution. Observation of the potential gradient at such a station usually show a maximum at local winter months which are mostly affected by anthropogenic pollution. The 1965–2005 series has been newly analysed to describe the Świder PG variations in greater detail, also in connection with an analysis of simultaneous measurements of condensation nuclei measured at 6, 12, 18 UT. An attempt is made to calculate the diurnal and seasonal variations at condensation nuclei number concentrations below 10000 cm<sup>-3</sup>. There is a decrease of the PG in the diurnal variation by up to 11% in the winter, and no significant change in the summer. The reduction in the annual variation is 11–26% with the biggest difference in February. In the summer months, this difference is negligible. Despite the efforts to minimise the aerosol effect on the PG, the character of the PG seasonal and annual variation preserves its character with a maximum in the Northern Hemisphere winter and the minimum in the summer. When investigating the effect of reduced CN concentrations on the measured positive conductivity, an increase of 7–17 % is found.

## 1 Introduction

### 1.1 Diurnal and seasonal variation of GEC as deduced from the atmospheric electricity stations observations

The Global Atmospheric Electric Circuit (GEC) manifests itself in the atmospheric potential gradient (PG) observed at any location on the globe (Rycroft et al., 2000). Diurnal and seasonal variations of the PG are the earliest recognised changes of the PG (e.g., Witkowski, 1902; Israël, 1973b). It has been also early established that even in fair weather (FW) conditions the curves of the diurnal variation had different shapes depending on the location of stations, season and local conditions (Israël, 1973a). At land stations a double oscillation was often observed in the PG variation, present in at least one (usually warm) season. For example, at Kew, Uppsala, Tokyo the diurnal variation of the PG exhibits the double oscillations at any time of the year while at Helsinki, Tortosa or Vassijaure it had the double oscillation only in the local summer. At polar stations and over

the oceans a single oscillation is usually observed throughout the year and its character is unitary in the Universal Time which has been firstly recognised by Hoffmann (1924). The average fair weather variation of the PG calculated from 1920–1925 observations over the oceans on board of the Carnegie Institution of Washington research vessel “Carnegie” is known as the Carnegie curve (Parkinson and Torreson, 1931; Israël, 1973a; Harrison, 2013). This curve is considered to reflect the variation of the diurnal activity of the global circuit and is often compared with the diurnal variation of the PG observed in fair weather conditions in other places on the globe. In the new century such analyses have been reported by (e.g., Kubicki et al., 2016; Burns et al., 2017; Nicoll et al., 2019; Tacza et al., 2020; Michnowski et al., 2021, and others). In the seasonal or annual variation many observations indicated a winter maximum in the atmospheric potential gradient compared to the summer. Adlerman and Williams (1996) investigated the connection between the PG local winter maximum and more intensive anthropogenic emission of Aitken nuclei which occurred independent of the station’s location in Northern or Southern Hemispheres. The annual variation of the air conductivity affected by the attachment of ions to aerosol particles which minimises during the local winter as compared to the local summer supported this relationship.

A question arises then when the true maximum of the GEC is (Williams, 2009), and Adlerman and Williams (1996) argued this should be investigated based on observations in locations relatively free from aerosol particles effects. Their analysis of Mauna Loa air-Earth current data and reanalysis of Carnegie and Maud cruises PG data indicated a GEC maximum in the Northern hemisphere summer in agreement with the maximum of global thunderstorm activity (e.g. Christian et al., 2003).

In this paper we reanalyse the diurnal, seasonal and annual variation in the atmospheric potential gradient at the land station Świder (Poland) which also has condensation nuclei monitoring. Previous analyses of the diurnal variation of the PG indicate that Świder is characterised by the double oscillation prevalent in the summer (Northern Hemisphere summer), and the single oscillation in the winter. Furthermore, in the annual variation, the PG values in the winter are higher in amplitude (fluctuation level of 30% of the mean), and the air conductivity in the summer (Kubicki et al., 2007). This could be explained by the higher emissions of dust and aerosol affecting the local electric conductivity in the winter (Kubicki et al., 2003, 2007), with the dust having the largest fluctuation over the year of 60% of the mean, and aerosols only 20% of the their mean. The thermal convection as well as the dependence of dust concentrations on the meteorological conditions is indicated as a significant factor affecting the variations of the PG and conductivity.

Here we particularly analyse these variations at possible low levels of the nuclei number concentration, i.e. less than 10000 particles in cubic centimetre, in order to investigate whether, with such a limit applied, this may have any considerable effect on the seasonal peaks of the PG or PC. We choose concentrations below 10000 particles per  $\text{cm}^{-3}$  because these are characteristic for rural, non-urban and remote regions of the globe (e.g. Landsberg, 1938; Mohnen and Hidy, 2010). For example, Schonland (1953, Table II) cites the results of CN measurements from Heligoland ( $6300 \text{ cm}^{-3}$ ) in the conditions of wind from the direction of land. Junge (1951) presents the results of Landsberg (1938) in a different way, with  $9500 \text{ cm}^{-3}$  being an average CN concentration for an open country, and  $5900 \text{ cm}^{-3}$  a mean minimum concentration for a town – these are the conditions we aim to investigate. It is also worth noting the remark by Podzimek et al. (1982) that experiments showed that up to a value of 10000 particles in  $\text{cm}^3$ , the readings of different particle counters were highly consistent and the chemical composition of the aerosol became less important for their readings. This further facilitates comparison of the results obtained in this study with

those of other studies conducted under low particle concentration conditions, despite different instruments used for the particle count measurements.

## 60 2 Świder published atmospheric electricity data sets

Stanisław Kalinowski Geophysical Observatory of the Institute of Geophysics, Polish Academy of Sciences was founded as the Magnetic Observatory, and since 1929 also worked as an atmospheric electricity station except during the World War II till 1947. **Most of the** pre-war electric measurements were lost, and publication of new results of the resumed electric measurements started in 1958. Measurements reports from years 1957–2005 were subsequently published in the form of  
65 yearbooks such as Warzecha (1960, 1968) or Kubicki (2006). The list of the yearbooks used in this study can be found in Appendix A. Since the beginning of the atmospheric electric measurements, the PG was the main variable to be measured. After 1949 the measurements were complemented by air conductivity recordings and after 2000 by the Maxwell current density measurements. In this paper we aimed at analysing of mainly PG data since it is the most commonly measured parameter of atmospheric electricity worldwide.

### 70 2.1 Site location

Świder is located in the central part of Poland in the Warsaw suburban area, about 25 km south-west of Warsaw (52.12°N, 21.23°E). It used to be a popular holiday and health resort village located on the Świder river. The distance to the nearest urban center, which is the district town of Otwock, is 2.5 km. There is no major industry in the area but there are local anthropogenic sources of air pollution from household heating, very typical for these suburban conditions. The architecture is dominated by  
75 residential buildings and mainly includes single and multifamily houses. The Observatory is located in a less populated area nearer the river. It includes the main office and three observation pavilions, two residential buildings are in a distance. The entire area covers about 7 ha and is partly covered and surrounded by, predominantly, pine trees, with several clearings. In one of these clearings of an area of approximately 1 ha, one pavilion and the station's instruments for atmospheric electricity and meteorology observations are located.

### 80 2.2 PG and conductivity data

The measurements of fair weather atmospheric potential gradient (or the atmospheric electric field strength) at the Świder observatory have been made with radioactive collectors (the activity of about 30  $\mu Ci$ ). Initially two independent collectors were used, each consisting of the radioactive sonde at a height of about 2 m connected to an electrometer and a recording milliammeter. At the beginning Benndorf electrometers were used, in 1964 replaced by an observatory-built electrometer  
85 (Warzecha, 1968, 1976). After 2000 it was replaced by commercially available electrometers. The recording milliammeter was also later replaced by an A/D logger. The PG measurements have absolute calibration and reduction to free plane value (e.g., Kubicki, 2006, and earlier yearbooks). The air conductivity has been measured using a Gerdien tube. Hourly averages for each

month of the PG measured with the collector as well as the positive conductivity (PC) data were tabulated and published in the observatory yearbooks of the period from 1957 to 2005.

## 90 2.3 Condensation nuclei data

Since the continuous atmospheric electricity observations simultaneous systematic measurements of aerosol particle number concentrations of the condensation nuclei type (CN, particle size from  $0.005\ \mu\text{m}$  to  $10\ \mu\text{m}$ ) were made (Warzecha, 1960, 1974). The measurements have been made at a height of 1 m, only in 1998–1999 at 3 m height. Observations were performed three times a day: first between 6:10 GMT and 6:30 GMT (5:50 – 6:20 GMT till 1971), next between 11:00 GMT and 11:30 GMT, 95 and the last between 18:10 UT and 18:30 UT (19:00 – 19:30 GMT till 1971). These three times of observations are referred to in the yearbooks as 6, 12, and 18 GMT or UT, respectively. The measurements of the concentration of condensation nuclei at the three terms have been included in the observatory yearbooks.

The measurements between 1965 and 2005 in Świder were carried out using two counters: a small Scholz counter, and from 1982 a photoelectric CN counter built in the observatory using a chamber of a Verzář counter as a base (the Verzář counter was 100 previously used for occasional continuous observations). The two counter models are described in more detail by Grabovskiy (1956). The measurement method used in the counters was the process of condensation of water vapor on atmospheric aerosol particles present in the measurement chamber, followed by a quantitative analysis of the resulting mist droplets. A Scholz counter is a type of condensation counter constructed by Scholz as an improvement of the Aitken counter, designed to measure the concentration of condensation and nuclei nearly the total concentration of aerosol. The main part of the small Scholz 105 counter was a brass cylindrical chamber with a volume of  $102\ \text{cm}^3$  and a height of 4 cm; the adiabatic volume expansion ratio was 1.25. Measurements were performed within the clearing of the meteorological station using the suction method, with an air sample of  $1\ \text{cm}^3$  volume. According to Miller and Bodhaine (1982), with such volume expansion ratio, the supersaturation is in the range 2.93–3.40 for initial temperatures between  $25^\circ\text{C}$  and  $10^\circ\text{C}$ , respectively, which is equivalent to the supersaturation between 3.93–4.40, according to Wilson (1897). Measurements with the Scholz counter were repeated five times, and one 110 measurement takes several minutes. The Scholz counter allows measurements in a wide range of CN concentrations from 5 to 960 000 particles in  $\text{cm}^3$ , and its experimental error should not be higher than the experimental error of the Aitken counter which is about 10% (Grabovskiy, 1956; McMurry, 2000).

Since January 1983, the measurements have been performed with the photoelectric CN counter that was placed inside a measurement pavilion. The basic measurement of the number of CN took place in a cylindrical chamber filled with the tested 115 air sample of the volume equal to  $680\ \text{cm}^3$ . Estimates of the number of droplets were obtained using a photoelectric counter system by measuring the extinction of light. The air samples were collected from the outside of the building at a height of 1 m above the ground. The suction of air was made through a 1 m long rubber pipe using an electric rotational pump. The measuring range of the counter was 4500 to 850 000 CN in  $1\ \text{cm}^3$ . The electronic circuit system was built (also patented) by Stanisław Warzecha. The measurement accuracy was 15% Warzecha and Warzechowa (1979). The yearbooks do not provide 120 any details on the cross-check of measurements during the instrument transition period.

## 2.4 Criteria of fair weather conditions

In order to investigate the global signal of atmospheric electricity and minimise the effects of local weather, criteria for fair weather conditions are used (e.g. Imyanitov and Chubarina, 1967). Since 1965 a WMO standard for the fair weather conditions has been applied at Świder: low cloudiness up to 3/8 in okta scale, no precipitation or fog, wind speed less than 6 m/s and  
125 PG amplitude less than 1000 V/m (e.g., Warzecha, 1968; Kubicki, 2006). The criteria for fair weather were assessed by the observatory staff on an ongoing basis. The fair weather hours are marked in the yearbooks' PG and conductivity data tables.

## 3 Overview of fair weather PG and CN data

In our analysis we consider all fair weather hourly values of the potential gradient (PG) and the condensation nuclei concentrations (CN) from 6, 12, 18 UT published in the Observatory yearbooks 1965–2005 (no criterion regarding the minimum number  
130 of points in a month has been applied). There is a total of 131634 hourly PG values with a subset of 16072 hourly values with corresponding CN concentrations and 3303 missing data (2.5%). Table B1 and Table B2 in the Appendix B include the number of FW hourly average mean values of PG and number of FW days (24 h) for each month and year of the analysed period. There are more FW values during the local warmer seasons from March to September. The percentage of FW hours is about 37% on average (see also Table 1) and varies between 49% in the summer and 23% during winter. The total number of FW  
135 days is 1414. The monthly percentage of FW days is about 9% on average and rises to 14% during summer and decreases to 4% during winter.

The time variations of monthly mean values of PG and CN during this period is shown in Fig. 1. There seems to be several periods of monotonic increase and decrease of the PG, and a general trend of increase of the PG by 2.5 V/m per year. The general trend for CN concentration is a decrease by  $-120 \pm 20 \text{ cm}^{-3}$  per year. Analysis of the long-term variation in these series  
140 will be a subject of another study.

Histogram of these values and main statistical parameters of the set are given in Fig. 2. The mean average, Ma, of the total fair weather PG of 1965–2005 is 230 V/m, and the standard deviation, SD, is 131 V/m. The median, Me, is 207 V/m, the first quartile, Q1, equals 139 V/m, and the third quartile, Q3, 294 V/m. The kurtosis, K, is equal to 5.4 and the skewness, S, is 1.3 what characterises a leptokurtic and right-skewed distribution. These parameters describe the total FW PG set but in next  
145 sections we concentrate on the subset of PG hours for which simultaneously measured CN data also exist, and particularly on the subset for limited CN concentrations up to  $10000 \text{ cm}^{-3}$ . The statistical parameters of the distribution of these subsets are also given in Fig. 2 separated by slashes. The selection of PG data measured at the same time as CN creates a new distribution with slightly higher mean (248 V/m) or median value (224 V/m) but similar kurtosis and skewness of  $\sim 5$  and  $\sim 1$ , respectively. When we further limit the CN concentration the mean and median value are lower, 221 V/m and 208 V/m, respectively.  
150 Selection of PG-CN pairs limits the number of values, N, to 16072 (12%) and with the condition of CN lower than  $10000 \text{ cm}^{-3}$  the number of pairs decreases to 3593 (3%) as indicated in the Table 1.

In Fig. 3 we show the histogram of the measured CN values. The mean value of the total fair weather CN concentration is  $18980 \text{ cm}^{-3}$  and the standard deviation is  $13240 \text{ cm}^{-3}$ . The median is  $15510 \text{ cm}^{-3}$ , the first quartile is  $10340 \text{ cm}^{-3}$  and the

**Table 1.** Number of total hours for the period January 1965 – December 2005 together with number of all fair weather PG hours and number of fair weather PG hours corresponding to the CN concentration values below 10000 cm<sup>-3</sup>, 8000 cm<sup>-3</sup>, 6000 cm<sup>-3</sup> and 4000 cm<sup>-3</sup>, respectively. The percentages refer to the number in bold in the same colour in the first raw of the table.

All hours 1965-2005	All FW hours 1965-2005	FW PG hours 1965-2005	FW PG all CN	FW PG CN<10000	FW PG CN<8000	FW PG CN<6000	FW PG CN<4000
<b>359400</b>	<b>134937</b>	<b>131634</b>	<b>16072</b>	<b>3593</b>	<b>2025</b>	<b>915</b>	<b>122</b>
	37.54%	97.5%	4.47%	1.00%	0.56%	0.25%	0.03%
			11.91%	2.66%	1.50%	0.69%	0.09%
			12.21%	2.72%	1.54%	0.71%	0.09%
				22.35%	12.60%	5.69%	0.76%

**Table 2.** Statistical measures of the distribution of all values FW PG and CN in the series 1965–2005.

	Spring		Summer		Autumn		Winter		all year	
	PG	CN	PG	CN	PG	CN	PG	CN	PG	CN
	(V/m)	(cm <sup>-3</sup> )	(V/m)	(cm <sup>-3</sup> )	(V/m)	(cm <sup>-3</sup> )	(V/m)	(cm <sup>-3</sup> )	(V/m)	(cm <sup>-3</sup> )
Ma	250	21180	193	15330	253	20020	351	20930	249	18970
Sd	129	14940	84	11220	130	12970	165	12800	134	13270
Me	230	16900	184	12310	238	16740	336	18200	224	15260
Q <sub>1</sub>	163	11330	131	8700	163	11330	236	11815	157	10200
Q <sub>3</sub>	317	26000	241	18200	329	24190	448	25600	317	22650
S	1.2	1.9	0.8	2.9	1.0	1.9	0.6	1.7	1.2	2.1
K	5.7	7.0	4.8	15.2	4.7	7.9	3.2	7.4	5.3	8.8

third quartile is 22650 cm<sup>-3</sup>. The kurtosis is equal to 8.8 and the skewness 2.1 what indicates a leptokurtic and right-skewed distribution. The number of fair weather CN values is 16756 and the subset of CN below 10000 cm<sup>-3</sup> (see green frame in Fig. 3) is 3736 which corresponds to 22% of the total CN values. Table 2 and Fig. 4 show the main statistical measures of the FW PG and CN distributions discussed above and separately for each season. Fig. 4 presents the seasonal distribution of CN concentrations. The green line separates the distributions of CN concentrations below 10000 cm<sup>-3</sup> which will be used in next sections.

The lowest concentrations of CN in fair weather conditions (there is almost twice more counts of the concentrations in the summer compared to winter) are observed in the summer with the mean value of ~15000 cm<sup>-3</sup>. In the other seasons the mean value is close to 20000 cm<sup>-3</sup> with the highest standard deviation values in the spring ~15000 cm<sup>-3</sup> and the lowest ~11000 cm<sup>-3</sup> in the summer and ~13000 cm<sup>-3</sup> in the winter and in the autumn. In terms of kurtosis and skewness the values in the

spring, autumn and winter are similar and vary between 7.0-7.9 and 1.7-1.9, respectively. In the summer these parameters are  
165 higher reaching values 15.2 and 2.9.

#### 4 Diurnal and seasonal variation of the potential gradient in Świder, 1965-2005

In this section we analyse the diurnal and seasonal variation of Świder FW PG and how it relates to different levels of CN concentrations. Previous works on the analysis of diurnal and seasonal variations of PG focused mainly on full 24 hours of FW, i.e. an FW day needed to accomplish FW conditions for each hour between 0 and 23 UT (Kubicki et al., 2016). This  
170 restriction limited the amount of available data. Michnowski et al. (2021) used FW PG values with at least 12 hours of fair weather conditions. In this work we increase the amount of data taking in account every FW hourly value to calculate the diurnal and seasonal variations.

##### 4.1 Diurnal variation of PG

Fig. 5 presents the PG diurnal variation by season (winter – December, January, February, spring – March, April, May, summer  
175 – June, July, August, autumn – September, October, November) and the all year mean variation. The shape of the PG curve is substantially different depending on the season. During the winter months there is a single broad maximum lasting from the noon (~13 UT) until late evening (~20 UT) whereas during the spring-summer-autumn period double maxima (~7 and ~19 UT) are present. The amplitude variation of the PG curve during the winter is larger than in the summer (Table 2). Świder is thus characterised by the type I of the “double oscillation” variation in the diurnal average curve as according to Israël (1973a),  
180 i.e. the double oscillation is prevalent except in local wintertime and its maxima occur at different hours depending on the season. Differences between the PG diurnal curves at Świder and the classical Carnegie diurnal curve come from the much greater variability and dynamics of the planetary boundary layer (PBL) characteristic for the location and at Świder affected by pollution, meteorological factors and seasonal effects (Kubicki et al., 2016; Nicoll et al., 2019; Michnowski et al., 2021, Section 4).

185 Since we cannot calculate the whole diurnal variation of CN we show results only for 6, 12, 18 UT (represented by symbols) in Fig. 6. On the background (solid lines) are shown the Świder 1968/69 diurnal curves from Warzecha (1976) obtained by the recordings of the condensation nuclei concentrations using a Verzář counter every 15 minutes. We note that in this comparison we calculated the mean values only for 24-hour FW days since Warzecha presented the diurnal curves for cloudless days. The mean CN values for the seasons are given in the Table 2. The highest mean CN concentrations at these hours are registered at  
190 12 UT although looking at diurnal curves of Warzecha (1976) this is already after the morning maximum of CN concentrations between 6 and 12 UT (all seasons) and before the afternoon/evening maximum. These values are especially high in the winter and the spring. The concentration values at 6 and 18 UT are similar, however the lowest values are noted in the morning. At 6 and 18 UT the mean CN concentrations in the spring and autumn are almost equal while at 12 UT the spring mean is higher, even than the winter value which generally is the highest among seasons. Compared with the diurnal curves of Warzecha  
195 (1976) the CN concentrations fairly agree with the diurnal variation except for the 18 UT (excluding summer) and for 12 UT in

summer and autumn. Some differences are likely since we compare long-term mean values with one year means. Nevertheless, it is worth noting that the diurnal CN curves fall between the CN averages at one standard deviation.

## 4.2 Seasonal and annual variation of PG

Analogically to Fig. 4 in Fig. 7 we show the seasonal differences in the distribution of FW PG at different CN concentrations. Their main statistical parameters for seasonal distributions are given in the figures as well as in Table 2. Fig. 7 presents the seasonal histograms of the hourly averaged (at 6, 12, 18 UT) FW PG values for 1965–2005. Each histogram contains two types of distributions: red colour indicates the FW PG values measured simultaneously with CN concentrations, and green colour indicates a subset of the FW PG values for CN concentrations less than  $10000 \text{ cm}^{-3}$ . The main statistical measures for both of these distributions are displayed within each plot. Depending on the season we observe different PG distribution characteristics. For spring, summer and autumn most values ranged between 100–300 V/m with mean value of  $\sim 250 \text{ V/m}$  in the spring and autumn and  $\sim 200 \text{ V/m}$  in the summer. Winter values ranged between 200–400 V/m and the mean was  $\sim 350 \text{ V/m}$ . The standard deviation is also the highest in the winter. The kurtosis values for the spring and autumn are 4.7–4.8 while the summer kurtosis is the highest at 5.7 and the winter is the lowest at 3.2. The skewness of the winter distribution is the lowest of 0.6, and the skewness of all the other distributions is between 0.8–1.2 indicating more values concentrated around the mean value. We have used the non-parametric test of Kolmogorov-Smirnov (KS) and Mann-Whitney U (MW) test to estimate the statistical significance of the difference between the PG distributions for decreasing CN concentrations. In regard to the distributions shown in Fig. 2 and in Fig. 7, both KS and MW tests indicated statistically significant differences (at the significance level of 0.05) between the PG-CN all and PG-CN<10000 distributions for the whole year as well as in the spring, autumn and winter. In case of the variations shown in Fig. 11, the KS test indicated statistically significant differences between the PG populations for all CNs and CN<10000, in January and February only. The MW test indicated such in April and November as well. When comparing PG and CN all with PG and CN<8000 the results of both tests indicate the statistically significant difference also in December, similarly to PG and CN<6000.

Another way to look at the relationship between aerosol or condensation nuclei concentration is presented in the Fig. 8. Intensity of colour indicates the number (counts) of data points at certain range of PG and CN. Here the PG values are plotted against CN concentrations by season. In the summer there seems to be little influence of CN concentration on PG as the similar range of PG values is observed at wide range of CN concentrations. Some kind of dependency is visible in the winter, and spring and autumn are intermediate in this sense with some high PG values at high CN concentrations.

## 5 Diurnal, seasonal and annual variation at different level of CN concentration

**When the continuity of the conduction current is fulfilled**, the atmospheric potential gradient depends on local conductivity of the air. The process of attaching of highly electrically ions to the aerosols causes reduction their mobility and, consequently, a decrease of local conductivity and an increase of the PG values (e.g., Israël, 1973a; Bennett and Harrison, 2008; Matthews et al., 2019; Wright et al., 2020). In this section we analyse if limiting CN concentrations to the lowest ranges in the CN



distribution affects the PG values. We consider CN concentrations below  $10000 \text{ cm}^{-3}$  (green histograms in Fig. 4) and decrease the concentration limit by  $2000 \text{ cm}^{-3}$  ( $10000 \text{ cm}^{-3}$ ,  $8000 \text{ cm}^{-3}$ ,  $6000 \text{ cm}^{-3}$ ) to investigate the diurnal, seasonal and annual variations of PG at the lowest CN concentrations. In earlier sections we showed that limiting the CN concentration to  $10000 \text{ cm}^{-3}$  did not change significantly the PG distributions, here we investigate if any possible extreme limits affect this behaviour. Right part of Table 1 presents the counts and percentages of the whole PG population of values within each CN concentration limit. In Fig. 9, in the upper panel we show the histograms of CN concentrations and in next panels are shown the histograms of PG values at the three selected limits of CN concentration. There are 3736 values below  $10000 \text{ cm}^{-3}$ , 2086 values below  $8000 \text{ cm}^{-3}$  and 939 values below  $6000 \text{ cm}^{-3}$ , which is about 23%, 13%, 6% of all CN concentrations, respectively. Below  $4000 \text{ cm}^{-3}$  there was only 123 values (0.8%) and therefore this subset was not taken into account. These values represent the whole span of the studied time period, as shown in the small panel in the top panel of Fig. 9. They also represent all seasons as shown in Table 6, even though there is proportionally more counts in the warmer season because of better FW conditions. In the lower panels of Fig. 9 the histograms of corresponding PG values are shown. We note a slightly decreasing tendency in the mean value of the population especially at CN concentrations below  $6000 \text{ cm}^{-3}$  approximately 210 V/m and lower kurtosis and skewness compared with the distributions below  $10000 \text{ cm}^{-3}$  and  $8000 \text{ cm}^{-3}$ .

Seasonal means are shown in Table 3 and Fig. 10 for each hour term (6, 12, 18 UT) and the mean average over the terms, together with the standard error of the mean. Here we also notice a decrease in the mean values of PG for each term at lower CN concentrations particularly at  $6000 \text{ cm}^{-3}$  limit (i.e., comparing PG values between CN-all and CN below  $6000 \text{ cm}^{-3}$ ). This is present in the winter ( $73 \pm 15 \text{ V/m}$  or  $\sim 13\%$  for total mean value), spring ( $30 \pm 8 \text{ V/m}$  or  $\sim 12\%$ ) and autumn ( $29 \pm 9 \text{ V/m}$  or  $\sim 12\%$ ) and the decrease are larger for 6 UT (e.g.,  $30 \pm 12 \text{ V/m}$  or 12% in the spring) and 18 UT (e.g.,  $57 \pm 29 \text{ V/m}$  or 15% in the winter,  $36 \pm 23 \text{ V/m}$  or 14% in the spring). However, in the summer there is practically no effect of CN and no clear tendency of decrease or increase (the largest decrease of several V/m only).

Finally, in Fig. 11 and Table 4 the annual variation of FW PG is presented, calculated as monthly averages. Additionally in Table B3 in the Appendix B the number of FW PG values calculated as monthly values for all and limited CN concentration is presented. PG variations calculated at 6, 12, 18 UT are plotted separately from the average curve. The maximum of the PG is in February which is also the maximum at 12, 18 UT. The minimum of PG is in June which is also the month of the minimum at 6, 12 and 18 UT. The lowest PG amplitude of the differences between the maximum and the minimum are in the 6 UT curve:  $91 \pm 12 \text{ V/m}$  (wide maximum Dec-Mar, minimum in June) and much higher PG differences are at 12 and 18 UT:  $213 \pm 12 \text{ V/m}$  (Feb-Jun) and  $217 \pm 13 \text{ V/m}$  (Feb-Jun/Jul), respectively,  $174 \pm 7 \text{ V/m}$  (Feb-Jun) in the average curve.

In the PG variations represented by the curves calculated with limitations of CN concentration there is a significant reduction of PG from October to March (11%–26%) and the biggest differences are seen in February at each term 6, 12, 18 UT and on average. In February the PG decreases by  $95 \pm 25 \text{ V/m}$  or 26%. In January and November the change is about 20% ( $75 \pm 24 \text{ V/m}$  and  $58 \pm 20 \text{ V/m}$ , respectively) and in December about 13% ( $45 \pm 28 \text{ V/m}$ ). From April to September there is a lack of or little influence of low CN concentration, and the PG differences are mostly of several V/m, which is of the order of the standard deviation of the mean error, except for April and September where these differences can be larger than 10 V/m. In September at 18 UT we note a larger decrease of PG of  $44 \pm 29 \text{ V/m}$  (18%). In addition, as we already have seen in the seasonal

**Table 3.** Mean values of fair weather PG by season for all and limited CN concentration values together with the one standard error of the mean corresponding to 6, 12, 18 UT and on the average.

Winter					Spring			
Hour	6 UT	12 UT	18 UT	Mean	6 UT	12 UT	18 UT	Mean
CN all	291 ± 6	370 ± 5	373 ± 6	350 ± 3	251 ± 3	227 ± 3	264 ± 3	250 ± 2
CN <10000	265 ± 9	339 ± 12	320 ± 10	305 ± 6	228 ± 5	217 ± 6	238 ± 7	228 ± 4
CN <8000	258 ± 12	333 ± 16	318 ± 13	299 ± 8	231 ± 7	219 ± 8	228 ± 11	227 ± 5
CN <6000	225 ± 15	342 ± 19	316 ± 23	277 ± 12	221 ± 9	215 ± 9	228 ± 20	220 ± 6
Autumn					Summer			
Hour	6 UT	12 UT	18 UT	Mean	6 UT	12 UT	18 UT	Mean
CN all	227 ± 4	253 ± 3	272 ± 4	253 ± 2	215 ± 2	169 ± 2	187 ± 2	193 ± 1
CN <10000	210 ± 8	238 ± 6	261 ± 10	235 ± 4	207 ± 3	168 ± 3	188 ± 3	192 ± 2
CN <8000	213 ± 10	231 ± 7	256 ± 13	231 ± 5	208 ± 4	169 ± 4	194 ± 4	193 ± 2
CN <6000	208 ± 12	227 ± 9	242 ± 19	224 ± 7	202 ± 7	174 ± 5	196 ± 7	190 ± 3

plots in Fig. 10 at 12 UT there is even an inverse relationship between the PG and CN. For example in July at 12, 18 UT and in August at 18 UT, although these differences are within the statistical error. In terms of the change of the amplitude of maximum–minimum differences at 6 UT there is a small change with decreasing CN concentration reaching  $77 \pm 41$  V/m (Dec–Jun) at  $6000 \text{ cm}^{-3}$  CN limit. At 12 UT it is  $165 \pm 35$  V/m (Jan–Jul) at 6000 CN limit while at 10000 and 8000 CN limit it is practically the same as with no limits ( $\sim 213 \pm 12$  V/m, Jan/Feb–Jun–Jul). At 18 UT at all CN limits the amplitude is  $\sim 160$  V/m compared with  $217 \pm 13$  V/m with no limits. There seems to be slight shift between the maximum or minimum month which is January or February for the maximum, and June and July for the minimum. In general, there are significantly lower differences between winter and summer in the annual variation of the FW PG at the lowest CN concentration limit considered, however, the maximum still stays in the winter (Dec–Jan–Feb).

## 6 PG and conductivity model with variable aerosol content

In this section we model a hypothetical change in the PG when we limit the number concentration of CN. The relationship between atmospheric electric potential gradient and aerosol concentration is complex. These two most often measured variables are related through the electrical conductivity of the air. The conductivity,  $\sigma$ , is influenced by the loss of small atmospheric ions due to attachment to aerosol particles, and decreases when the attachment intensifies. When we assume that the mobility and concentration of positive and negative ions are equal, and the ions are singly charged, then, in a steady state the ion conductivity is determined by Eq. 1 (e.g. Schonland, 1953; Makino and Ogawa, 1985; Sapkota and Varshneya, 1990).

$$\sigma = 2e\mu n = \frac{4e\mu q}{\beta N + \sqrt{(\beta N)^2 + 4\alpha q}} \quad (1)$$

280 where  $e = 1.6 \times 10^{-19}$  C is the elementary electric charge,  $\mu$  is the ion mobility, and  $n$  is the ion concentration,  $q$  - the ion production,  $\alpha$  is the ion recombination rate coefficient, and  $\beta N$  represents the ion loss due to attachment to particles and droplets of concentration  $N$ , and  $\beta$  the attachment rate coefficient. In the atmosphere the loss of ions in fair weather is mainly due to the attachment to aerosol particles. The process of attachment is dependent both on the number concentration and the spectrum of the size of aerosol particles, and is usually expressed by the ion-aerosol loss term, in general being a  
 285 sum of contributions over different aerosol types and sizes (e.g., Israël, 1973a; Tinsley and Zhou, 2006). The term can be replaced by the effective coefficient  $\beta$  and the total aerosol concentration  $N$ , i.e.  $\beta N$  as in Eq. 1. We set the other parameters in Eq. 1 as follows:  $q = 2.2 \text{ cm}^3 \text{ s}^{-1}$ ,  $\alpha = 1.3 \times 10^{-6} \text{ cm}^{-3}$ ,  $\mu = 1.5 \text{ cm}^2 \text{ V s}^{-1}$ , similarly to other conductivity models based on other empirical models and measurements (Makino and Ogawa, 1985; Sapkota and Varshneya, 1990; Tinsley and Zhou, 2006; Kulkarni, 2022). The ion production by cosmic rays of the order of  $1.0 \text{ s}^{-1} \text{ cm}^{-3}$  is appropriate for the production at  
 290 the ground level, and of the order of  $1.0 \text{ s}^{-1} \text{ cm}^{-3}$  for the production by radioactivity of radon. We set the total ion production value at Świder to  $2.2 \text{ s}^{-1} \text{ cm}^{-3}$ . The mobility of  $1.5 \pm 0.3 \text{ cm}^2 \text{ s}^{-1} \text{ V}^{-1}$  up to 15 km could be used for both positive and negative ions according to (Swider, 1988).

In case where the conduction current,  $j_C = \sigma \text{ PG}$ , is preserved, a decrease or increase in the electrical conductivity should result in proportional increase or, respectively, decrease in the potential gradient:

$$295 \quad \frac{\sigma_2}{\sigma_1} = \frac{PG_1}{PG_2} = s \quad (2)$$

where indices 1 and 2 refer to value before and after the change, respectively.

Rural continental aerosols of natural origin occur in Świder but there may be a lot of pollution from suburban traffic and  
 300 domestic heating (Majewski and Przewoźniczuk, 2009; Pietruczuk and Jarosławski, 2013). More recent observations to the considered CN concentration measurements at Świder, of aerosol size distribution, confirm the occurrence of polluted aerosol which consists of large concentrations of fine particles dominated by the nuclei mode of the mean diameter 20–30 nm, and by the accumulated mode of the mean diameter 100–120 nm (Kubicki et al., 2016). We want to focus on the winter conditions when the decrease of the potential gradient at lower CN concentrations, expected to be due to decreasing effect of pollution on  
 305 the air conductivity, is the biggest as obtained in Sec. 4, and next, to estimate the effect during the summer. We will emulate the conditions by calculating the conductivity at different aerosol composition and concentrations. Let us consider a mix of continental aerosol types the Świder aerosol may be composed of, and calculate the corresponding conductivities as well as the ratio of the conductivities as in Eq. 2.

According to Hess et al. (1998) a land aerosol may consist of 3-4 main aerosol types: continental clean, continental average, continental polluted, and urban. These aerosol types consist of three basic aerosol components: water insoluble (*WatInsol*), water soluble (*WatSol*) and soot (*Soot*), the two latest contributing almost 99% of the total concentration and contained in the range of CN particle size. Their individual  $\beta$  coefficients are:  $\beta_{WatInsol} = 2.80 \times 10^{-5} \text{ cm}^3/\text{s}$ ,  $\beta_{WatSol} = 1.18 \times 10^{-6} \text{ cm}^3/\text{s}$ , and  $\beta_{Soot} = 5.56 \times 10^{-7} \text{ cm}^3/\text{s}$ , respectively, obtained from calculations based on the equations used in Tinsley and Zhou (2006). In this model the effect of relative humidity on the conductivity is not included.

$$\beta N = \beta_{WatSol} N_{WatSol} + \beta_{WatInsol} N_{WatInsol} + \beta_{Soot} N_{Soot} \quad (3)$$

where  $N_{WatInsol}$ ,  $N_{WatSol}$ , and  $N_{Soot}$  are the number concentrations of the three aerosol components.

At first we set the limiting values of concentration  $N$  for 100% relative composition of soot versus 100% water soluble aerosol at  $26000 \text{ cm}^{-3}$  ( $N_{Soot}$ ) and  $6000 \text{ cm}^{-3}$  ( $N_{WatSol}$ ), respectively. The first value corresponds to the highest average aerosol concentrations in the winter (Fig. 4, Table 2), and the second value refers to the lowest concentrations measured at Świder ( $\sim 4000$  occur too, but they are very rare). A small, constant contribution of insoluble aerosol of  $500 \text{ cm}^{-3}$  is also assumed. In Tab. 5 we present the results of the model calculations when the relative proportions of water soluble and soot components vary (columns 1-2), but the total concentration is dominated by the soot component, except at the lowest ratios of  $N_{Soot}/N_{WatSol}$ , which result in water soluble aerosol at a low number concentration. Columns 3-5 give the resulting number concentrations of  $N_{Soot}$ ,  $N_{WatSol}$ , and the total concentration CN. In the column 6 we give the calculated ion-aerosol loss term  $\beta N$ , and in the column 7 the effective ion attachment rate  $\beta_{eff} = (\beta_{eff} N)/N$  which here varies from 1.07 to  $3.24 \times 10^{-6} \text{ cm}^3 \text{ s}^{-1}$ . Sapkota and Varshneya (1990) mention that Hoppel predicts a range of  $0.8$  to  $3.0 \times 10^{-6}$  of  $\beta_{eff}$  for the continental aerosol. In the last column we give the calculated conductivity. Twice the polar (positive) conductivity calculated from newly digitised 1965-2005 data:  $4.4 \pm 0.2 \times 10^{-15} \text{ S/m}$  for the winter, and  $8.0 \pm 0.2 \times 10^{-15} \text{ S/m}$  for the summer. The value given equals twice the seasonal average of the positive polar conductivity calculated from the hourly values reported in the observatory yearbooks (in fair weather conditions).

An average winter situation may correspond to the concentrations of  $1200 \text{ cm}^{-3}$  of water soluble aerosol and  $18200 \text{ cm}^{-3}$  of soot which give the total CN concentration of  $20500 \text{ cm}^{-3}$ , since the average observed mean is  $\sim 20900 \text{ cm}^{-3}$ . For the observed median,  $\sim 18200 \text{ cm}^{-3}$ ,  $1800 \text{ cm}^{-3}$  of water soluble aerosol and  $15600 \text{ cm}^{-3}$  of soot could be more appropriate. The conductivity in these cases,  $4.56$  and  $4.69 \times 10^{-15} \text{ S/m}$ , are close to  $4.4 \times 10^{-15} \text{ S}$ , an average observational winter value. This has been adjusted by assuming  $q = 2.2 \text{ cm}^3 \text{ s}^{-1}$ . When we look for concentration characteristic for the summer it would rather be reflected by  $4200 \text{ cm}^{-3}$  of water soluble aerosol and  $7800 \text{ cm}^{-3}$  of soot for the summer mean average mean  $\sim 15300 \text{ cm}^{-3}$ , and the median  $\sim 12300 \text{ cm}^{-3}$ . The total CN concentration is  $14500$ – $12500 \text{ cm}^{-3}$ , since the observational (Fig. 4, Tab. 2). However, the conductivity for such concentration is  $4.97 \times 10^{-15} \text{ S/m}$  and remains much too low compared with the observations, since the summer average is  $\sim 8.0 \times 10^{-15} \text{ S/m}$ . Such conductivity level requires the ion-aerosol loss term  $\beta N$  be about  $1.5 \times 10^{-2} \text{ s}^{-1}$ , and the effective  $\beta_{eff} = 1.25 \times 10^{-6}$  at  $N = 12000 \text{ cm}^{-3}$  or  $\beta_{eff} = 1.0 \times 10^{-6}$  at  $N =$

15000  $\text{cm}^{-3}$ . Such values of  $\beta$  are more of the order of  $\beta_{WatSol}$ , therefore we now consider a mix of the three selected aerosol components dominated by the water soluble component. We set the limiting values of concentration  $N$  for 100% relative composition of soot versus 100% water soluble aerosol at 6000  $\text{cm}^{-3}$  ( $N_{Soot}$ ) and 28000  $\text{cm}^{-3}$  ( $N_{WatSol}$ ),  
 345 respectively. Now the lowest concentrations are only due to soot, and small additions of water soluble concentrations. A small, constant contribution of insoluble aerosol of 200  $\text{cm}^{-3}$  is assumed in this case. The results of the conductivity calculations are shown in Tab. 6.

An average summer situation may correspond to the concentration ratio of 40% vs 60% of the water soluble and soot components which give the total CN concentration of 15000  $\text{cm}^{-3}$ . The conductivity in this case is  $5.73 \times 10^{-15}$  S/m. When  
 350 we look for concentrations characteristic for the winter they would rather be reflected by the ratio of 70% vs 30%, i.e. CN total 21600  $\text{cm}^{-3}$ , and the model conductivity is close to the observational value of  $\sim 4.4 \times 10^{-15}$  S/m. It seems this conductivity model could describe both winter and summer conditions, however situations from the model of Tab. 5 cannot be excluded.

There may be other atmospheric particles that cause a larger difference between the winter and summer conductivities, like the dust particles, even though these are in general much less numerous than CN. The conductivity value very much  
 355 depends on both  $\beta_{eff}N$  and  $q$ , with  $\beta_{eff}N$  depending on the distribution of the sizes of the aerosol particles. In particular, the insoluble component also plays an important part through the high attachment rate. These may also vary between the summer and the winter. More analysis, and observational data from Świder are needed to develop a more realistic conductivity model, particularly of the aerosol size distributions.

## 7 Effect of low CN concentrations on the air positive conductivity

360 In this section we analyse the annual variation of the positive conductivity (PC or  $\lambda_+$ ) and the effect of decreased CN concentrations, similarly as in Sec. 5. The total number of PC hourly values is lower than the total number of hourly PG: 124216 hourly values (about 95% of the PG set), and 15189 values have the corresponding hourly value of CN concentration at 6, 12 or 18 UT. These are also data sets of considerable size for the calculation of the annual variation which we present in Fig. 12. The all average mean of the conductivity is  $3.28 \pm 0.01 \times 10^{-15}$  S/m, as indicated in the first row of Tab. 7. There we give also  
 365 seasonal values with their standard errors. The winter positive conductivity  $2.24 \pm 0.01 \times 10^{-15}$  S/m is 1.8 times the highest seasonal, summer conductivity of  $3.98 \pm 0.01 \times 10^{-15}$  S/m. For each season we observe an increase in the mean values of PC at lower CN concentrations. The biggest differences were noted when comparing PC values for all CN and CN below 6000  $\text{cm}^{-3}$  (summer, winter and the whole year), or CN below 8000  $\text{cm}^{-3}$  (in the spring and autumn). An increase of about 17% was noted for the spring ( $0.58 \times 10^{-15}$  S/m), the autumn ( $0.52 \times 10^{-15}$  S/m) and for the winter ( $0.36 \times 10^{-15}$  S/m). The  
 370 lowest increase of about 8% was observed in the summer ( $0.33 \times 10^{-15}$  S/m).

## 8 Discussion

At many land atmospheric electricity stations, including Świder, the increase of the potential gradient during local wintertime or rush hours, which is usually related to increase of emission from intensive domestic heating and transport, is considered to be due to a decrease in the electrical conductivity (Harrison, 2006; Kubicki et al., 2016; Tacza et al., 2021). Having at disposal all digitised PG hourly values we calculated the diurnal variation of PG from all available fair weather hours. As in the previous results of such analysis at Świder (e.g., Warzecha, 1991; Kubicki et al., 2007), made on the basis of 24-hour fair weather days 1965-2000, or fair weather hours during days with 12 hours of FW conditions over the time period 2004–2011 (Michnowski et al., 2021) the PG diurnal variation shown in Fig. 5 confirmed the different behaviour according to season. The double oscillation is evident during spring, summer and autumn (in the summer with peaks at 7 and 20 UT), and in winter a single oscillation (a minimum at ~3UT and a peak at 19 UT). Similar PG diurnal curves were found in Reading for summer and winter (Nicoll et al., 2019). One hypothesis for the two PG peaks found in summer at urban sites could be due to pollution of vehicular traffic during rush hours. Majewski and Przewoźniczuk (2009) performed Particulate Matter ( $10\mu\text{m}$ ) measurements for eleven stations in Warsaw. They found clear two peaks (7 and 20 UT) in warm weather (April to September) and less pronounced peaks in cold weather (October to March). As mentioned earlier, Świder is located in a suburban site so vehicular transportation is not expected affect too much in the PG diurnal variation. Then, the hypothesis for the difference in the PG diurnal variation in summer and winter is a combination of “sunrise effect”, generally thought to be related to mixing of the near-surface electrode layer (which is an accumulation of positive charge next to the negatively charged Earth’s surface). The first peak (at ~7 UT) could be associated both with the dynamic changes in the planetary boundary layer and generation of the secondary aerosol (Kubicki et al., 2007). Then, as the temperature is continuously increasing, the convection intensify producing mixing processes in the PBL causing transport of aerosol to higher altitudes and therefore, the PG decreases. As the temperature decrease after 16 UT the PG return to normal values. In winter, the reduced variability in PBL height (due to diminished convection) therefore leads to more quiescent meteorological conditions which results in a more stable diurnal variation in PG, and the disappearance of the morning maximum peak (see also Nicoll et al., 2019).

The annual cycle of emissions with maxima in the winter and minima in the summer affecting the land PG measurements (Adlerman and Williams, 1996; Nicoll et al., 2019; Shatalina et al., 2019) are considerable at Świder, as confirmed in the analysis in Sec. 3 (Figs. 5, 6, 8). Similar behavior in the PG seasonal variation was found in Kew and Reading, in the UK, and Nagycenk, in Hungary (Märzcz et al., 1997; Harrison and Aplin, 2002; Nicoll et al., 2019). Harrison and Aplin (2002) associated the PG higher values during winter, at Kew, to the influence of smoke pollution. Märzcz et al. (1997) suggested that the PG seasonal variation at Nagycenk is likely associated to the condensation nuclei variation. Nicoll et al. (2019) suggested that the PG higher values during winter at Reading is due to more use of domestic heating, and it is very likely that at Świder the PG values are higher during winter due to domestic heating producing soot. This is also in agreement with the simple conductivity model in this work. Measurements of  $\text{PM}_{2.5}$  in Krakow (south of Poland) in 2020/2021 found higher  $\text{PM}_{2.5}$  concentrations during winter month compared with summer months (Ryś and Samek, 2022). The time intervals of this study

and the PG measurement at Świder are different (in fact both site locations are very far away), however, a similar behavior in  
405 PM is very likely at Świder.

The aerosol concentration changes also annually due to variation in convection processes in the planetary boundary layer (PBL), and the PBL height, more influential during warm season and especially in the local summertime. At Świder, as discussed in Sec. 3, the summer CN concentrations differ much more from the rest of the year (Fig. 4) being minimal through all the year. As opposed to winter the change in the summer concentrations does not seem to be so significant on average  
410 (Fig. 9). The effect of reduced aerosol concentration to its lowest levels did not change the average PG amplitude in the summer, while in the other seasons this effect was observed particularly in the winter (Sec. 4).

According to the conductivity model for varying ground-level CN concentrations characteristic for Świder, described in Sec. 5, there could be increases in the ion conductivity, and decreases of the PG, but the relative change differs from the observed conditions.

415 The changes of PG due to changes in the conductivity obscure the real changes of the PG resulting from the activity of the GEC, and the annual maximum of PG occurs usually during local winter. In this work we wanted to investigate in what way taking into account fair weather PG values at low CN concentrations affects the PG annual maximum. In general, our results show that even though the PG decreases at low CN concentrations in the winter and in the autumn, the maximum of PG remains in the winter, and when investigating similar effects on the conductivity they are not so much dependent on season  
420 (Tab. 7, Fig. 12).

## 9 Summary

The findings of this work could be summarised as follows:

- We calculated diurnal, seasonal and annual variations of fair weather potential gradient based on 1965–2005 digitized time series of data from Świder observatory yearbooks, and we provided details of the statistics for the PG and the  
425 condensation nuclei CN number concentrations, measured during this period at three specific hours of measurement (6, 12 and 18 UT). The general average mean of PG is 230 V/m, and the average PG at the CN concentration observation terms is 248 V/m while the CN average mean is about  $19000 \text{ cm}^{-3}$ . The PG and CN distributions are both leptokurtic (i.e., narrow) and right-skewed. A subset of PG values for CN concentration below  $10000 \text{ cm}^{-3}$  (which correspond to 20% of the whole CN-PG pair data) was selected for further analysis of the effect of aerosol concentration on the PG  
430 variation. In particular, we aimed to investigate the PG annual variation in connection with the GEC variability. The PG average mean for reduced levels of CN concentrations is 221 V/m, and the CN average mean in this subset is  $7290 \text{ cm}^{-3}$ .
- Summer and winter months have a clear signature in the PG diurnal variation for all CN concentrations, likely associated with mixing processes in the planetary boundary layer. For the subset with CN below  $10000 \text{ cm}^{-3}$ , we found that PG  
435 values do not show any significant variation in the summer. On the other hand, for the winter we found a PG difference of 11%. Furthermore, this PG difference is about 20% when we consider CN below  $6000 \text{ cm}^{-3}$ .

- Local winter months have higher PG values compared with local summer months, a common feature of continental atmospheric electricity stations. This PG seasonal variation is maintained even at lower CN concentrations (e.g., 10000, 8000, 6000 cm<sup>-3</sup>). We found that there is a reduction in the PG values by 11-26% with the biggest difference in February. In the summer months this difference is negligible. From this, we can conclude that aerosol at these concentrations has more effect in the PG during winter compared with summer.
- On the opposite, local summer months have higher positive conductivity values compared with local winter months, for Świder there is factor of 1.8 between the corresponding mean average values  $3.98 \pm 0.01 \times 10^{-15}$  S/m and  $2.24 \pm 0.01 \times 10^{-15}$  S/m, respectively. At lower CN concentrations (e.g., 10000, 8000, 6000 cm<sup>-3</sup>) we found an increase in the conductivity by 7–17% over the winter, spring and autumn and summer. This is different from the effect on the PG, and also could indicate a non-Ohm PG component or the component which is not very sensitive to CN concentration levels or both. Future calculations of the annual variation of the PG in the hope of finding an annual variation of the GEC have to take these effects into account.
- A simplified modelling of electrical conductivity affecting the PG is created. The ion loss in this model is due to attachment mainly to water soluble aerosol particles and soot particles, the main components of continental aerosol.

*Data availability.* Data is available on request from co-author A. Odzimek (aodzimek@igf.edu.pl).

## Appendix A

This Appendix contains the list of Świder Observatory yearbooks used in this study.

- Warzecha S. (Ed.), 1968: Rocznik Elektryczności Atmosferycznej i Meteorologii 1965, Prace Obs. Geof. Świder 38.
- 455 Warzecha S. (Ed.), 1968: Électricité atmosphérique et météorologie Observatoire Géophysique de St. Kalinowski à Świder 1966, Mater. Prace Zakł. Geofizyki 23.
- Warzecha S. (Ed.), 1969: Électricité atmosphérique et météorologie Observatoire Géophysique de St. Kalinowski à Świder 1967, Mater. Prace Zakł. Geofizyki 28.
- Warzecha S. (Ed.), 1970: Électricité atmosphérique et météorologie Observatoire Géophysique de St. Kalinowski à Świder 460 1968, Mater. Prace Zakł. Geofizyki 38.
- Warzecha S. (Ed.), 1971: Électricité atmosphérique et météorologie Observatoire Géophysique de St. Kalinowski à Świder 1969, Mater. Prace Zakł. Geofizyki 44.
- Warzecha S. (Ed.), 1972: Électricité atmosphérique et météorologie Observatoire Géophysique de St. Kalinowski à Świder 1970, Mater. Prace Zakł. Geofizyki 53.
- 465 Warzecha S. (Ed.), 1973: Électricité atmosphérique et météorologie Observatoire Géophysique de St. Kalinowski à Świder 1971, Mater. Prace Zakł. Geofizyki 63.



- Warzecha S. (Ed.), 1974: Électricité atmosphérique et météorologie Observatoire Géophysique de St. Kalinowski à Świder 1972, Mater. Prace Zakł. Geofizyki 77.
- Warzecha S. (Ed.), 1974: Électricité atmosphérique et météorologie Observatoire Géophysique de St. Kalinowski à Świder 470 1973, Mater. Prace Zakł. Geofizyki 80.
- Warzecha S. (Ed.), 1975: Électricité atmosphérique et météorologie Observatoire Géophysique de St. Kalinowski à Świder 1974, Mater. Prace Zakł. Geofizyki 92.
- Warzecha S. (Ed.), 1976: Électricité atmosphérique et météorologie Observatoire Géophysique de S. Kalinowski à Świder 1975, Publs. Inst. Geophys. Pol. Acad. Sci. 104 (D-2).
- 475 Warzecha S. (Ed.), 1978: Électricité atmosphérique et météorologie Observatoire Géophysique de S. Kalinowski à Świder 1976, Publs. Inst. Geophys. Pol. Acad. Sci. 121 (D-6).
- Warzecha S. (Ed.), 1979: Électricité atmosphérique et météorologie Observatoire Géophysique de S. Kalinowski à Świder 1977, Publs. Inst. Geophys. Pol. Acad. Sci. 131 (D-8).
- Warzecha S. (Ed.), 1980: Électricité atmosphérique et météorologie Observatoire Géophysique de S. Kalinowski à Świder 480 1978, Publs. Inst. Geophys. Pol. Acad. Sci. 140 (D-10).
- Warzecha S. (Ed.), 1981: Électricité atmosphérique et météorologie Observatoire Géophysique de S. Kalinowski à Świder 1979, Publs. Inst. Geophys. Pol. Acad. Sci. 148 (D-12).
- Warzecha S. (Ed.), 1982: Électricité atmosphérique et météorologie Observatoire Géophysique de S. Kalinowski à Świder 1980, Publs. Inst. Geophys. Pol. Acad. Sci. 151 (D-14).
- 485 Warzecha S. (Ed.), 1982: Électricité atmosphérique et météorologie Observatoire Géophysique de S. Kalinowski à Świder 1981, Publs. Inst. Geophys. Pol. Acad. Sci. 158 (D-16).
- Warzecha S. (Ed.), 1983: Électricité atmosphérique et météorologie Observatoire Géophysique de S. Kalinowski à Świder 1982, Publs. Inst. Geophys. Pol. Acad. Sci. 168 (D-17).
- Warzecha S. (Ed.), 1984: Électricité atmosphérique et météorologie Observatoire Géophysique de S. Kalinowski à Świder 490 1983, Publs. Inst. Geophys. Pol. Acad. Sci. 177 (D-19).
- Warzecha S. (Ed.), 1985: Électricité atmosphérique et météorologie Observatoire Géophysique de S. Kalinowski à Świder 1984, Publs. Inst. Geophys. Pol. Acad. Sci. 190 (D-23).
- Warzecha S. (Ed.), 1986: Électricité atmosphérique et météorologie Observatoire Géophysique de S. Kalinowski à Świder 1984, Publs. Inst. Geophys. Pol. Acad. Sci. 194 (D-24).
- 495 Warzecha S. (Ed.), 1987: Électricité atmosphérique et météorologie Observatoire Géophysique de S. Kalinowski à Świder 1985, Publs. Inst. Geophys. Pol. Acad. Sci. 209 (D-27).
- Warzecha S. (Ed.), 1988: Électricité atmosphérique et météorologie Observatoire Géophysique de S. Kalinowski à Świder 1984, Publs. Inst. Geophys. Pol. Acad. Sci. 219 (D-29).
- Warzecha S. (Ed.), 1989: Électricité atmosphérique et météorologie Observatoire Géophysique de S. Kalinowski à Świder 500 1988, Publs. Inst. Geophys. Pol. Acad. Sci. 229 (D-31).
- Warzecha S. (Ed.), 1990: Électricité atmosphérique et météorologie Observatoire Géophysique de S. Kalinowski à Świder

- 1989, *Publs. Inst. Geophys. Pol. Acad. Sci.* 234 (D-34).
- Warzecha S. (Ed.), 1991: *Électricité atmosphérique et météorologie Observatoire Géophysique de S. Kalinowski à Świder* 1990, *Publs. Inst. Geophys. Pol. Acad. Sci.* 247 (D-37).
- 505 Warzecha S. (Ed.), 1992: *Électricité atmosphérique et météorologie Observatoire Géophysique de S. Kalinowski à Świder* 1991, *Publs. Inst. Geophys. Pol. Acad. Sci.* 253(D-39).
- Warzecha S. (Ed.), 1993: *Électricité atmosphérique et météorologie Observatoire Géophysique de S. Kalinowski à Świder* 1992, *Publs. Inst. Geophys. Pol. Acad. Sci.* 264 (D-41).
- Warzecha S. (Ed.), 1994: *Électricité atmosphérique et météorologie Observatoire Géophysique de S. Kalinowski à Świder* 1993, *Publs. Inst. Geophys. Pol. Acad. Sci.* 271 (D-43).
- 510 Warzecha S. (Ed.), 1995: *Électricité atmosphérique et météorologie Observatoire Géophysique de S. Kalinowski à Świder* 1994, *Publs. Inst. Geophys. Pol. Acad. Sci.* 280 (D-44).
- Warzecha S. (Ed.), 1996: *Électricité atmosphérique et météorologie Observatoire Géophysique de S. Kalinowski à Świder* 1995, *Publs. Inst. Geophys. Pol. Acad. Sci.* 290 (D-47).
- 515 Kubicki M. (Ed.), 1997: *Results of atmospheric electricity and meteorological observations. S.Kalinowski Geophysical Observatory at Świder - 1996*, *Publs. Inst. Geophys. Pol. Acad. Sci.* 299 (D-49).
- Kubicki M. (Ed.), 1998: *Results of atmospheric electricity and meteorological observations. S.Kalinowski Geophysical Observatory at Świder - 1997*, *Publs. Inst. Geophys. Pol. Acad. Sci.* 307 (D-51).
- Kubicki M. (Ed.), 1999: *Results of atmospheric electricity and meteorological observations. S.Kalinowski Geophysical Observatory at Świder - 1998*, *Publs. Inst. Geophys. Pol. Acad. Sci.* 321 (D-52).
- 520 Kubicki M. (Ed.), 2000: *Results of atmospheric electricity and meteorological observations. S.Kalinowski Geophysical Observatory at Świder - 1999*, *Publs. Inst. Geophys. Pol. Acad. Sci.* 324 (D-54).
- Kubicki M. (Ed.), 2001: *Results of atmospheric electricity and meteorological observations. S.Kalinowski Geophysical Observatory at Świder - 2000*, *Publs. Inst. Geophys. Pol. Acad. Sci.* 333 (D-55).
- 525 Kubicki M. (Ed.), 2002: *Results of atmospheric electricity and meteorological observations. S.Kalinowski Geophysical Observatory at Świder - 2001*, *Publs. Inst. Geophys. Pol. Acad. Sci.* 342 (D-58).
- Kubicki M. (Ed.), 2003: *Results of atmospheric electricity and meteorological observations. S.Kalinowski Geophysical Observatory at Świder - 2002*, *Publs. Inst. Geophys. Pol. Acad. Sci.* 355 (D-61).
- Kubicki M. (Ed.), 2004: *Results of atmospheric electricity and meteorological observations. S.Kalinowski Geophysical Observatory at Świder - 2003*, *Publs. Inst. Geophys. Pol. Acad. Sci.* 372 (D-65).
- 530 Kubicki M. (Ed.), 2005: *Results of atmospheric electricity and meteorological observations. S.Kalinowski Geophysical Observatory at Świder - 2004*, *Publs. Inst. Geophys. Pol. Acad. Sci.* 383 (D-68).
- Kubicki M. (Ed.), 2006: *Results of atmospheric electricity and meteorological observations. S.Kalinowski Geophysical Observatory at Świder - 2005*, *Publs. Inst. Geophys. Pol. Acad. Sci.* 391 (D-71).

## Appendix B

This Appendix contains the list of additional Tables: Table B1, Table B2 and Table B3.

*Author contributions.* AO digitised PG and CN data from Świder yearbooks and conceived the subject of the study. IZ and AO performed the investigation and wrote the paper. DK and JT revised and edited the paper. All authors discussed the results.

540 *Competing interests.* The authors declare that they have no competing interests.

*Acknowledgements.* The work is supported by Poland National Science Centre grant No 2021/41/B/ST10/04448 at the Institute of Geophysics, Polish Academy of Sciences.

## References

- Adlerman, E. J. and Williams, E. R.: Seasonal variation of the global electrical circuit, *J. Geophys. Res.-Atmos.*, 101, 29 679–29 688, <https://doi.org/10.1029/96JD01547>, 1996.
- Bennett, A. J. and Harrison, R. G.: Variability in surface atmospheric electric field measurements, *Journal of Physics: Conference Series*, 142, 012 046, <https://doi.org/10.1088/1742-6596/142/1/012046>, 2008.
- Burns, G. B., Frank-Kamenetsky, A. V., Tinsley, B. A., French, W. J. R., Grigioni, P., Camporeale, G., and Bering, E. A.: Atmospheric Global Circuit Variations from Vostok and Concordia Electric Field Measurements, *J. Atmos. Sci.*, 74, 783–800, <https://doi.org/10.1175/JAS-D-16-0159.1>, 2017.
- Christian, H. J., Blakeslee, R. J., Boccippio, D. J., Boeck, W. L., Buechler, D. E., Driscoll, K. T., Goodman, S. J., Hall, J. M., Koshak, W. J., Mach, D. M., and Steward, D. F.: Global frequency and distribution of lightning as observed from space by the Optical Transient Detector, *J. Geophys. Res.-Atmos.*, 108, ACL 4–1–ACL 4–15, <https://doi.org/10.1029/2002JD002347>, 2003.
- Grabovskiy, Y. T.: *Atmosfernye jadra kondensacii*, Gidrometeoizdat, Leningrad, 1956.
- Harrison, R.: Urban smoke concentrations at Kew, London, 1898–2004, *Atmospheric Environment*, 40, 3327–3332, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2006.01.042>, 2006.
- Harrison, R. and Aplin, K.: Mid-nineteenth century smoke concentrations near London, *Atmos. Env.*, 36, 4037–4043, [https://doi.org/https://doi.org/10.1016/S1352-2310\(02\)00334-5](https://doi.org/https://doi.org/10.1016/S1352-2310(02)00334-5), 2002.
- Harrison, R. G.: The Carnegie Curve, *Surv. Geophys.*, 34, 209–232, <https://doi.org/10.1007/s10712-012-9210-2>, 2013.
- Hess, M., Koepke, P., and Schult, I.: Optical Properties of Aerosols and Clouds: The Software Package OPAC, *Bulletin of the American Meteorological Society*, 79, 831 – 844, [https://doi.org/10.1175/1520-0477\(1998\)079<0831:OPOAAC>2.0.CO;2](https://doi.org/10.1175/1520-0477(1998)079<0831:OPOAAC>2.0.CO;2), 1998.
- Hoffmann, K.: Bericht uber die in Ebeltothafen auf Spitsbergen in der Jahren 1913/14 durchgefurten luftelektrischen Messungen (11 36' 15" E, 79 09' 14" N), *Breitr. Phys. Frei. Atmos.*, 11, 1–19, Report about the Atmospheric Electric Measurements Performed in 1913/14 in Ebeltothafen in Spitsbereg (11 36' 15" E, 79 09' 14" N), 1924.
- Imyanitov, I. M. and Chubarina, Y. V.: Electricity of free atmosphere, Gidrometeoizdat, Leningrad, 1965, technical Translation from Russian, NASA, Washington, TT F-425, 1967.
- Israël, H.: *Atmospheric Electricity*, vol. I, Fundamentals, Conductivity, Ions, Israel Program for Scientific Translations, Jerusalem, 1973a.
- Israël, H.: *Atmospheric Electricity*, vol. II, Fields, Charges, Currents, Israel Program for Scientific Translations, Jerusalem, 1973b.
- Junge, C.: Nuclei of Atmospheric Condensation, in: *Compendium of Meteorology*, edited by Byers, H. R., pp. 182–191, American Meteorological Society, 1951.
- Kubicki, M.: Results of Atmospheric Electricity and Meteorological Observations at S. Kalinowski Geophysical Observatory at Świder, 2005, *Publ. Inst. Geophys. Pol. Acad. Sci.*, D-71, 3–27, 2006.
- Kubicki, M., Michnowski, S., Myslek-Laurikainen, B., and Warzecha, S.: Long term variations of some atmospheric electricity, aerosol, and extraterrestrial parameters at Swider Observatory, Poland, in: *Proceedings of the 12th International Conference on Atmospheric Electricity*, 9-13 June 2003, Versailles, France, pp. 291–294, 2003.
- Kubicki, M., Michnowski, S., and Myslek-Laurikainen, B.: Seasonal and daily variations of atmospheric electricity parameters registered at the Geophysical Observatory at Świder (Poland) during 1965-2000, in: *Proceedings of the 13th International Conference on Atmospheric Electricity*, 13-17 August 2007, Beijing, China, vol. I, p. 4 pp., 2007.

- Kubicki, M., Odzimek, A., and Neska, M.: Relationship of ground-level aerosol concentration and atmospheric electric field at three observation sites in the Arctic, Antarctic and Europe, *Atmos. Res.*, 178–179, 329–346, <https://doi.org/10.1016/j.atmosres.2016.03.029>, 2016.
- 580 Kulkarni, N. M.: Non-conventional approach to computation of the atmospheric global electric parameters: Resultant data and analysis, *J. Earth Syst. Sci.*, 131, <https://doi.org/10.1007/s12040-022-01981-3>, 2022.
- Landsberg, H.: Atmospheric condensation nuclei, *Ergeb. Kosm. Phys.*, 48, 410, 1938.
- Majewski, G. and Przewoźniczuk, W.: Study of Particulate Matter Pollution in Warsaw Area, *Pol. J. Environ Studies*, 18, 293–300, <https://www.pjoes.com/Study-of-Particulate-Matter-Pollution-r-nin-Warsaw-Area,88234,0,2.html>, 2009.
- 585 Makino, M. and Ogawa, T.: Quantitative estimation of global circuit, *J. Geophys. Res.*, 90, 5961–5966, <https://doi.org/10.1029/JD090iD04p05961>, 1985.
- Matthews, J., Wright, M., Clarke, D., Morley, E., Silva, H., Bennett, A., Robert, D., and Shallcross, D.: Urban and rural measurements of atmospheric potential gradient, *Journal of Electrostatics*, 97, 42–50, <https://doi.org/https://doi.org/10.1016/j.elstat.2018.11.006>, 2019.
- 590 McMurtry, P. H.: The History of Condensation Nucleus Counters, *Aerosol Sci. Technol.*, 33, 297–322, <https://doi.org/10.1080/02786820050121512>, 2000.
- Michnowski, S., Odzimek, A., Kleimenova, N. G., Kozyreva, O. V., Kubicki, M., Klos, Z., Israelsson, S., and Nikiforova, N. N.: Review of Relationships Between Solar Wind and Ground-Level Atmospheric Electricity: Case Studies from Hornsund, Spitsbergen, and Swider, Poland, *Surv. Geophys.*, 42, 757–801, <https://doi.org/10.1007/s10712-021-09639-3>, 2021.
- 595 Miller, S. and Bodhaine, B.: Supersaturation and expansion ratios in condensation nuclei counters: an historical perspective, *J. Aerosol Sci.*, 13, 481–490, [https://doi.org/10.1016/0021-8502\(82\)90014-3](https://doi.org/10.1016/0021-8502(82)90014-3), 1982.
- Mohnen, V. and Hidy, G. M.: Measurements of Atmospheric Nanoparticles (1875–1980), *Bull. Amer. Meteorol. Soc.*, 91, 1525 – 1540, <https://doi.org/10.1175/2010BAMS2929.1>, 2010.
- März, F., Sători, G., and Zieger, B.: Variations in Schumann resonances and their relation to atmospheric electric parameters at Nagycenk station, *Ann. Geophys.*, 15, 1604–1614, <https://doi.org/10.1007/s00585-997-1604-y>, 1997.
- 600 Nicoll, K., Harrison, R., Barta, V., Bor, J., Brugge, R., Chillingarian, A., Chum, J., Georgoulas, A., Guha, A., Kourtidis, K., Kubicki, M., Mareev, E., Matthews, J., Mkrtchyan, H., Odzimek, A., Raulin, J.-P., Robert, D., Silva, H., Tacza, J., Yair, Y., and Yaniv, R.: A global atmospheric electricity monitoring network for climate and geophysical research, *J. Atmos. Sol.-Terr. Phys.*, 184, 18–29, <https://doi.org/10.1016/j.jastp.2019.01.003>, 2019.
- 605 Parkinson, W. L. and Torreson, O. W.: The diurnal variation of the electric potential of the atmosphere over the oceans, *Union Terr. Magn. Electr. Bull.*, 8, 340–341, 1931.
- Pietruczuk, A. and Jarosławski, J.: Analysis of particulate matter concentrations in Mazovia region, Central Poland, based on 2007–2010 data, *Acta Geophys.*, 61, 445–462, <https://doi.org/10.2478/s11600-012-0069-x>, 2013.
- Podzimek, J., J.C., C., and Yue, P.: Comparison of several aitken nuclei counters, *Atmospheric Environment* (1967), 16, 1–11, [https://doi.org/https://doi.org/10.1016/0004-6981\(82\)90309-2](https://doi.org/https://doi.org/10.1016/0004-6981(82)90309-2), 1982.
- 610 Rycroft, M. J., Israelsson, S., and Price, C.: The global atmospheric electric circuit, solar activity and climate change, *J. Atmos. Sol.-Terr. Phys.*, 62, 1563–1576, [https://doi.org/10.1016/S1364-6826\(00\)00112-7](https://doi.org/10.1016/S1364-6826(00)00112-7), 2000.
- Ryś, A. and Samek, L.: Yearly Variations of Equivalent Black Carbon Concentrations Observed in Krakow, Poland, *Atmosphere*, 13, 539, <https://doi.org/10.3390/atmos13040539>, 2022.
- 615 Sapkota, B. K. and Varshneya, N. C.: On the global atmospheric electric circuit, *J. Atmos. Terr. Phys.*, 52, 1–22, [https://doi.org/10.1016/0021-9169\(90\)90110-9](https://doi.org/10.1016/0021-9169(90)90110-9), 1990.

- Schonland, B. F. J.: Atmospheric electricity, Methuen and Co., London, second ed., 1953.
- Shatalina, M. V., Mareev, E. A., Klimenko, V. V., Kuterin, F. A., and Nicoll, K. A.: Experimental Study of Diurnal and Seasonal Variations in the Atmospheric Electric Field, *Radiophys. Quant. Electr.*, 62, 183–191, <https://doi.org/10.1007/s11141-019-09966-x>, 2019.
- 620 Swider, W.: Ionic mobility, mean mass, and conductivity in the middle atmosphere from near ground level to 70 km, *Radio Sci.*, 23, 389–399, <https://doi.org/10.1029/RS023i003p00389>, 1988.
- Tacza, J., Raulin, J.-P., Macotela, E., Marun, A., Fernandez, G., Bertoni, F., Lima, L., Samanes, J., Buleje, Y., Correia, E., Alves, G., and Makita, K.: Local and global effects on the diurnal variation of the atmospheric electric field in South America by comparison with the Carnegie curve, *Atmos. Res.*, 240, 104 938, <https://doi.org/10.1016/j.atmosres.2020.104938>, 2020.
- 625 Tacza, J., Raulin, J. P., Morales, C. A., Macotela, E., Marun, A., and Fernandez, G.: Analysis of long-term potential gradient variations measured in the Argentinian Andes, *Atmos. Res.*, 248, 105200, <https://doi.org/10.1016/j.atmosres.2020.105200>, 2021.
- Tinsley, B. A. and Zhou, L.: Initial results of a global circuit model with variable stratospheric and tropospheric aerosols, *J. Geophys. Res.-Atmos.*, 111, D16 205, <https://doi.org/10.1029/2005JD006988>, 2006.
- Warzecha, S., ed.: *Annuaire Meteorologique et de l'electricite atmospherique 1957*, vol. 16 of *Travaux de l'Observatoire Géophysique de St. Kalinowski à Świder*, 1960.
- 630 Warzecha, S., ed.: *Annuaire Meteorologique et de l'electricite atmospherique 1965*, vol. 38 of *Travaux de l'Observatoire Géophysique de St. Kalinowski à Świder*, 1968.
- Warzecha, S.: Wpływ czynników meteorologicznych na koncentrację atmosferycznych jąder kondensacji w Świdrze., Phd thesis, Instytut Geofizyki, Polska Akademia Nauk, 1974.
- 635 Warzecha, S.: Periodical variations of the condensation nuclei content at Świder, *Publs. Inst. Geophys. Pol. Acad. Sci.*, 99, 17–35, 1976.
- Warzecha, S.: Variations of the Electric Field and air conductivity at Swider in the years 1958-1985, *Publs. Inst. Geophys. Pol. Acad. Sci.*, 238, 193pp., 1991.
- Warzecha, S. and Warzechowa, K.: Wyniki pomiarów stężenia atmosferycznych jąder kondensacji w przybrzeżnej strefie morza w Lubiatowie, *Archive of the Institute of Geophysics Polish Academy of Sciences, Institute of Geophysics Polish Academy of Sciences*, 1979.
- 640 Williams, E. R.: The global electrical circuit: A review, *Atmos. Res.*, 91, 140–152, <https://doi.org/10.1016/j.atmosres.2008.05.018>, 2009.
- Wilson, C.: Condensation of Water Vapour in the Presence of Dust-Free Air and Other Gases., *Phil.Trans.R.Soc.London*, 189, 265–307, 1897.
- Witkowski, A.: Spostrzeżenia nad elektrycznością atmosferyczną w Zakopanem, *Bull. Acad. Sci. Cracovie A*, 42, 7–10, 1902.
- Wright, M., Matthews, J., Silva, H., Bacak, A., Percival, C., and Shallcross, D.: The relationship between aerosol concentration and atmospheric potential gradient in urban environments, *Science of The Total Environment*, 716, 134 959, <https://doi.org/https://doi.org/10.1016/j.scitotenv.2019.134959>, 2020.
- 645

**Table 4.** Mean values of PG measured for fair weather conditions by month for all and limited CN concentration values together with the standard deviation of the mean value corresponding to 6, 12, 18 UT.

Hour	PG (V/m)	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
All	CN all	349±5	361±5	320±4	239±3	205±2	187±2	191±2	200±2	230±3	257±4	290±5	339±6
	CN<10000	323±11	283±9	307±10	229±7	202±4	184±4	190±3	201±4	214±6	247±8	255±9	309±11
	CN<8000	321±15	274±13	300±14	230±8	202±6	187±5	192±4	198±5	212±8	235±9	254±12	303±14
	CN<6000	274±19	266±20	270±14	219±10	195±8	187±7	187±5	197±6	216±10	229±14	232±15	294±22
6 UT	CN all	294±9	286±9	298±8	248±5	227±4	207±4	216±3	220±4	226±6	213±6	256±10	295±11
	CN<10000	273±15	236±14	283±16	233±10	211±6	195±5	213±5	212±6	197±10	208±12	232±17	279±15
	CN<8000	258±22	215±16	277±20	240±13	215±9	195±7	218±7	206±6	188±15	208±14	243±21	287±21
	CN<6000	209±20	200±20	228±20	236±21	211±11	179±12	209±10	211±10	212±30	210±18	204±19	256±29
12 UT	CN all	372±8	378±8	298±6	202±4	170±4	165±4	171±3	170±3	214±4	259±5	298±7	356±9
	CN<10000	367±21	310±14	299±13	208±7	169±7	155±7	168±4	174±5	222±7	240±9	262±12	343±24
	CN<8000	380±29	295±18	311±17	210±9	173±8	158±10	167±5	175±6	216±9	233±10	269±17	332±37
	CN<6000	334±29	333±39	303±13	206±10	166±11	184±13	169±6	175±7	221±11	233±17	233±32	357±36
18 UT	CN all	364±10	395±10	355±7	253±5	202±3	178±3	178±3	201±3	247±6	287±8	301±10	351±11
	CN<10000	339±17	303±17	359±24	250±15	205±6	181±5	179±4	211±7	221±12	304±18	271±19	316±16
	CN<8000	338±22	309±24	333±48	261±22	196±10	188±6	186±5	216±11	232±23	289±25	254±19	303±22
	CN<6000	344±38	293±35	305±47	240±35	190±22	194±11	185±7	229±21	203±26	271±73	268±27	331±55

**Table 5.** Model of ion conductivity with CN concentrations dominated by soot: concentrations of the CN components, the ion-aerosol loss term  $\beta N$ , ion conductivity, and conductivity ratios at different concentration ratio of water soluble particles and soot components. A constant concentration of insoluble aerosol at  $500 \text{ cm}^{-3}$  is added to the total CN concentration.

<i>Soot</i> (%)	<i>WatSol</i> (%)	$N_{Soot}$ ( $\text{cm}^{-3}$ )	$N_{WatSol}$ ( $\text{cm}^{-3}$ )	CN ( $\text{cm}^{-3}$ )	$\beta N$ ( $10^{-2} \text{ s}^{-1}$ )	$\beta_{eff}$ ( $10^{-6} \text{ cm}^3 \text{ s}^{-1}$ )	Conductivity ( $10^{-15} \text{ S/m}$ )	s	
100	0	26000	0	26500	2.85	1.07	4.20	1.00	
90	10	23400	600	24500	2.77	1.13	4.32	1.03	
80	20	20800	1200	22500	2.70	1.20	4.43	1.05	
70	30	18200	1800	20500	2.62	1.28	4.56	1.08	W mean
60	40	15600	2400	18500	2.55	1.38	4.69	1.11	W median
50	50	13000	3000	16500	2.48	1.50	4.82	1.15	
40	60	10400	3600	14500	2.40	1.66	4.97	1.18	S mean
30	70	7800	4200	12500	2.33	1.86	5.13	1.22	S median
20	80	5200	4800	10500	2.26	2.15	5.29	1.26	
10	90	2600	5400	8500	2.18	2.57	5.47	1.30	
0	100	0	6000	6500	2.11	3.24	5.66	1.35	

**Table 6.** Model of ion conductivity with CN concentrations dominated by water soluble particles: concentrations of the CN components, the ion-aerosol loss term  $\beta N$ , ion conductivity, and conductivity ratios at different concentration ratio of water soluble particles and soot components. A constant concentration of insoluble aerosol at  $200 \text{ cm}^{-3}$  is added to the total CN concentration.

<i>Soot</i> (%)	<i>WatSol</i> (%)	$N_{Soot}$ ( $\text{cm}^{-3}$ )	$N_{WatSol}$ ( $\text{cm}^{-3}$ )	CN ( $\text{cm}^{-3}$ )	$\beta N$ ( $10^{-2} \text{ s}^{-1}$ )	$\beta_{eff}$ ( $10^{-6} \text{ cm}^3 \text{ s}^{-1}$ )	Conductivity ( $10^{-15} \text{ S/m}$ )	s	
0	100	0	28000	28200	3.57	1.37	3.10	1.00	
10	90	600	25200	26000	3.57	1.37	3.36	1.08	
20	80	1200	22400	23800	3.27	1.37	3.66	1.18	
30	70	1800	19600	21600	2.97	1.38	4.03	1.30	W mean
40	60	2400	16800	19400	2.67	1.38	4.47	1.44	W median
50	50	3000	14000	17200	2.38	1.38	5.02	1.62	
60	40	3600	11200	15000	2.08	1.38	5.73	1.85	S mean
70	30	4200	8400	12800	1.78	1.39	6.67	2.15	S median
80	20	4800	5600	10600	1.48	1.40	7.97	2.57	
90	10	5400	2800	8400	1.19	1.41	9.88	3.18	
100	0	6000	0	6200	0.89	1.44	13.0	4.17	



**Table 7.** Mean values of fair weather positive conductivity by season for all and limited CN concentration values together with the one standard error of the mean.

	Winter	Spring	Summer	Autumn	All year
All values	$2.24 \pm 0.01$	$3.20 \pm 0.01$	$3.98 \pm 0.01$	$3.09 \pm 0.01$	$3.28 \pm 0.01$
CN all	$2.16 \pm 0.02$	$3.14 \pm 0.03$	$4.15 \pm 0.02$	$2.98 \pm 0.02$	$3.27 \pm 0.01$
CN<10000	$2.40 \pm 0.06$	$3.69 \pm 0.09$	$4.37 \pm 0.04$	$3.35 \pm 0.06$	$3.79 \pm 0.03$
CN<8000	$2.38 \pm 0.07$	$3.72 \pm 0.10$	$4.40 \pm 0.05$	$3.50 \pm 0.07$	$3.85 \pm 0.04$
CN<6000	$2.52 \pm 0.11$	$3.71 \pm 0.15$	$4.48 \pm 0.06$	$3.36 \pm 0.09$	$3.91 \pm 0.06$

**Table B1.** Number of hourly values of PG at fair weather hours divided into months for January 1965 – December 2005. Last column indicates the number of hours without PG measurements.

Season Y\M	Winter			Spring			Summer			Autumn			total	missing
	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov		
1965	105	211	148	315	205	275	286	342	332	440	270	58	2987	59
1966	125	85	193	226	340	289	462	391	476	407	324	99	3417	91
1967	104	242	268	256	288	383	370	443	445	395	385	222	3801	152
1968	138	100	146	340	427	342	384	339	529	364	244	116	3469	109
1969	181	257	177	308	346	361	401	374	354	394	235	110	3498	111
1970	31	199	197	152	224	271	410	297	324	203	91	113	2512	24
1971	33	275	106	241	309	404	300	512	526	221	207	143	3277	99
1972	300	214	276	279	200	320	347	433	237	149	251	127	3133	68
1973	174	184	46	346	287	311	326	273	552	305	209	181	3194	107
1974	104	236	217	501	347	239	261	201	432	376	150	207	3271	142
1975	137	195	204	256	191	339	309	373	473	430	167	182	3256	103
1976	94	124	321	178	281	321	318	397	382	289	199	140	3044	64
1977	96	252	98	291	227	299	390	304	287	281	189	146	2860	68
1978	129	237	120	230	271	374	351	268	143	67	69	41	2300	14
1979	128	45	237	150	233	401	290	119	171	156	346	63	2339	110
1980	95	126	86	242	190	247	167	90	181	143	179	99	1845	47
1981	109	102	165	225	228	309	229	243	277	185	132	35	2239	87
1982	21	129	158	316	136	383	230	373	408	361	239	200	2954	98
1983	126	60	131	180	264	271	359	356	461	253	150	89	2700	67
1984	56	63	259	281	346	279	130	234	441	149	275	156	2669	87
1985	70	164	153	124	326	399	205	358	373	176	89	76	2513	42
1986	108	95	303	295	323	421	351	311	334	146	259	136	3082	85
1987	69	212	147	292	208	284	182	343	288	288	431	132	2876	80
1988	84	174	168	187	409	328	286	419	370	203	403	171	3202	42
1989	156	75	152	262	256	425	201	403	296	346	145	64	2781	54
1990	106	119	208	264	318	355	328	282	392	174	320	25	2891	82
1991	128	197	123	165	309	311	281	405	333	422	291	183	3148	73
1992	224	166	100	275	243	391	470	485	543	459	195	174	3725	74
1993	119	326	164	366	401	517	411	398	427	252	274	157	3812	56
1994	104	178	225	169	373	407	380	614	378	245	248	231	3552	81
1995	184	156	244	174	317	352	341	553	496	188	185	161	3351	42
1996	304	302	220	272	456	279	398	329	447	186	247	227	3667	152
1997	155	191	221	370	331	373	334	285	536	319	219	180	3514	65
1998	172	281	222	379	290	405	297	312	274	342	265	146	3385	243
1999	202	220	110	360	303	442	271	471	434	478	259	214	3764	79
2000	137	123	221	200	444	543	464	265	426	425	595	265	4108	100
2001	143	156	245	310	255	479	335	316	511	232	249	171	3402	90
2002	273	158	336	370	348	491	398	441	525	377	180	289	4186	39
2003	298	216	277	314	352	482	382	373	501	455	186	166	4002	50
2004	180	153	162	221	431	319	333	415	463	453	292	187	3609	25
2005	41	219	233	356	428	453	390	430	536	552	455	206	4299	42
total	5543	7217	7787	11038	12461	14874	13358	14570	16314	12286	10098	6088	131634	3303
	20547			38373			44242			28472				

**Table B2.** Number of fair weather days (24h) divided into months for the period January 1965 – December 2005.


Season Y\M	Winter			Spring			Summer			Autumn			total
	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	
1965	–	2	–	3	–	4	3	5	6	10	2	–	35
1966	–	–	3	1	5	5	11	1	11	4	2	–	43
1967	1	4	3	1	3	10	3	6	2	7	3	3	46
1968	1	–	1	3	10	–	6	3	7	4	2	2	39
1969	3	5	1	4	6	5	3	7	5	8	1	–	48
1970	–	2	1	1	2	4	6	1	5	2	–	–	24
1971	–	2	–	1	5	4	3	12	11	3	2	–	43
1972	4	4	6	6	2	3	5	9	2	1	1	–	43
1973	1	3	–	3	2	4	5	1	11	2	1	1	34
1974	–	4	3	11	4	2	1	–	10	6	–	2	43
1975	–	3	1	3	1	2	3	7	12	6	4	1	43
1976	–	2	6	1	1	5	7	7	5	4	1	–	39
1977	2	2	–	2	1	2	5	4	4	2	1	1	26
1978	–	5	1	2	3	7	5	1	1	–	–	–	25
1979	–	–	3	–	3	3	2	–	2	–	3	–	16
1980	1	–	–	–	2	–	–	–	–	–	–	–	3
1981	–	–	1	1	3	2	1	1	3	–	–	–	12
1982	–	–	–	4	–	10	4	2	6	2	3	2	33
1983	–	–	–	3	2	1	3	2	7	3	–	–	21
1984	–	–	5	3	3	1	–	3	5	–	1	2	23
1985	–	2	1	–	3	7	1	–	5	–	–	–	19
1986	1	–	6	4	5	6	2	3	3	1	2	–	33
1987	–	1	1	6	1	3	1	4	4	2	7	–	30
1988	1	2	1	–	7	3	2	5	1	1	6	3	32
1989	1	–	–	2	3	7	1	7	2	5	–	–	28
1990	–	–	–	4	6	6	1	2	6	–	2	–	27
1991	–	1	–	1	3	3	1	6	2	4	3	–	24
1992	2	–	–	3	1	5	4	9	9	10	1	–	44
1993	–	6	–	5	8	7	7	4	6	2	3	1	49
1994	–	2	1	–	2	3	7	16	2	–	3	4	40
1995	1	1	–	–	4	3	3	10	7	–	–	–	29
1996	2	4	–	1	6	–	9	–	5	1	2	1	31
1997	1	–	1	3	2	6	3	–	13	3	1	3	36
1998	1	2	–	5	1	5	4	3	3	4	2	1	31
1999	1	1	–	6	4	6	3	9	2	10	2	2	46
2000	1	–	1	–	6	12	8	1	5	6	16	1	57
2001	–	–	1	–	3	6	1	–	6	1	1	–	19
2002	–	–	3	6	2	6	5	8	14	5	–	3	52
2003	4	1	4	6	4	8	6	3	5	6	–	–	47
2004	2	–	3	1	10	2	1	3	3	5	4	–	34
2005	–	2	4	5	4	9	–	5	10	15	11	2	67
total	31	63	62	111	143	187	146	170	228	145	93	35	1414
	156			441			544			273			

**Table B3.** Number of PG values measured in FW conditions by month for all and limited CN concentrations corresponding to 6, 12, 18 UT.


Hour	PG length	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
6UT	all CN	239	269	418	545	737	653	702	754	475	347	190	190
	PG with all CN	236	258	397	526	709	628	670	723	449	326	178	183
	CN<10000	52	45	63	102	240	217	242	218	88	55	50	55
	CN<8000	27	25	37	52	139	118	138	123	36	34	35	34
	CN<6000	16	13	22	21	56	45	58	55	9	13	18	18
12 UT	all CN	363	388	467	382	409	329	377	523	481	508	336	301
	PG with all CN	356	377	453	374	395	322	367	514	476	497	329	298
	CN<10000	41	47	55	76	79	74	137	160	109	94	56	35
	CN<8000	20	25	31	63	51	44	108	110	72	52	25	16
	CN<6000	8	7	19	42	27	25	73	72	36	17	9	8
18 UT	all CN	322	393	544	621	718	676	713	772	599	467	293	255
	PG with all CN	318	377	526	592	682	638	671	722	564	418	277	246
	CN<10000	47	44	36	54	153	222	270	166	65	48	43	55
	CN<8000	29	26	14	22	70	123	151	79	25	18	25	28
	CN<6000	8	14	8	8	19	49	65	23	12	4	13	5

# Time series of monthly values of fair weather PG and CN


SWI 1965-2005 yearbook data




**Figure 1.** Time series of monthly averaged values of fair weather PG and CN values over January 1965 – December 2005. The dashed red lines indicate positive trend for PG ( $2.5 \pm 0.3$  V/m per year) and negative trend for CN ( $-120 \pm 20$   $\text{cm}^{-3}$ ).



**Figure 2.** Histogram of the PG values measured during fair weather conditions in January 1965 – December 2005: black colour – all PG values, red colour – PG values recorded simultaneously with CN concentrations, green colour – the PG values at CN concentrations below  $10000 \text{ cm}^{-3}$ . The black dashed line represents mean PG value (230 V/m). The other statistical parameters separately for the three PG groups are given in the legend, including the number of data values, N.




**Figure 3.** Histogram of hourly averaged CN concentrations at fair weather conditions in January 1965 – December 2005: CN below 10000  $\text{cm}^{-3}$  are framed in green, red colour – all CN. The black dashed line represents mean CN concentration value of 18980  $\text{cm}^{-3}$ .




**Figure 4.** Seasonal histograms of the hourly averaged (6, 12, 18 UT) CN concentration values measured during fair weather conditions for the period January 1965 – December 2005: red colour – all CN concentration values, green colour – CN concentration values below  $10000 \text{ cm}^{-3}$ . Note that Y-axis ranges for summer (0-1000 counts) is twice compared with the other seasons (0-500 counts).







**Figure 5.** Diurnal variation of PG measured during fair weather conditions by seasons and for the whole year in January 1965 – December 2005. The error bars represent  $\pm 1$  standard deviation.




**Figure 6.** Diurnal variation of CN concentration values for cloudless days in spring 1968 – winter 1968/1969 and mean CN concentrations measured for 6, 12, 18 UT in 1965–2005.




**Figure 7.** Seasonal histograms of the hourly averaged (6, 12, 18 UT) PG values measured during fair weather conditions in January 1965 – December 2005: red colour–PG values measured simultaneously with all CN concentration values (6, 12, 18 UT), green colour–PG values measured simultaneously with CN concentrations below  $10000 \text{ cm}^{-3}$ . Note that the Y-axis ranges for summer (0-3000 counts) is twice compared with the other seasons (0-1500 counts).




**Figure 8.** Seasonal dependence of hourly averaged values of fair weather PG and CN concentration values in January 1965 – December 2005. Horizontal dashed line indicates the limit of CN concentration at 10000 cm<sup>-3</sup>.




**Figure 9.** Histograms of the limited CN concentration values measured during fair weather conditions divided into CN range: below 10000  $\text{cm}^{-3}$  (purple), 8000  $\text{cm}^{-3}$  (green), and 6000  $\text{cm}^{-3}$  (blue), Bottom panels: histograms of hourly averaged PG values measured at limited CN concentration.



**Figure 10.** Diurnal variation of PG values measured during fair weather conditions at 6, 12, 18 UT, divided into seasons, for the period January 1965 – December 2005 for all CN concentration values (black), and CN concentration values below: 10000  $\text{cm}^{-3}$  (brown), 8000  $\text{cm}^{-3}$  (green), 6000  $\text{cm}^{-3}$  (blue). Error bars indicate one standard error of the mean.



**Figure 11.** Annual variation of PG values measured during fair weather conditions for all and limited CN concentration values for all hours (upper chart) and separately for each hour (6, 12, 18 UT). Error bars indicate one standard error of the mean.



**Figure 12.** Annual variation of positive conductivity values measured during fair weather conditions for all and limited CN concentration values for all hours (upper chart) and separately for each hour (6, 12, 18 UT). Error bars indicate one standard error of the mean.