



# Acoustic–gravity waves and their role in ionosphere–lower thermosphere coupling

Gordana Jovanovic

University of Montenegro, Faculty of Science and Mathematics

Dzordza Vasingtona bb, 81000 Podgorica, Montenegro

**Correspondence:** Gordana Jovanovic (gordanaj@ucg.ac.me)

**Abstract.** The properties of acoustic–gravity waves (AGWs) in the ionospheric D layer and their role in the D layer–lower thermosphere coupling are studied using the dispersion equation and the reflection coefficient. These analytical equations are an elegant tool for evaluating the contribution of upward–propagating acoustic and gravity waves to the dynamics of the lower thermosphere. It was found that infrasound waves with frequencies  $\omega > 0.035s^{-1}$ , which propagate almost vertically, can reach the lower thermosphere. Also, gravity waves with frequencies lower than  $\omega < 0.0087s^{-1}$ , with horizontal phase velocities in the range  $159m/s < v_h < 222m/s$ , and horizontal wavelength  $115 km < \lambda_p < 161 km$ , are important for the lower thermosphere dynamics. These waves can cause temperature rise in the lower thermosphere and have the potential to generate middle–scale traveling ionospheric disturbances (TIDs). The reflection coefficient for AGWs is highly temperature dependent. During maximum solar activity, the temperature of the lower thermosphere can rise several times. This is the situation where infrasound waves become a prime candidate for the ionospheric D layer–lower thermosphere coupling, since strongly reflected gravity waves remain trapped in the D layer. Knowing the temperatures of the particular atmospheric layers, we can also know the characteristics of AGWs and vice versa.

## 1 Introduction

Acoustic–gravity waves (AGWs) are able to transport energy and momentum between different layers of the atmosphere. Understanding these waves is essential if we want to comprehend the atmosphere as a system where the layers are coupled. The ionosphere is a part of the Earth’s atmosphere located between about 60 km and 1000 km where the charged particles significantly influence its physical and chemical properties (Mitra , 1974; Bothmer and Daglis , 2007). Knowledge of typical AGW characteristics in the ionosphere is important for modelling of coupling between the ionized and neutral atmosphere. Ionosphere is constantly exposed to various influences from outer space as well as from the terrestrial atmosphere and lithosphere. The non–periodic and sudden events, such as solar flares (Singh et al. , 2014; Nina et al. , 2017; Chum et al. , 2018), coronal mass ejection (Bochev and Dimitrova , 2003; Balan et al. , 2008), solar eclipses (Singh et al. , 2012), supernova explosions



followed by hard X and  $\gamma$  radiation (Inan et al. , 2007), lightnings (Voss et al. , 1998), and some processes in the terrestrial lithosphere like volcanic eruptions and earthquakes (Nenovski et al. , 2010; Argunov and Gotovtsev, 2019), induce space and time varying ionospheric perturbations. These disturbances cause numerous complex physical, chemical and dynamical phenomena in the ionosphere (Rozhnoi , 2012; Hayakawa et al. , 2010) and may directly affect human activities, especially in the telecommunications.

The atmospheric monitoring depends on the altitude of the considered atmospheric layer. The ionospheric D layer at an altitude of about 60 to 90 km, lies below the area being studied by satellite observations and above the region where balloon measurements find their application. Therefore, its monitoring is based on rocket and radar measurements and on the propagation of very low and low-frequency (VLF/LF) radio waves (Nina and Čadež , 2013). In this way, it is possible to observe a large part of the low ionosphere, detect local perturbations, and sudden events.

The ionospheric D layer and lower thermosphere below 140 km, where AGWs with the specific frequencies and wavelengths are detected, are the focus of this article. We considered the conditions for propagation of AGWs in the D ionospheric layer and their reflection/transmission on the plane boundary between this layer and the lower thermosphere. This is a way to study the coupling between the ionosphere and the lower thermosphere and to analyze the influence of AGWs on thermospheric processes and characteristics.

The article is structured as follows: Section 2 contains the basic theory of AGWs and the derivation of their dispersion equation. Section 3 presents the analytical equation for the AGW reflection coefficient. In Section 4, the propagation of AGWs through the ionospheric D layer as well as their reflection/transmission properties are analyzed. Discussion and conclusions are displayed in Sections 5 and 6, respectively.

## 2 Basic equations

The D layer is part of the ionosphere, where typical atmosphere models give  $n_n \sim 10^{21} m^{-3}$  for the neutral particle density and  $n_p \sim 10^8 m^{-3}$  for charged plasma particles, and where electric and magnetic effects play a minor role in the local atmosphere dynamics. This is why hydrodynamic (HD) equations, rather than magneto-hydrodynamic (MHD) equations, can be used to analyze wave propagation. The standard set of HD equations describes the dynamics of adiabatic processes in a fully ionized hydrogen plasma in the presence of gravity  $\mathbf{g} = -g\mathbf{e}_z$  with constant acceleration  $g = 9.81 m/s^2$ :

continuity and ideal gas equation

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{v}) = 0, \quad p = \rho RT, \quad (1)$$

momentum equation

$$\rho \left( \frac{\partial \mathbf{v}}{\partial t} + \mathbf{v} \cdot \nabla \mathbf{v} \right) = -\nabla p + \rho \mathbf{g} \quad (2)$$

and an adiabatic law for a perfect gas

$$\frac{\partial p}{\partial t} + \mathbf{v} \cdot \nabla p = \frac{\gamma p}{\rho} \left( \frac{\partial \rho}{\partial t} + \mathbf{v} \cdot \nabla \rho \right). \quad (3)$$



Here,  $R = R_0/M$  is the individual gas constant for molecules with molar mass  $M$ ,  $R_0 = 8.314 \text{ J/molK}$  is the universal gas constant and  $\gamma = c_p/c_v = (j + 2)/j$  is the ratio of specific heats for gas particle with  $j = 5$  degrees of freedom.

## 2.1 Dispersion equation for AGWs

In what follows, we consider waves whose wavelengths are sufficiently small in comparison with the Earth radius  $R_E = 6371$  km. Therefore, the plane parallel geometry can be applied in a locally isothermal medium. Under these assumptions, the atmosphere is taken to be vertically stratified, initially in hydrostatic equilibrium, and then perturbed by harmonic waves of small amplitude. This means that Eqs. (1)–(3) can be linearized by taking any physical quantity  $\psi(x, y, z, t)$  as a sum of its basic state unperturbed value  $\psi_0(z)$  and a small first order perturbation  $\delta\psi(x, y, z, t)$ , i.e.  $\psi(x, y, z, t) = \psi_0(z) + \delta\psi(x, y, z, t)$ , where:  $\delta\psi(x, y, z, t) = \psi'(z)e^{i(k_x x + k_y y - \omega t)}$ , and  $|\psi'| \ll |\psi_0|$ . Eqs. (1)–(3), linearized with these perturbations, reduce to three equations: one for the unperturbed basic state and two for small perturbations. The unperturbed basic state is described by:

$$\frac{d}{dz} \ln \rho_0(z) + \frac{1}{H} = 0, \quad p_0 = \rho_0 R T_0, \quad \text{with } T_0 = \text{const},$$

whose solution is:

$$\rho_0(z) = \rho_0(0)e^{-z/H} \quad \text{or} \quad p_0(z) = p_0(0)e^{-z/H}, \quad (4)$$

where  $H = p_0(0)/\rho_0(0) = v_s^2/\gamma g = \text{const}$  is the characteristic scale-height of the isothermal atmosphere.

The small perturbations are governed by equations (Jovanovic, 2016):

$$\frac{d\xi'_z}{dz} = C_1 \xi'_z - C_2 p', \quad \frac{dp'}{dz} - g \frac{d\rho_0}{dz} \xi'_z = C_3 \xi'_z - C_1 p', \quad (5)$$

where  $\xi'_z = i v'_z / \omega$  is the  $z$ -component (i.e. the vertical component) of the fluid displacement, while  $p'$  is the pressure perturbation. The coefficients in Eqs. (5) are:

$$C_1 = \frac{g}{v_s^2}, \quad C_2 = \frac{\omega^2 - k_p^2 v_s^2}{\rho_0(z) v_s^2 \omega^2}, \quad C_3 = \rho_0(z) \left( \omega^2 + \frac{g^2}{v_s^2} \right). \quad (6)$$


The density distribution  $\rho_0(z)$  is given by Eq. (4) and  $k_p^2 = k_x^2 + k_y^2$  designates square of the horizontal wavenumber. The Eqs. (5)–(6) allow the following solutions for the vertical displacement  $\xi'_z$  and the pressure perturbation  $p'$ :

$$\xi'_z(z) = \xi'_z(0) e^{\frac{z}{2H}} e^{i k_z z}, \quad p'(z) = p'(0) e^{\frac{z}{2H}} e^{i k_z z}. \quad (7)$$

Eqs. (5) with solutions Eqs. (7) yield the dispersion equation for AGWs:

$$k_z^2 = \frac{\omega^2(\omega^2 - \omega_{co}^2) - k_p^2 v_s^2(\omega^2 - \omega_{BV}^2)}{\omega^2 v_s^2}. \quad (8)$$

Here,  $k_z$  is the vertical wavenumber,  $\omega_{co}^2 = \gamma^2 g^2 / 4 v_s^2 = v_s^2 / 4 H^2$  is the square of the acoustic wave cutoff frequency, and  $\omega_{BV}^2 = (\gamma - 1) g^2 / v_s^2$  is the square of the Brunt–Väisälä frequency. This equation is quadratic in  $\omega^2$  which indicates the existence of two wave modes in the considered stratified atmosphere: the acoustic and gravity modes. Stratification in a vertical



**Figure 1.** Dispersion curves for AGWs. Two sets of curves are related to acoustic and gravity waves, which cannot propagate below the acoustic cutoff frequency  $\Omega_{co} = \omega_{co}H/v_s$  and above the Brunt–Väisälä frequency  $\Omega_{BV} = \omega_{BV}H/v_s$ , respectively.

direction, caused by gravity and given by Eq. (4), introduces cutoff frequencies—acoustic cutoff frequency below which acoustic waves cannot propagate and Brunt–Väisälä frequency above which gravity waves cannot propagate. Therefore, the branches of acoustic and gravity waves are present. Between them are evanescent waves that do not propagate, Fig. 1. The dispersion equation Eq. (8) can be expressed in terms of wavelengths and wave frequencies  $\omega$  in the following way:

$$\lambda_z^2(\omega) = \frac{A_2(\omega)\lambda_p^2}{\lambda_p^2 - A_0(\omega)}, \quad (9)$$

where

$$A_0(\omega) = \frac{4\pi^2 v_s^2 (\omega^2 - \omega_{BV}^2)}{\omega^2 (\omega^2 - \omega_{co}^2)}, \quad A_2(\omega) = \frac{4\pi^2 v_s^2}{\omega^2 - \omega_{co}^2}.$$

This equation will be useful for further analysis.

The physical quantities in the dispersion equation can be made dimensionless by appropriate scalings:  $K_p = k_p H$ ,  $K_z = k_z H$ ,  $\Omega = \omega H/v_s$ ,  $\Omega_{co} = \omega_{co}H/v_s = 0.5$  and  $\Omega_{BV} = \omega_{BV}H/v_s = \sqrt{\gamma-1}/\gamma = 0.45$ . Now, the dispersion equation Eq. (8) has the dimensionless form:

$$K_z^2 = \Omega^2 - \Omega_{co}^2 - \frac{K_p^2(\Omega^2 - \Omega_{BV}^2)}{\Omega^2}. \quad (10)$$

The AGWs propagate in the vertical direction if  $K_z^2 > 0$ . This is fulfilled when

$$K_p^2 < \frac{\Omega^2(\Omega^2 - \Omega_{co}^2)}{\Omega^2 - \Omega_{BV}^2}, \quad (11)$$

i.e. when dimensionless horizontal phase velocity is

$$V_h^2 = \frac{\Omega^2}{K_p^2} > \frac{\Omega^2 - \Omega_{BV}^2}{\Omega^2 - \Omega_{co}^2}. \quad (12)$$



The AGWs become evanescent when  $K_p^2 > \frac{\Omega^2(\Omega^2 - \Omega_{co}^2)}{\Omega^2 - \Omega_{BV}^2}$  and  $V_h^2 < \frac{\Omega^2 - \Omega_{BV}^2}{\Omega^2 - \Omega_{co}^2}$ . The boundary between propagating and evanescent regions is given by  $K_z = 0$ . Acoustic waves with frequencies close to the acoustic cutoff frequency  $\Omega \approx \Omega_{co} = 0.5$ , are more influenced by gravity than those with high frequencies, when  $\Omega \gg \Omega_{co}$ . Hence, gravity-modified acoustic waves and pure acoustic waves coexist in the stratified atmosphere (Mihalas, 1984). The Eq. (10) shows that the vertical wavenumber  $K_z$  has a maximum value for  $K_p = 0$ , i.e.

$$K_{zmax} = K = \sqrt{\Omega^2 - \Omega_{co}^2}. \quad (13)$$

This equation describes acoustic waves that propagate only in the vertical direction.

Gravity waves, in contrast to acoustic waves, are not able to travel vertically with  $K_p = 0$ , which means there are no pure vertically propagating gravity waves (Mihalas, 1984). Therefore, they propagate obliquely through the stratified atmosphere in accordance with Eq. (10). For the very low frequencies, when  $\Omega \ll \Omega_{BV} = 0.45$ , gravity waves propagate with:

$$K_z \approx \frac{K_p \Omega_{BV}}{\Omega}, \quad i.e., \quad \frac{\lambda_p}{\lambda_z} \approx \frac{\omega_{BV}}{\omega}. \quad (14)$$

Dimensionless equations are used because they are valid in each stratified medium, like Earth's, planets or the solar atmosphere.

When we rewrite them using characteristic frequencies and temperatures, we obtain the equations for particular atmospheric layers as will be done in Section 4 for the ionospheric D layer.

### 3 Reflection coefficient of AGWs

The considered basic state in the stratified atmosphere is composed of two half-spaces with constant sound speeds, separated by a horizontal plane boundary  $z = 0$ . The two regions are characterized by the corresponding plasma densities  $\rho_{01}$  and  $\rho_{02}$  adjacent to the lower and upper side of the boundary  $z = 0$ . The unperturbed density profile can be expressed as follows:

$$\rho_0(z) = \rho_{01} e^{-z/H_1}, \quad z < 0, \quad \text{region (1)}, \quad \rho_0(z) = \rho_{02} e^{-z/H_2}, \quad z > 0, \quad \text{region (2)}, \quad (15)$$

where  $H(n) = v_{sn}^2 / \gamma g$ ,  $n = 1, 2$ . There is a density, pressure, and temperature jump across  $z = 0$ . The boundary condition that has to be applied at  $z = 0$  in the basic state is the continuity of the unperturbed pressure  $p_0$  at  $z = 0$ , (Jovanovic, 2016), which yield:

$$\frac{\rho_{02}}{\rho_{01}} = \frac{v_{s1}^2}{v_{s2}^2} = \frac{T_1}{T_2} = s = const. \quad (16)$$

The boundary conditions for perturbations are continuity of both the vertical fluid displacement  $\xi'_z$  and the pressure perturbation  $p' - g\rho_0(z)\xi'_z$  at the boundary  $z = 0$ . Also, the energy density of the perturbations has to diminish to zero as  $|z|$  tends to infinity. The harmonic wave, which propagates through regions (1) and (2), does not change its frequency, and the horizontal wavevector component  $K_p$ , parallel to the boundary  $z = 0$ . However, the vertical wavevector component  $K_z$  has a discontinuity at the boundary  $z = 0$ , where it changes from  $K_{z1}$  to  $K_{z2}$  according to the dispersion equation Eq. (10). We assume that a wave propagates from the lower region (1) upward towards the boundary  $z = 0$ , and that the waves continuing past it are absorbed



with no reflection in the upper region (2). In this case, in the lower region, the perturbations are the superposition of the incident and reflected waves, while in the upper region, there is only the transmitted wave. The reflection coefficient of AGWs is defined as the square of the absolute value of the reflection amplitude. Using dimensionless physical values for brevity, the reflection coefficient can be written as (see details in Jovanović, 2014):

$$R = \left[ \frac{\left[ \left(1 - \frac{\gamma}{2}\right) \left( \frac{1}{V_h^2 - 1} - \frac{s^2}{sV_h^2 - 1} \right) + \frac{(s-1)}{V_h^2} \right]^2 + \frac{\gamma^2 \Omega^2}{V_{v1}^2} \left( \frac{V_{v1}^2}{V_{v2}^2} \cdot \frac{s^2}{(sV_h^2 - 1)^2} - \frac{1}{(V_h^2 - 1)^2} \right)}{\left[ \left(1 - \frac{\gamma}{2}\right) \left( \frac{1}{V_h^2 - 1} - \frac{s^2}{sV_h^2 - 1} \right) + \frac{(s-1)}{V_h^2} \right]^2 + \frac{\gamma^2 \Omega^2}{V_{v1}^2} \left[ \frac{V_{v1}}{V_{v2}} \cdot \frac{s}{sV_h^2 - 1} + \frac{1}{V_h^2 - 1} \right]^2} \right]^2 + \left[ \frac{\frac{2\gamma\Omega}{V_{v1}(V_h^2 - 1)} \left[ \left(1 - \frac{\gamma}{2}\right) \left( \frac{1}{V_h^2 - 1} - \frac{s^2}{sV_h^2 - 1} \right) + \frac{(s-1)}{V_h^2} \right]}{\left[ \left(1 - \frac{\gamma}{2}\right) \left( \frac{1}{V_h^2 - 1} - \frac{s^2}{sV_h^2 - 1} \right) + \frac{(s-1)}{V_h^2} \right]^2 + \frac{\gamma^2 \Omega^2}{V_{v1}^2} \left[ \frac{V_{v1}}{V_{v2}} \cdot \frac{s}{sV_h^2 - 1} + \frac{1}{V_h^2 - 1} \right]^2} \right]^2. \quad (17)$$

Here,  $V_{v1}$  and  $V_{v2}$  are the vertical phase velocities of AGWs in regions (1) and (2) respectively, given by the equations:

$$V_{v1} = \frac{\Omega}{K_{z1}} = \frac{V_h \Omega}{\sqrt{V_h^2(\Omega^2 - \Omega_{co}^2) - (\Omega^2 - \Omega_{BV}^2)}}, \quad (18)$$

and

$$V_{v2} = \frac{\Omega}{K_{z2}} = \frac{V_h \Omega}{\sqrt{sV_h^2(\Omega^2 - s\Omega_{co}^2) - (\Omega^2 - s\Omega_{BV}^2)}}, \quad (19)$$


while  $V_h$  is horizontal phase velocity given by Eq. (12). If  $V_{v1}^2$  and  $V_{v2}^2$  are positive, AGWs propagate through regions (1) and (2), respectively. If  $V_{v1}^2, V_{v2}^2 < 0$ , these waves are evanescent and not of interest to this study.

## 4 Results

In this Section, the analytical equations derived in Sections 2 and 3 are used to analyze the propagation of AGWs and their reflection/transmission properties in the ionospheric D layer.

### 4.1 AGWs in the ionospheric D layer


Acoustic-gravity waves which propagate in the lower ionosphere below 90 km can be generated from below, where hydrodynamic motions can be induced by atmospheric convective motions (Sindelarova, Buresova, and Chum, 2009), in the lithosphere (Nina et al. 2021; Boudjada et al., 2024), and from above, due to sunrise and sunset effects (Afraimovich et al., 2009; Nina and Čadež, 2013; Nina et al., 2017). These perturbations may result into various patterns of either eigenmodes or driven linear waves in the atmosphere. The focus of this research is on the driven AGWs and their role in the ionosphere and the lower thermosphere coupling. Therefore, propagation of AGWs in the vertical direction is particularly important. For the considered isothermal ionospheric D layer with a temperature  $T = 250$  K and  $\gamma = 1.4$ , sound velocity is  $v_s = \sqrt{\gamma RT} = 317$  m/s, and  $H = 7317$  m. This is in accordance with Lizunov and Hayakawa (2004). For a gravity modified acoustic wave with frequency




**Figure 2.** Vertical wavelength  $\lambda_z$  of acoustic waves from dispersion equation (9) as a function of horizontal wavelength  $\lambda_p$ , for given frequencies  $\omega > \omega_{co} = 0.021s^{-1}$ .

near acoustic cutoff frequency  $\omega = 0.022s^{-1} \geq \omega_{co} = 0.021s^{-1}$ , Eq. (13) enables the calculation of  $\lambda_z \geq \lambda_{zmin} \approx 460$  km. For the pure acoustic wave, with frequency much greater than acoustic cutoff frequency,  $\omega \gg \omega_{co}$ , i.e.  $\omega = 10 \cdot \omega_{co} = 0.21s^{-1}$ , this value is  $\lambda_z \geq \lambda_{zmin} \approx 9.2$  km. It can be noticed that gravity–modified acoustic waves have much longer vertical wave-  
 150 lengths than pure acoustic waves. Therefore, acoustic waves with frequencies near the acoustic cutoff frequency  $\omega_{co}$  have the best chance for vertical propagation through the ionospheric D layer towards the lower thermosphere. Acoustic waves in Fig. 2 are detected in the ionospheric D layer using VLF waves (Nina and Čadež, 2013).

Gravity waves with high frequencies,  $\omega \approx \omega_{BV}$ , and with low frequencies,  $\omega \ll \omega_{BV}$ , are presented in Figs. 3 and 4, respectively. The Eq. (14) shows that low–frequency gravity waves have much longer horizontal than vertical wavelengths, i.e. they  
 155 propagate more horizontally than vertically, Fig. 4. In addition, for a given  $\lambda_p$ , the vertical wavelengths of low–frequency gravity waves are shorter than those of gravity waves with a frequency that is close to the Brunt–Väisälä frequency, Figs. 3 and 4. The vertical phase velocities of these waves are smaller than those of high–frequency gravity waves. Therefore, high–frequency gravity waves propagate faster upward through the ionospheric D layer towards the lower thermosphere. Figures 3 and 4 show gravity waves that were found in the ionospheric D layer (Nina and Čadež, 2013). They can be induced in situ at sunrise and  
 160 sunset due to motions of the solar terminator. Low–frequency gravity waves are observed near OH layer at an altitude of about 87 km and near  $O_2$  layer at an altitude of about 94 km by the mesospheric temperature mapper (Yuan et al., 2016). Their frequencies are  $\omega = 0.0011s^{-1}$  and  $\omega = 0.0014s^{-1}$ , respectively, and are even lower than the gravity wave frequencies in Fig. 4.



**Figure 3.** Vertical wavelength  $\lambda_z$  of gravity waves from dispersion Eq. (9) as a function of horizontal wavelength  $\lambda_p$ , for given frequencies  $\omega \approx \omega_{BV} = 0.0195s^{-1}$ .



**Figure 4.** Vertical wavelength  $\lambda_z$  of gravity waves from dispersion Eq. (9) as a function of horizontal wavelength  $\lambda_p$ , for given frequencies  $\omega \ll \omega_{BV} = 0.0195s^{-1}$ .





## 4.2 Reflection coefficient of AGWs at the D layer–lower thermosphere boundary

165 We assume that  $z = 0$ , Eq. (15), is the plane boundary between the ionospheric D layer at an altitude of 60–90 km, region (1), and lower thermosphere at an altitude of 90 to about 140 km, region (2). At this boundary, AGWs coming from below can be reflected in the D layer or transmitted into the lower thermosphere. The temperature of the D layer is  $T_1 = 250$  K, while the temperature of the lower thermosphere is  $T_2 = 500$  K. Therefore, Eq. (16) gives  $s = 0.5$ . The reflection coefficients of acoustic and gravity waves will be analyzed separately.

### 170 4.2.1 Reflection coefficient of the acoustic waves


Figure 5 shows the reflection coefficient as a function of frequency  $\Omega$ , for acoustic waves at the D layer–lower thermosphere plane boundary  $z = 0$ , when  $s = 0.5$ . Acoustic waves in the frequency range  $\Omega_{co} < \Omega < 0.8$  are reflected at this boundary to a somewhat greater extent. The reflection coefficient strongly decreases with increasing frequency, and acoustic waves with frequencies  $\Omega > 0.8$ , i.e.  $\omega = 0.035s^{-1}$ , can easily propagate through the D layer–lower thermosphere boundary. These waves  
175 could affect the thermospheric temperature and dynamics by depositing their momentum and energy in the lower thermosphere. The value of the horizontal phase velocity  $V_h$  does not significantly affect the reflection coefficient, except in the case when  $V_h = 1/\sqrt{s} = 1.41$ , i.e. for the horizontal phase velocity of the acoustic waves  $v_{ph} = 1.41v_s = 447$  m/s, when total internal reflection occurs. Waves with this horizontal velocity cannot penetrate the thermosphere. Acoustic waves with horizontal phase velocities  $V_h > 1.41$  can propagate through the D layer–lower thermosphere boundary and extend further into the ther-  
180 mosphere, especially if  $\Omega > 1$ . Their reflection coefficient slowly decreases with the increase of  $V_h$  for a given frequency  $\Omega$ .

### 4.2.2 Reflection coefficient of the gravity waves


The reflection coefficient for gravity waves increases when the frequency  $\Omega$  increases and decreases with increasing horizontal phase velocity  $V_h$  for a given frequency  $\Omega$ , Fig. 6. These waves can propagate in both regions, in the ionospheric D layer and  
185 in the lower thermosphere, if their frequencies are lower than the cutoff frequency  $\Omega = \sqrt{s}\Omega_{BV} = 0.32$ , or  $\omega = 0.014s^{-1}$ , and their horizontal phase velocities are lower than  $V_h = \Omega_{BV}/\Omega_{co} = 0.9$ , i.e.  $v_h = 0.9v_s = 285$  m/s. Gravity waves with frequencies much lower than the Brunt–Väisälä frequency and with high horizontal phase velocities are candidates for crossing the D layer–lower thermosphere boundary. Contrary to this, gravity waves with frequencies near the cutoff frequency  $\Omega = \sqrt{s}\Omega_{BV}$  are strongly reflected at the D layer–lower thermosphere boundary. For the horizontal phase velocity  $V_h = 0.9$ , total internal  
190 reflection occurs, and the reflection coefficient is equal to unity.

## 5 Discussion

It is known that high–frequency acoustic waves are strongly absorbed by the atmosphere (Sindelarova, Buresova, and Chum, 2009). The rate of absorption is proportional to the wave frequency squared. Therefore, only acoustic waves with the low



**Figure 5.** Reflection coefficient for acoustic waves at the D layer–lower thermosphere plane boundary  $z = 0$ , as a function of frequency  $\Omega$  and parameter  $s = 0.5$ . If  $V_h = 1/\sqrt{s} = 1.41$ , the reflection coefficient is  $R = 1$  and total internal reflection occurs.




**Figure 6.** Reflection coefficient for gravity waves at the D layer–lower thermosphere plane boundary  $z = 0$ , as a function of frequency  $\Omega$  and parameter  $s = 0.5$ . If  $V_h = \Omega_{BV}/\Omega_{co} = 0.9$ , the reflection coefficient is  $R = 1$  and total internal reflection occurs.



frequencies (infrasound) may propagate through ionospheric D layer and eventually through the lower thermosphere. Indeed, it was found that only acoustic waves with periods less than 4 minutes, i.e.  $\Omega > 0.6$ , or  $\omega > 0.026s^{-1}$  propagating almost vertically are able to reach the lower thermosphere (Blanc , 1985; Schulthess , 2022). In Fig. 5, the reflection coefficient for infrasound waves is presented since the dimensionless frequency  $\Omega = 1.6$  corresponds to frequency  $\omega = 0.069s^{-1}$ , i.e.  $\nu = \omega/2\pi = 0.01$  Hz. These waves, with horizontal phase velocities  $v_h > 447$  m/s and with minimum vertical phase velocity  $v_{vmin} > 317$  m/s, have the best chance of reaching the thermosphere if they propagate almost vertically with infrasound frequencies  $\omega > 0.035s^{-1}$ . Although infrasound waves dissipate their energy in the lower thermosphere, they are not the first option for raising its temperature. Namely, the influence of acoustic wave energetics into the ionosphere/lower thermosphere is weak (Lizunov and Hayakawa , 2004). It appears that the temperature in the thermosphere is increased by low-frequency gravity waves coming from below (Sindelarova, Buresova, and Chum , 2009). Their reflection coefficient for the ionospheric D layer–lower thermosphere boundary is shown in Fig. 6. Gravity waves with horizontal phase velocities  $V_h < 0.5$  are easily reflected from the boundary between the D layer and lower thermosphere and will likely remain trapped at lower altitudes. Only waves with horizontal phase velocities  $0.5 < V_h < 0.7$ , i.e.  $159$  m/s  $< v_h < 222$  m/s, and with low frequencies  $\Omega < 0.2$ , or  $\omega < 0.0087s^{-1}$ , are important for the dynamics of the middle atmosphere. Horizontal wavelengths for these waves are in the range of  $115$  km  $< \lambda_p < 161$  km. This is consistent with the results known from the scientific literature (Fritts et al. , 2014; Bakhmetieva et al. , 2019), which emphasize that gravity waves with periods as short as 10 minutes (i.e.  $\Omega < 0.24$ , or  $\omega < 0.01s^{-1}$  ) can carry significant momentum flux vertically. These waves with wavelengths  $\lambda_p \approx 100 - 200$  km, play an important role in the coupling between the ionospheric D layer and the lower thermosphere. They are responsible for the generation of middle-scale traveling ionospheric disturbances (TIDs) with periods from 15 minutes to 3 hours, velocities from 100 to 250 m/s and horizontal wavelength of approximately a few hundred kilometers (Lizunov and Hayakawa , 2004). It seems that they are causing a rise in temperature in the lower thermosphere through the process of gravity wave breaking and dissipation due to kinematic viscosity and thermal diffusivity (Vadas , 2007; Sindelarova, Buresova, and Chum , 2009; Yuan et al. , 2016). A similar situation can be found in the solar atmosphere (Fleck et al. , 2021), and at the photosphere–chromosphere boundary (Marmolino et al. , 1993; Jovanović , 2014), with the parameter  $s = 0.6$ .

Gravity waves dissipate their energy contributing to local heating of the thermosphere at higher altitudes during extreme solar minimum, since the kinematic viscosity is much smaller in warmer than in colder thermosphere at the same altitude (Sindelarova, Buresova, and Chum , 2009). During extreme solar minimum the lower thermosphere is relatively cold,  $T \approx 500$  K, while during active solar conditions the temperature in thermosphere can be  $T \approx 2000$  K (Vadas , 2007). The current 25<sup>th</sup> solar cycle, which began in Decemeber 2019, is expected to have maximum activity in July 2025. This solar activity could increase the temeperature in the lower thermosphere several times. The reflection coefficient for acoustic waves in active solar conditions varies with the frequency  $\Omega$  and the parameter  $s = 250K/2000K = 0.125$ , as depicted in Fig. 7. The reflection coefficient decreases in the frequency range  $\Omega_{co} < \Omega < 1.5$ . Acoustic waves with  $V_h \geq 1/\sqrt{s} \approx 2.83$  are the best candidates to pass through D layer–lower thermosphere boundary and propagate further in the thermosphere. Acoustic waves with  $V_h \gg 1/\sqrt{s}$  are the most susceptible to reflection. This is opposite situation compared to the reflection coefficient for acoustic waves with  $s = 0.5$ , Fig. 5, where the waves with  $V_h \approx 1/\sqrt{s}$  are the ones that are most prone to reflection. For frequencies  $\Omega > 1.5$ , the



**Figure 7.** Reflection coefficient for acoustic waves at the D layer–lower thermosphere plane boundary  $z = 0$ , as a function of frequency  $\Omega$  and parameter  $s = 0.125$ .


reflection coefficient decreases very slowly and remains almost constant.

230 The reflection coefficient for gravity waves in active solar conditions as a function of frequency  $\Omega$  and the parameter  $s = 0.125$ , is shown in Fig. 8. It has very high values for all gravity waves propagating with allowed  $V_h < 0.9$ . These waves can hardly pass the boundary between the ionospheric D layer and the lower thermosphere. It seems that they are trapped in the ionospheric D layer and cannot propagate through the thermosphere. Therefore, infrasound could be instrument in coupling between ionospheric D layer and lower thermosphere during solar maximum activity.

235 The conditions for AGW propagation, as well as their reflection coefficient, strongly depend on the temperature through  $v_s$  and parameter  $s$ , Eqs. (8) and (17). Therefore, any change in temperature can affect the propagation of AGWs and their reflection and transmission features. This means that the detection of these waves depends on the current temperature in the region being observed. A similar situation exists with the detection of AGWs by lidar or any other instrument because their positions relative to the wave source region will determine which AGW characteristic can be observed (Yuan et al. , 2016).

240 One of the important effects of AGWs and especially gravity waves is their influence on the concentration of charged particles in the ionospheric E layer embedded in the lower thermosphere at an altitude of 90 – 140 km. Namely, the concentration of charged particles becomes time–dependent in the presence of waves. The changed characteristics of this layer affect the reflection of radio waves and telecommunication connections (Zawdie et al. , 2022). A similar situation is seen in strong natural hazards when earthquakes of magnitude  $M_v 5.5+$  are studied by VLF/LF radio waves. A physical interpretation is based on  
245 atmospheric gravity waves which could alter the ionospheric E layer and modulate the height of the VLF/LF waves reflection (Eichelberger et al. , 2024).

An interesting approach to the study of linear AGWs has been made by Cheremnykh (2020). He suggests that AGWs in an



**Figure 8.** Reflection coefficient for gravity waves at the D layer–lower thermosphere plane boundary  $z = 0$ , as a function of frequency  $\Omega$  and parameter  $s = 0.125$ .

isothermal atmosphere can be considered as a superposition of oscillations that occur simultaneously at two natural frequencies–acoustic and gravitational for a fixed wavelength.

## 250 6 Conclusions

In this article, analytical equations are used to study AGWs propagation through the ionospheric D layer and the D layer–lower thermosphere coupling. The dispersion equation and the reflection coefficient show that infrasound waves with frequencies  $\omega > 0.035s^{-1}$  that propagate almost vertically can reach the lower thermosphere. Gravity waves propagate in both regions–the ionospheric D layer and the lower thermosphere if their frequencies are  $\omega < 0.014s^{-1}$  and their horizontal phase velocities are  $v_h < 285m/s$ . Gravity waves with frequencies much lower than the Brunt–Väisälä frequency  $\omega_{BV} = 0.0195s^{-1}$  propagate more horizontally than vertically because  $\lambda_p \ll \lambda_z$ . These waves have lower vertical phase velocities than high–frequency gravity waves with  $\omega \approx \omega_{BV}$  which travel faster through the ionospheric D layer towards the lower thermosphere. The reflection coefficient is the smallest for the gravity waves with the frequencies  $\omega < 0.0087s^{-1}$ , horizontal phase velocities  $159m/s < v_h < 222m/s$ , and horizontal wavelengths  $115 \text{ km} < \lambda_p < 161 \text{ km}$ , which is in accordance with the results known in the scientific literature (Lizunov and Hayakawa, 2004; Fritts et al., 2014; Bakhmetieva et al., 2019). These waves can generate the middle–scale TIDs and cause temperature rise in the lower ionosphere.

The reflection coefficient is highly temperature dependent. It changes significantly during the pronounced solar maximum when the temperature in the lower thermosphere can rise several times. A strong increase in the reflection coefficient for gravity waves indicates that they cannot pass D layer–lower thermosphere boundary. Therefore, infrasound waves are better



265 coupling instruments.

There is broad scientific interest in the future study of AGWs. This is particularly attributed to the study of natural hazards, telecommunications and navigation, and space weather. Due to the complex nature of this process, differences between model results and observations are expected (Klymenko et al. , 2021).

270

*Author contributions.* There is only one author and the article is the result of the work of one author.

275 *Competing interests.* The author declares that they have no conflict of interest.

*Acknowledgements.* The research and writing of this work was supported by the Montenegrin National Project "Physics of Ionized Gases and Ionized Radiation".



## References

280

Afraimovich, E. L., Edemskiy I. K., Leonovich, A. S., Leonovich, L. A., Voeykov, S. V., and Yasyukevich, Y. V.: MHD nature of night-time MSTIDs excited by the solar terminator, *Geophys. Res. Lett.*, 36, L15106, doi:10.1029/2009GL039803, 2009.

Argunov, V. and Gotovtsev, M.: Researching of quasiperiodic variations in the amplitude of VLF electromagnetic signals from lightning discharges passing over strong earthquakes, *E3S Web of Conferences* 127, 03005, <https://doi.org/10.1051/e3sconf/201912703005>, 2019.

285 Bakhmetieva, N. V., Grigoriev, G. I., Tolmacheva, A. V., Zhemyakov, I. N.: Investigations of Atmospheric Waves in the Earth Lower Ionosphere by Means of the Method of the Creation of the Artificial Periodic Irregularities of the Ionospheric Plasma, *Atmosphere*, 10, 8, <https://www.mdpi.com/2073-4433/10/8/450>, 2019.

Balan, N., Alleyne, H., Walker, S., Reme, H., McCrea, I., and Aylward, A.: Magnetosphere-ionosphere coupling during the CME events of 07-12 November 2004, *Journal of Atmospheric and Solar-Terrestrial Physics*, 70, 2101, <https://doi.org/10.1016/j.jastp.2008.03.015>, 2008.

290 Blanc, E.: Observations in the upper atmosphere of infrasonic waves from natural or artificial sources: A summary, *Ann. Geophys.*, 3, 673 – 688, 1985.

Bochev, A. Z. and Dimitrova, I. I. A.: Magnetic cloud and magnetosphere-ionosphere response to the 6 November 1997 CME, *Advances in Space Research*, 32, 10, 1981-1987, [https://doi.org/10.1016/S0273-1177\(03\)90636-3](https://doi.org/10.1016/S0273-1177(03)90636-3), 2003.

Bothmer, V. and Daglis, I. A.: *Space weather: physics and effects*, Springer Science and Business Media, [http://dx.doi.org/10.1007/978-3-](http://dx.doi.org/10.1007/978-3-540-34578-7)

295

Boudjada, M. Y., Biagi, P. F., Eichelberger, H. U., Galopeau, P. H. M., Schwingenschuh, K., Solovieva, M., Nico, G., Lammer, H., Voller, W., and Stachel, M.: Study of VLF phase and amplitude variations before the Turkey Syria Mw 7.8 EQs, *EGU General Assembly 2024*, Vienna, Austria, 14–19 Apr 2024, EGU24-10341, <https://doi.org/10.5194/egusphere-egu24-10341>, 2024.

Cheremnykh, O. K., Kryuchkov, E. I., Fedorenko, A. K. et al.: Two-Frequency Propagation Mode of Acoustic-Gravity Waves in the Earth's

300

Atmosphere, *Kinemat. Phys. Celest. Bodies*, 36, 64–78, <https://doi.org/10.3103/S0884591320020026>, 2020.

Chum, J., Urbar, J., Laštovička, J., Carbera, M. A., Liu, J. Y., Bonomi, F. A. M., Fagre, M., Fišer, J., and Mošna, Z.: Continuous Doppler sounding of the ionosphere during solar flares, *Earth, Planets and Space*, 70, 198, doi:10.1186/s40623-018-0976-4, 2018.

Eichelberger, H., Boudjada, M. Y., Schwingenschuh, K., Besser, B. P., Wolbang, D., Solovieva, M., Biagi, P. F., Galopeau, P. H. M., Jaffer, G., Schiringer, C., Nina, A., Jovanovic, G., Nico, G., Stachel, M., Aydogar, Ö., Muck, C., Wilfinger, J., Jernej, I., and Magnes, W.:

305

Investigation of VLF/LF electric field variations related to magnitude  $M_w \geq 5.5$  earthquakes in the Mediterranean region for the year 2023, *EGU General Assembly 2024*, Vienna, Austria, 14–19 Apr 2024, EGU24-6001, <https://doi.org/10.5194/egusphere-egu24-6001>, 2024.

Fleck, B., Carlsson, M., Khomenko, E., Rempel, M., Steiner, O., and Vigeesh, G.: Acoustic-gravity wave propagation characteristics in three-dimensional radiation hydrodynamic simulations of the solar atmosphere, *Phil. Trans. R. Soc. A.*, 379, doi:10.1098/rsta.2020.0170, 2021.

310

Fritts, D. C., Pautet, P.-D., Bossert, K., Taylor, M. J., Williams, B. P., Limura, H., Yuan, T., Mitchell, N. J., and Stöber, G.: Quantifying Gravity Wave Momentum Fluxes with Mesosphere Temperature Mappers and Correlative Instrumentation, *J. Geophys. Res. Atmosphere*, 119, doi: 10.1002/2014JD022150, 2014.

315

Hayakawa, M., Kasahara, Y., Nakamura, T., Hobar, Y., Rozhnoi, A., Solovieva, M., and Molchanov, O. A.: On the correlation between ionospheric perturbations as detected by subionospheric VLF/LF signals and earthquakes as characterized by seismic intensity, *Journal of Atmospheric and Solar-Terrestrial Physics*, 72, 13, 982-987, <https://doi.org/10.1016/j.jastp.2010.05.009>, 2010.



- Inan, U. S., Lehtinen, N. G., Moore, R. C., Hurley, K., Boggs, S., Smith, D. M., and Fishman, G. J.: Massive disturbance of the daytime lower ionosphere by the giant  $\gamma$ -ray flare from magnetar SGR 1806-20, *Geophysical Research Letters*, 34, 8103, <https://doi.org/10.1029/2006GL029145>, 2007.
- Jovanović, G.: Reflection Properties of Gravitational-MHD Waves in an Inhomogeneous Horizontal Magnetic Field, *Sol. Phys.*, 289, 4085–4104, <https://doi.org/10.1007/s11207-014-0579-6>, 2014.
- Jovanovic, G.: Gravitational-acoustic wave reflection, *Romanian Reports in Physics*, 68, 2, 459-472, 2016.
- Klymenko, Y. O., Fedorenko, A. K., Kryuchkov, E. I. et al.: Identification of Acoustic-Gravity Waves According to the Satellite Measurement Data, *Kinemat. Phys. Celest. Bodies*, 37, 273–283, <https://doi.org/10.3103/S0884591321060052>, 2021.
- Lizunov, G. and Hayakawa, M.: Atmospheric Gravity Waves and their Role in the Lithosphere-troposphere-ionosphere Interaction, *IEEJ Transactions on Fundamentals and Materials*, 124, 12, 1109-1120, <https://doi.org/10.1541/ieejfms.124.1109>, 2004.
- Marmolino, C., Severino, G., Deubner, F.L., and Fleck, B.: Phases and Amplitudes of Acoustic-Gravity Waves II. The Effects of Reflection, *Astron. Astrophys.*, 278, 617-626, 1993.
- Mihalas, D.: *Foundations of Radiation Hydrodynamics*, Oxford University Press, Oxford, 1984.
- Mitra, A. P.: *Ionospheric effects of solar flares*, D. Reidel, Publishing Company, US, 1974.
- 320 Nenovski, P., Spassov, C., Pezzopane, M., Villante, U., Vellante, M., and Serafimova, M.: Ionospheric transients observed at mid-latitudes prior to earthquake activity in Central Italy, *Natural Hazards and Earth System Sciences*, 10, 1197, <https://doi.org/10.5194/nhess-10-1197-2010>, 2010.
- Nina, A., Biagi, P. F., Mitrović, S. T., Pulnits, S., Nico, G., Radovanović, M., and Popović, L. Č.: Reduction of the VLF Signal Phase Noise Before Earthquakes, *Atmosphere*. 2021; 12(4):444. <https://doi.org/10.3390/atmos12040444>, 2021.
- 335 Nina, A. and Čadež, V.: Detection of acoustic-gravity waves in lower ionosphere by VLF radio waves, *Geophysical Research Letters*, 40, 18, 4803-4807, <https://doi.org/10.1002/grl.50931>, 2013.
- Nina, A., Čadež, V., Popović, L., and Srećković V.: Diagnostics of plasma in the ionospheric D-region: detection and study of different ionospheric disturbance types, *The European Physical Journal, D* 71, 7, 1-12, <https://doi.org/10.1140/epjd/e2017-70747-0>, 2017.
- Rozhnoi, A., Shalimov, S., Solovieva, M., Levin, B., Hayakawa, M., and Walker, S.: Tsunami-induced phase and amplitude perturbations of subionospheric VLF signals, *Journal of Geophysical Research*, 117, A09313, doi: 10.1029/2012A017761, 2012.
- 340 Schulthess, G. B.: *Acoustic Waves in the Upper Atmosphere*, All Graduate Plan B and other Reports, 1625, <https://digitalcommons.usu.edu/gradreports/1625>, 2022.
- Sindelarova, T., Buresova, D., and Chum, J.: Observations of acoustic-gravity waves in the ionosphere generated by severe tropospheric weather, *Stud. Geophys. Geod.*, 53, 403-418, <https://doi.org/10.1007/s11200-009-0028-4>, 2009.
- 345 Singh Ashutosh, K., Singh, A. K., Singh R., and Singh, R. P.: Solar flare induced D-region ionospheric perturbations evaluated from VLF measurements, *Astrophys Space Sci*, 350, 1-9, [10.1007/s10509-013-1699-4](https://doi.org/10.1007/s10509-013-1699-4), 2014.
- Singh, Ashutosh K., Singh, R., Veenadhari, B., and Singh, A. K.: Response of low latitude D-region ionosphere to the total solar eclipse of 22 July 2009 deduced from ELF/VLF analysis, *Advances in Space Research*, 50, 10, 1352, <https://doi.org/10.1016/j.asr.2012.07.005>, 2012.
- 350 Vadas, S. L.: Horizontal and vertical propagation and dissipation of gravity waves in the thermosphere from lower atmospheric and thermospheric sources, *J. Geophys. Res.*, 112, A06305, doi:10.1029/2006JA011845, 2007.
- Voss, H. D., Walt, M., Imhof, W. L., Mobilia, J., and Inan, U. S.: Satellite observations of lightning-induced electron precipitation, *Journal of Geophysical Research*, 103, 11725, <https://doi.org/10.1029/97JA02878>, 1998.





- 355 Yuan, T., Heale, C. J., Snively, J. B., Cai, X., Pautet, P-D., Fish, C., Zhao, Y., Taylor, M. J., Pendleton, W. R., Wickwar, V., and Mitchell, N. J.: Evidence of dispersion and refraction of a spectrally broad gravity wave packet in the mesopause region observed by the Na lidar and Mesospheric Temperature Mapper above Logan, Utah, *Journal of Geophysical Research: Atmospheres*, 121, 2, <https://doi.org/10.1002/2015JD023685>, 2016.
- 360 Zawdie, K., Belehaki, A., Burleigh, M., Chou, M-Y., Dhadly, M. S., Greer, K., Halford, A. J., Hickey, D., Inchin, P., Kaeppler, S. R., Klenzing, J., Narayanan, V. L., Sassi, F., Sivakandan, M., Smith, J. M., Zabolin, N., Zettergren, M. D., and Zhang, S-R.: Impacts of acoustic and gravity waves on the ionosphere, *Front. Astron. Space Sci.* 9:1064152, doi: 10.3389/fspas.2022.1064152, 2022.